

DOI: 10.29289/259453942018V28S1032

PHARMACOKINETIC MODEL FOR QUANTITATIVE EVALUATION OF THE EPR EFFECT ON ALBUMIN-BASED NANOCARRIERS

Ailton A. Sousa-Junior¹, Gustavo C. P. Leite¹, Nicholas Zufelato¹, Franciyelli Mello-Andrade², Wanessa C. Pires², Elisângela P. Silveira-Lacerda², Andris F. Bakuzis¹

¹Instituto de Física, Universidade Federal de Goiás – Goiânia (GO), Brazil.

²Instituto de Ciências Biológicas, Universidade Federal de Goiás – Goiânia (GO), Brazil.

Understanding the dynamics of the tumor microenvironment is fundamental for the success of new anti-cancer therapies based on tumor targeting drug delivery. It is known that, in an effort to guarantee a continuous flow of oxygen and nutrients, the tumor cells emit molecular signals (VEGF, Vascular Endothelial Growth Factors) to the neighboring endothelial cells, inducing the formation of new blood vessels. It is also known that the endothelium of these new blood vessels shows abnormalities, like intercellular fenestrations, which allow an enhanced permeability and retention of nanostructures within the tumor extracellular matrix – an effect known as EPR (Enhanced Permeability and Retention). However, although this effect is known for over 30 years, very little is known about its kinetics. Therefore, we designed a 4-compartment pharmacokinetic model in order to quantitatively evaluate (both analytically and numerically) of the role of the EPR effect on the absorption of multifunctional nanocarriers by tumor tissues. Aliquots (150 μ L) of colloidal suspensions containing albumin-based nanocarriers were retro-orbitally administered to a control group (healthy Swiss mice) and to a treatment group (murine-Ehrlich tumor-bearing Swiss mice). Our nanocarriers, named after MalbIR, consisted of manganese ferrite (MnFe_2O_4) superparamagnetic nanoparticles associated both with molecules of bovine serum albumin (BSA) and with molecules of IR-780 iodide, a near infrared (NIR) fluorescent dye with chemotherapeutic action. The time-evolution of the amount of IR-780 (in pmol) within the retro-orbital, tail and tumor regions was determined via FMT (Fluorescence Molecular Tomography) during the first 8 days after the administration. According to the experimental data, in the first 48 h, the albumin-based nanocarriers accumulate within the tumor region presumably due to: the EPR effect (passive targeting); and to the affinity between BSA (and/or IR-780) molecules and tumor cell membrane receptors (active targeting). The results also enabled the determination of the pharmacokinetic model parameters (first-order intercompartment exchange rate constants, in h^{-1}), consequently bringing a better understanding of the EPR effect kinetics for this type of nanocarrier. Intratumoral accumulation (via EPR effect and presumably via other internalization mechanisms) was clearly observed. The time-evolution profile of the amount of IR-780 in different body regions (retro-orbital sinus, tail and tumor) could be successfully determined by FMT. Albumin-based nanocarriers have long blood half-lives (post-distributive phase), with no toxicity concerns at the assessed concentrations. In particular, our nanocarriers (MalbIR, i.e. MnFe_2O_4 nanoparticles + BSA + IR-780) show delivery efficiencies (DE) around 15% of the initial dose (ID); about 5 times higher than the DE measured for the free drug (free IR-780); and about 21 times higher than the median DE value reported by 232 papers over one decade (2005–2015).

DOI: 10.29289/259453942018V28S1033

THE RADIATION THERAPY IN OCCULT PRIMARY BREAST CANCER: CASE SERIES

Ana Flávia de Paula Guerra Campedelli¹, Nilceana Maya Aires Freitas¹, Thais Franco Simionatto¹, Nathalya Ala Yagi¹, Jean Teixeira de Paiva¹, Carolina Martinelli Bezerril¹

¹Hospital Araújo Jorge – Goiânia (GO), Brazil.

Objectives: Approximately 0.1–0.8% of all breast cancers is occult. The occult primary breast cancer (OPBC) is a breast cancer that first presents through regional nodal or distant disease without clinical or mammographic evidence of disease in the breast. Management of OPBC has been inconsistent and controversial since its initial description by Halsted in 1907, but the rarity of this form of breast cancer has made it difficult to standardize management. Furthermore, the natural history of OPBC remains unclear: most case series have reported outcomes that are better than those of non-occult breast cancers with similar nodal involvement, but a few have reported outcomes that are comparable or significantly worse. Here, we describe the management and outcomes of OPBC patients treated at a private and a public institution. Reviewing the management and outcome of OPBC patients we hope to elucidate both the benefits and disadvantages associated with a multidisciplinary treatment approach to OPBC. **Methods:** We described a four case series of women diagnosed with OPBC between 2015 and 2017 at Araujo Jorge Hospital and Cebrom Clinic, as well as the multimodal treatment received. All four patients were followed by a single observator. **Case Report:** The patients' ages ranged from 45 to 75 years, with a mean age of 60 years. The four patients presented with a palpable mass at the axillar area and all of them underwent level I and II axillary lymph node dissection pre radiotherapy. The pathologic results are invasive ductal carcinoma; 3 patients are grade 3 and 1 patient is grade 2; the imuno histologic results are 2 patients were luminous and 2 patients were triple negative; microscopic features revealed metastatic breast carcinoma in all of the four cases. All patients were submitted to adjuvant chemotherapy and then.

DOI: 10.29289/259453942018V28S1034

IR-780 BASED-NANOPARTICLE CYTOTOXIC EFFECT ON MURINE BREAST CANCER CELLS (EHRlich)

André Luiz Silva Oliveira^{1,2*}, Nicholas Zufelato³, Letícia Reis Silva², Andris Figueiroa Bakuzis³
Elisângela de Paula Silveira-Lacerda²

¹Bolsista FAPEG, Programa de Pós-Graduação em Ciências Biológicas, Instituto de Ciências Biológicas(ICB), Universidade Federal de Goiás (UFG) – Goiânia (GO), Brazil.

²Laboratório de Genética Molecular e Citogenética, ICB/UFG – Goiânia (GO), Brazil.

³Laboratório Ressonância Ferromagnética e Nanomagnetismo, IF/UFG – Goiânia (GO), Brazil.

*Corresponding author: andreluiz@ufg.br

Objectives: The aim of this study was to evaluate the cells viability of the murine breast cancer cells (Ehrlich) after treatment with magnetic nanocarriers (MALBIR). **Methodology:** Ehrlich's ascites carcinoma cells were aspirated from the peritoneal cavity of the animals and washed with 1x PBS solution, and then maintained in a humidified atmosphere for 24h at 37°C in 5% CO₂. RPMI-1640 medium was supplemented with 10% FBS, 1% penicillin/streptomycin and 0.3% amphotericin. The MALBIR was used as treatment for 24h, which consist of ferrite-based magnetic nanoparticles, bovine albumin proteins and IR-780 iodide molecules, forming an aggregate nanostructure with a diameter around 100nm. Because IR-780 is cytotoxic, all concentration used (0.0; 0.437; 0.875; 1.75; 3.5; 7; 14) were based in quantity of the IR-780 (µg.mL⁻¹) present in nanocarrier. The MALBIR cytotoxic effects were evaluated using the MTT assay with Ehrlich tumor cells as described by Mosman (1983). The results as presented as mean ± SD obtained from three independent experiments each done in four replicates. Statistical significance was considered at p<0.05 performed by one-way ANOVA with Tukey post test. The IC50 (IR-780 concentration in nanocarrier that results in a 50% reduction in cellular viability) was acquired by dose-response curves (nonlinear regression curves from sigmoidal dose-response) using GraphPad Prism 5 for Windows. **Results:** The IR-780 present in MALBIR has a statistical significantly effect inducing cell death in a dose dependent manner on Ehrlich cells. At concentrations of 3.5, 7 and 14µg.mL⁻¹ obtained a significantly different result from negative control with p<0.001. By nonlinear regression curve was estimated an IC50 value on Ehrlich cells of 14.2 ± 1.7µg.mL⁻¹. Similar results had previously been found for Sarcoma-180 tumor cell line and L-929 normal cell line treated with this same nanocarrier. **Conclusions:** In summary, MALBIR treatment has significant cytotoxic activity on Ehrlich cells line when compared to the reference negative control. Thus, these results suggest that the MALBIR treatment may share a distinct and important dose-dependent biological property. Additional studies are needed to compare normal cell line, determine the molecular mechanisms and to evaluate the potential *in vivo* anticancer activity of the MALBIR. Nevertheless, the nanocarrier shows potential for synergetic effect between the chemotherapeutic agent IR-780 and the nanoparticles, if magnetic hyperthermia treatment is also performed, that could decrease even further the IC50 of the nanoparticle.

DOI: 10.29289/259453942018V28S1035

TIME BETWEEN DIAGNOSIS AND BEGIN OF THE TREATMENT IN A PUBLIC BRAZILIAN HOSPITAL IN SÃO PAULO

André Mattar¹, Jorge Y. Shida¹, Roberto Hegg¹, Felipe A. Cavagna¹, Ana Luiza A. Faria¹, Luiz H. Gebrim¹

¹Women's Reference Center, Hospital Pérola Byington – São Paulo (SP), Brazil.

Objectives: Breast cancer affects about 59,000 Brazilian women annually according to data from INCA 2018 (National Health Institute) and almost 55% of patients are diagnosed in advanced stages. In most Brazilian cancer centers the treatment is made only with the definitive diagnosis (biopsy). It can lead to a delay of 5 to 8 months and may even take longer. On November 22, 2012, a federal law established in 60 days the maximum period between the diagnosis and the first treatment of any malignant disease. Pérola Byington Hospital is one of the most important cancer centers specialized Breast Cancer treatment in Brazil. In 2005 a model of visits based on quick diagnosis and treatment was implemented named CARE (Centro de Alta Resolutividade – High Resolutivity Center). This model is based in “One Stop Clinic” approach contemplates the integration between the professionals and the physical space in the same physical area. In this model a trained physician can have the results in the same day of mammography and ultrasonography. Biopsy is performed in the same day and after 2 or 3 weeks our pathologists trained in mammary pathology can give us the results with the immunohistochemistry analyses. With this approach, we have been able to optimize resources and gain the agility to diagnose breast cancer suspects, whether palpable or not. Many breast cancer cases require primary (neoadjuvant) treatments since many diagnoses were already made in late phases. In 2016, there was a need for integration with the clinical oncology service, allowing a clinical visit by the oncologist in the same day and the possibility to initiate the treatment between 7 to 10 days. This model based on a single visit requires an integrated and trained multidisciplinary team, not only in the treatment of the disease but also training in this model of approach. The idea of this model was born due to the observation that patients in Brazil usually spent a long time in lines to the first visit and had to wait long time to perform the specific exams (mammography/ultrasound) and then a long time to have an evaluation with the specialist and more time spend to undergo invasive diagnostic procedures. Usually patients had to return 3 or 4 times to have their diagnosis made and this process could take 2 or 3 months to start the treatment. With our approach we have been able to reduce the waiting time substantially, since all the tests are made on the same day (including biopsy), anticipating the diagnosis and only for this reason reducing the number of advanced cases. Since the prognosis is closely linked to the stage, this approach can substantially increase disease cure rates. Our Center annually treats about a thousand new cases of breast cancer and about 1400 surgical procedures are required. We have been spreading this idea promoting workshops, continued medical education; training courses in mammography and ultrasound diagnosis and medical training in percutaneous biopsies, both guided by ultrasonography and stereotactic. Our mission is to promote humanized treatment and achieve early diagnosis and timely treatment to achieve a better prognosis. **Methodology:** This study was performed at the Pérola Byington Hospital evaluating patients treated at our hospital from 2011 to 2017. We have reviewed the cases diagnosed with breast cancer at the hospital and the date of diagnosis and the beginning of treatment were evaluated, being it surgical or performed chemotherapy. **Results:** In this period 6,488 cases of breast cancer were included in our evaluation. The mean between the initial histological or cytologic diagnosis and the beginning of the treatment was calculated in each year: In 2011 63.82% of our cases had their first treatment beyond 60 days. In 2017 most of our patients were treated in at least 60 days (67.23%). We know that the mean can be very variable if one of the numbers is too large or too small and because of that we have been able to median between the diagnosis and the first treatment. In 2011 the median was 76, in 2012 was 83 and this number was going down year by year and in 2017 the median is 53 days. **Conclusions:** In our Hospital the “One Stop Clinic” approach have changed the way of treating the patients allowing more agility in the treatment of breast cancer including quick diagnosis and initiation of treatment as soon as possible.

DOI: 10.29289/259453942018V28S1036

THE VALUE OF STEREOTACTIC VACUUM-ASSISTED BIOPSY IN THE INVESTIGATION OF MICROCALCIFICATIONS IN 1354 PATIENTS IN PUBLIC BRAZILIAN HOSPITAL

Andressa G. Amorim¹, Marcellus N. M. Ramos¹, André Mattar¹, Jorge Y. Shida¹, Luiz H. Gebrim¹¹Women's Reference Center, Hospital Pérola Byington – São Paulo (SP), Brazil.

Objectives: The gold standard for breast biopsy procedures is currently an open excision of the suspected lesion. However, an excisional biopsy inevitably makes a scar. The cost and morbidity associated with this procedure has prompted many physicians to evaluate less invasive, alternative procedures. More recently, image-guided percutaneous core-needle biopsy has become a frequently used method for diagnosing palpable and non-palpable breast lesions. Although sensitivity rates for core-needle biopsy are high, it has the disadvantage of histological underestimation, which renders the management of atypical ductal hyperplasia, papillary lesions, and fibroepithelial lesions somewhat difficult. Stereotactic vacuum assisted breast biopsy (VABB) was developed to overcome some of these negative aspects of core-needle biopsy. VABB allows for a sufficient specimen to be obtained with a single insertion and can provide a more accurate diagnosis and completely remove the lesion under real-time ultrasonic guidance. The advantage of complete lesion removal with VABB is to reduce or eliminate sampling error, to decrease the likelihood of a histological underestimation, to decrease imaging-histological discordance, to decrease the re-biopsy rate, and to diminish the likelihood of subsequent growth on follow-up, especially when stereotactic VABB is used to investigate microcalcifications. This method is expensive but cost effective when used to investigate microcalcifications classified as *BI-RADS* 4 and 5. **Methodology:** We performed a review in 1,354 patients with suspicious mammography microcalcifications, classified as *BI-RADS* 4 or 5 that were seen in Perola Byington Hospital from July 2012 to July 2017 in São Paulo-Brazil. We have used a Hologic Lorad Multicare Platinum Stereotactic Prone Breast Biopsy and a Surus Pearl (Hologic, Malbolrough, Massachusetts, USA), with *gauge* 9 probe. Four to eight fragments (median of 6) were obtained with the placement of a metal clip in the bed that the biopsy was performed, and histopathological analysis was made. **Results:** The histopathological study of the lesions revealed benign alterations in 956 (68%) of our patients. The findings were positive for malignancy in 358 patients (29%) and the precursor lesions were diagnosed in 40 (3%). In 81 cases (5.9%) the findings were discordant. The sensitivity of the method was 84.4%, specificity was 96.1%, false negative rate was 4.5%, positive predictive value (PPV) was 89.8%, negative predictive value (NPV) was 93.8%. In literature review the sensitivity varies 91.5–100%, specificity 81.9–110%, false negative rate 0–3.3%, PPV 92.2–100% and NPV 80.5–99.5%. All patients with positive or discordant cases underwent surgical treatment to increase the margin or complete removal of the lesion. **Conclusions:** The VABB is an outpatient procedure that avoids hospital admissions for diagnostic elucidation in most of cases suspected of malignancy. It has high predictive value in both benign and malignant lesions, guiding therapeutic planning. In addition to presenting the cost lower than the surgical biopsy it indirectly increases the supply of hospital beds for cancer treatment.

DOI: 10.29289/259453942018V28S1037

PYODERMA GANGRENOSUM: REPORT OF A CASE INVOLVING THE BREAST

Andressa G. Amorim¹, Marcellus N. M. Ramos¹, Felipe A. Cavagna¹, Maria I. B. A. C. Sawada¹, Alexandre S. Melitto¹, André Mattar¹, Jorge Y. Shida¹, Luiz H. Gebrim¹

¹Women's Reference Center, Hospital Pérola Byington – São Paulo (SP), Brazil.

Background: Pyoderma gangrenosum (PG) is an inflammatory, ulcerative skin disorder characterized at histopathology by the presence of neutrophils in the skin. The disease is a rare but serious ulcerating skin disease, the treatment of which is mostly empirical. The condition may either develop spontaneously or in association with surgical trauma or systemic or neoplastic diseases. Pyoderma can present to a variety of health professionals and several variants exist that may not be recognized immediately. This can delay the diagnosis and have serious clinical consequences. Clinical presentation varies, but is basically characterized by ulcerated, painful, dark red or purple skin lesions with irregular, inflamed and raised borders and a necrotic base, with single or multiple small abscesses. The mainstay of treatment is long term immunosuppression, often with high doses of corticosteroids or low doses of ciclosporin. Recently, good outcomes have been reported for treatments based on anti-tumour necrosis factor α , and infliximab proved effective in a randomized controlled trial.

Case report: This report refers to a female patient who was 54 years old and that was seen in Perola Byington Hospital. She presented with spontaneous pyoderma gangrenosum and the lesions were seen on her right breast, left axilla, left eyelid and chin. She was submitted to a skin biopsy in 07/06/2017: chronic non-specific inflammatory disease. After the clinical and pathologic result, the hypothesis of PG was made, and she received systemic corticosteroids with prednisone 60 mg once a day orally for 30 days and after that more 90 days with prednisone 40mg. She had responded very well, and the lesions disappeared after 4 months. **Discussion:** The clinical course of pyoderma gangrenosum can be unpredictable and highly variable from onset and throughout progression of the condition; hence diagnosis of this pathology may constitute a challenge. Many patients develop painful lesions that grow progressively, accompanied by fever. In others, lesions are chronic, with ulcerations that progress slowly. Pathogenesis remains to be fully clarified. Since diagnosis is reached exclusively on the basis of clinical characteristics, it is very important to take this disease into consideration when evaluating differential diagnoses.

DOI: 10.29289/259453942018V28S1038

STAGE AND IMMUNOHISTOCHEMISTRY OF EIGHT THOUSAND CASES OF BREAST CANCER TREATED IN A BRAZILIAN REFERENCE CENTRE

Luiz Henrique Gebrim¹, André Mattar¹, Jorge Yoshinori Shida¹, Roberto Hegg¹¹Women's Reference Center, Hospital Pérola Byington – São Paulo (SP), Brazil.

Objetives: Breast cancer affects about 59,000 Brazilian women annually according to data from INCA 2018 (National Cancer Institute) and 14,388 deaths are related to the disease. There are few data about the stage of the disease in our Country. Most of the cases are confirmed in late stages mainly due to the delay in diagnosis and time to initiate the treatment. Pérola Byington Hospital is one of the most important cancer centers specialized breast cancer treatment in Brazil and we are responsible for treating about a thousand new cases of breast cancer annually. In 2005 a model of visits based on quick diagnosis and treatment was implemented named CARE (Centro de Alta Resolutividade – High Resolutivity Center) based on a model of One Stop Clinic. This approach can lead to a downstage of breast cancer stage once the diagnosis and the treatment is substantially quicker. In this study we have evaluated the stage and immunohistochemistry (IHC) subtype of the patients treated in our institution. **Methodology:** In 2016 we have created a web-based system that helped us to input information about the breast cancer cases treated in Pérola Byington Hospital. This program helped us to better understand our cases of breast cancer regarding the initial stage and the subtype of breast cancer based on IHQ. The main idea was to analyze our data and compare it with the national data and the international literature. **Results:** We have reviewed 8,420 cases of breast cancer treated in our Hospital between 2011 and 2017. We were able to access the clinical stage in 8,096 cases. The distribution was 4.9% of stage 0, 22.9% of stage I, 39.2% of stage II, 30% of stage III and 3.1% of stage IV at the time of diagnosis. The data regarding the immunohistochemistry subtype was able to be recovered in 7,934 patients. Most of our cases were luminal (A or B) – 56.4%. There was 10.3% of both Her2 and hormonal receptor positive and 7.9% of only Her2 positive. The prevalence in our patients of triple negative disease was 25.4% much higher than the literature (around 15%). We have also looked at the Her2 doubtful (Her2 2+) where usually you need a specific test to define it (CISH or FISH test). There were 1,450 cases in this condition and in 28.6% of them the test was not made. Among the cases that made the test we found that 74.7% resulted in CISH negative, 14.1% positive and 11.2% were still inconclusive. The Ministry of Health forces all the Her2 positive in the IHC (3/3) to be tested with a FISH or a CISH test. We found 616 cases of Her2 positive in the IHQ test and negative when CISH was made (7.2%). **Conclusions:** In our analysis we have found that our institution has more cases of triple negative disease than reported in the literature. Based on national statistics we had substantially fewer cases of advanced disease (only 30% of stage III compared to 60% of INCA report). Probably these findings are related to the quick diagnosis and immediate treatment (surgical or neoadjuvant). There was 17.4% of cases found to have Her2 uncertain and most of them were tested and became a Her2 negative when CISH was performed (74.7%). We have to look carefully to those patients with Her2 positive in IHC but negative when CISH was done specially because they do not receive trastuzumab according to Brazilian Ministry of Health guidelines a further study will be made to evaluate that.

DOI: 10.29289/259453942018V28S1039

ANALYSIS OF 41 PATIENTS WHO UNDERWENT ADENOMASTECTOMY AFTER A BREAST CANCER DIAGNOSIS IN A PUBLIC BRAZILIAN HOSPITAL

Andressa G. Amorim¹, Marcellus N. M. Ramos¹, André Mattar¹, Jorge Y. Shida¹, Luiz H. Gebrim¹¹Women's Reference Center, Hospital Pérola Byington – São Paulo (SP), Brazil.

Objectives: to describe the indications and complications of 41 patients with breast cancer submitted to skin-sparing mastectomy with or without preservation of the areola-papillary complex at a Public Hospital in Brazil. **Methods:** We've performed a retrospective analysis of 41 patients diagnosed with breast cancer that were treated from January 2008 to May 2015 and had adenomastectomy at Pérola Byington Hospital. Description data is shown. **Results:** The age of the patients ranged from 28 to 62 years, with a mean of 46 years. Invasive ductal carcinoma was diagnosed in 24 cases (58.4%), the invasive lobular type in 3 cases (7.3%). The other 14 (34.1%) cases were patients with multicentric ductal carcinoma *in situ*. In the invasive tumors there was predominance of hormone receptor positive (81.4%). In the preoperative clinical staging predominant was TisN0M0 in 31.7% cases. The preservation of areola-papillary complex occurred in 28 cases (68.2%). The sentinel lymph node biopsy it was made in 70.3% cases the invasive tumors. Regarding immediate repair reconstruction, the expander prosthesis was used in 38 cases (92.6%). Among the complications, there were no reports of bruising or infection. Extrusion of the expander prosthesis occurred in only 2 cases (4.8%). After a mean follow-up of 48 months, no local recurrences were reported. **Conclusions:** Skin sparing mastectomy with or without preservation of the areola-papillary complex is a safe option in the oncologic treatment of *in situ* and invasive cancer (T1 and T2), especially in multicentric forms in patients with small breasts. Immediate reconstruction was possible in all patients. The main complication observed was the late extrusion of the prosthesis in 4.8% of cases.

DOI: 10.29289/259453942018V28S1040

BIOMARKER EXPRESSION IN BREAST CANCER. CORRELATIONS WITH OUTCOME FOLLOWING NEOADJUVANT CHEMOTHERAPY

Augusto Ribeiro Gabriel¹, Ruffo Freitas-Junior¹¹CORA - Centro Avançado de Diagnóstico da Mama, Universidade Federal de Goiás, Rede Brasileira de Pesquisa em Mastologia – Goiânia (GO), Brazil.

Purpose: Breast cancer is the most common type of cancer in women in Brazil and worldwide, corresponding to around 20% of all malignant tumors in women. Neoadjuvant chemotherapy improves survival rates in around 30% of patients with locally advanced disease. This study evaluated biomarker expression in tumor samples and assessed outcome variables following neoadjuvant chemotherapy. **Methods:** This longitudinal, analytical study included 86 patients of 28 to 80 years of age who were evaluated according to clinical data, histology and laboratory analysis of selected markers. **Results:** Positivity for estrogen (ER) and progesterone receptors (PR) was 58.54% and 65.79%, respectively. The tumors were predominantly negative for the HER2, CK5/6, EGFR, p53 and Topo II biomarkers and predominantly positive for the Ki-67, PTEN and IGF-1 biomarkers. The results obtained for ER showed associations with PR ($p=0.002$), Ki-67 ($p=0.049$), CK5/6 ($p=0.027$) and p53 ($p=0.032$). Likewise, there were associations between HER2 and Ki-67 ($p=0.028$) and between Ki-67 and Topo II ($p=0.001$). Eighteen patients (20.93%) achieved complete clinical response, while 9 (10.47%) achieved pathologic complete response (pCR). There was an association between pCR and the HER2 ($p=0.05$) and Topo II ($p=0.007$) biomarkers. More patients with HER2 tumors (40%) achieved pCR compared to those with the other subtypes, while none of the patients with luminal A tumors achieved pCR ($p=0.043$). **Conclusions:** In this study, HER2 and Topo II expression, as well as tumors classified as HER2 subtype or luminal A subtype were predictive of pCR.

DOI: 10.29289/259453942018V28S1041

OBESITY INCREASES THE RISK OF BREAST CANCER IN POSTMENOPAUSAL WOMEN?

Carolina R. Mendonça¹, Magno B. Cirqueira¹, Jalsi T. Arruda², Nilza N. Guimarães³, Erika A. Silveira¹, Waldemar N. do Amaral¹

¹Programa de Pós-Graduação em Ciências da Saúde, Universidade Federal de Goiás – Goiânia (GO), Brazil.

²Faculdade de Medicina, Centro Universitário de Anápolis (UniEVANGÉLICA) – Anápolis (GO), Brazil.

³Departamento de Morfologia, Instituto de Ciências Biológicas, Universidade Federal de Goiás – Goiânia (GO), Brazil.

Objective: To describe the impact of obesity on the increased risk of breast cancer in postmenopausal women.

Methodology: The search for scientific articles was carried out in the databases: PubMed and Scopus. Articles published in English in the last five years have been selected. The surveys were conducted with the following combinations: (“Obesity”[MeSH Terms] AND “Breast Neoplasms”[MeSH Terms] AND “postmenopausal women”[All Fields]). Criteria for definition of obesity was the Body Mass Index (BMI) ≥ 30 . **Results:** We included 25 studies. Most epidemiological studies report that women with obesity and overweight are at higher risk of developing postmenopausal breast cancer. BMI ≥ 30 , at the time of diagnosis, is associated with more advanced stages of breast cancer with more aggressive tumor characteristics, larger tumor size and worse histological grade. An increase of 5 units in BMI is associated with a 12% increased risk of breast cancer. Among postmenopausal women, obesity has a 20% to 40% increased risk of developing breast cancer compared to women of normal weight. The greatest risks are observed mainly in women who have never used menopausal hormone therapy. Weight gain after diagnosis indicated an increase in mortality rates and obesity would be linked to an increased risk of secondary neoplasms. Obese women compared to normal weight women had a 21–41% higher overall mortality risk and a 25–68% risk of breast cancer mortality, depending on how much time had elapsed since the diagnosis of the disease. Another study suggests that increasing body fat percentage 6 months after breast cancer surgery is an important predictor of post-menopausal distant metastases. In another study, high BMI was protective in premenopausal women (OR=0.14 per; 95%CI 0.02–0.77), while in postmenopausal women it was a risk factor for development of breast cancer (OR=2.39 per; 95%CI 1.02–5.55). One explanation for the increased risk is that obesity leads to the release of cytokines into the bloodstream that affect the metabolism of breast cancer cells, making them more aggressive. Further, the impact of obesity on risk differs by tumor hormone receptors status and, among postmenopausal women, use of hormone replacement therapy. **Conclusion:** Obesity is associated with an increased risk of postmenopausal breast cancer.

DOI: 10.29289/259453942018V28S1042

BRCA1 AND MICRORNAS 7, 10B, 205AB, 218A EXPRESSION AS PROGNOSTIC MARKERS IN PRIMARY BREAST CANCERS – A RETROSPECTIVE COHORT STUDY

Cesar A. S. T. Vilanova-Costa^{1,2}, Jéssica E. P. Ramos³, Juliana F. Paes⁴, Daniel R. Bastos¹, Nathália A. Nogueira^{1,4}, Sílvia H. Rabelo-Santos⁶, Raphael B. Parmigiani⁷, Vera A. Saddi^{1,3,4,5}

¹Laboratório de Genética e Biodiversidade, Programa de Pós-graduação em Ciências Ambientais e Saúde, Pontifícia Universidade Católica de Goiás – Goiânia (GO), Brazil.

²Laboratório de Biologia Tumoral, Hospital Araújo Jorge, Associação de Combate ao Câncer em Goiás – Goiânia (GO), Brazil.

³Escola de Ciências Médicas, Farmacêuticas e Biomédicas, Pontifícia Universidade Católica de Goiás – Goiânia (GO), Brazil.

⁴Programa de Pós-graduação em Ciências da Saúde, Universidade Federal de Goiás – Goiânia (GO), Brazil.

⁵Laboratório de Oncogenética e Radiobiologia, Instituto de Ensino e Pesquisa, Associação de Combate ao Câncer em Goiás – Goiânia (GO), Brazil.

⁶Instituto de Patologia Tropical, Universidade Federal de Goiás – Goiânia (GO), Brazil.

⁷Idogene Medicina Diagnóstica – São Paulo (SP), Brazil.

Micro-RNAs (miRs) are post-transcriptional regulators of gene expression involved in several important biological processes. BRCA1 is a tumor suppressor gene and BRCA1-silent breast cancers (BC) tend to be more aggressive. Since BRCA1 may be regulated at post-transcriptional level by miRNAs, the purpose of this study was to evaluate the prognostic value of human miR-7, miR-10b, miR-205ab and miR-218b and BRCA1 expression levels in BC. A set of 36 triple-negative (TN) and 56 nontriple-negative (NTN) breast tumors was analyzed. Total miRNA was extracted from formalin-fixed paraffin-embedded (FFPE) BCs collected from the Pathology Department of Araújo Jorge Hospital-ACCG (Goiânia, Goiás, Brazil). MiRs expression was quantified by Quantitative Real-Time PCR (qRT-PCR) and BRCA1 expression was evaluated by immunohistochemistry (IHC). The present study was approved by the institutional Ethics Committee of Araújo Jorge Hospital (Report n° 948.930, 2015). The relative expression levels of miRs and clinic pathological features of breast cancers were compared. Overall survival in 60 months was 72.8%, and it was influenced by TNBC phenotype ($p=0.044$), tumor size ($p=0.007$), lymph node involvement ($p=0.038$), distant metastasis ($p=0.0008$), BRCA1 negative expression ($p=0.039$), miR-7 ($p=0.026$) and miR-10b ($p=0.011$) overexpression. MicroRNA hsa-miR-7 overexpression was associated with larger tumors (>2 cm) ($p=0.041$), higher histological grade ($p=0.028$), TN phenotype ($p=0.012$), BRCA1-negative expression ($p=0.047$) and worse survival ($p=0.026$). Overexpression of hsa-miR-10b was associated with larger tumors ($p=0.047$), lymph node ($p=0.032$) and distant metastases ($p=0.019$), higher histological grade ($p=0.009$), TN phenotype ($p=0.027$), BRCA1-negative expression ($p=0.006$) and worse survival ($p=0.011$). Meanwhile, underexpression of hsa-miR-205ab was associated with larger tumors ($p=0.027$), lymph node ($p=0.046$) and distant metastases ($p=0.014$), BRCA1-negative expression ($p=0.027$), TN phenotype ($p=0.038$) and worse survival ($p=0.024$). While, hsa-miR-218a underexpression was associated with a larger tumor size ($p=0.032$), lymph node ($p=0.011$) and distant metastases ($p=0.022$), TN phenotype ($p=0.019$), negative expression of BRCA1 ($p=0.039$) and worse survival ($p=0.003$). Our results show that BRCA1 protein expression assessment, miR-7 and miR-10b overexpression and miR-205ab and miR-218b underexpression could be useful in evaluating BCs prognosis, especially for patients with triple negative tumors.

DOI: 10.29289/259453942018V28S1043

PROSTHESIS EXPOSURE MANAGEMENT AFTER IMMEDIATE BREAST RECONSTRUCTION

Cícero de Andrade Urban¹, Rafael Amin Menezes Hassan¹, Rubens Silveira de Lima²¹Universidade Positivo – Curitiba (PR), Brazil.²Departamento de mastologia, Hospital Nossa Senhora das Graças – Curitiba (PR), Brazil.

Introduction: Breast cancer is the most frequent malignant neoplasm among women. When diagnosed at an early stage, it has a good prognosis. With the biotechnological development of the prostheses and the advent of oncoplastics, the surgeries started to present a better cosmetic result, bringing with it psychological and functional benefits, without compromising the oncological character of the treatment. Breast reconstruction with the use of prostheses is far from achieving low complication rates, making studies of a great relevance that can help to define and improve the behavior in cases of complicating prosthesis surgeries. The objective of the study was to evaluate the flowchart adopted by a single team of breast surgery for guidance of therapeutic management in patients submitted to breast reconstruction who evolved with prosthesis exposure. **Methods:** Retrospective analysis of the database included reconstructed patients with use of definitive or temporary breast implants between 2014 and June 2016. The epidemiological characteristics of the population were evaluated; type of surgery and prosthesis mark; postoperative follow-up; and application of the conduit flow chart, which includes history of radiotherapy, presence of severe infection and extensive necrosis. **Results:** There were 56 complications (20.2%), 33 cases were exposed prostheses and had criteria for application of the conduit protocol, of which 14 had criteria for simple suture, with a success rate of 57.1%. At the final outcome, nine had loss of breast reconstruction and 24 remained with reconstructed breasts. **Conclusions:** The protocol for ducts for exposed prostheses, evaluated in this study, showed results with acceptable efficacy and good reproducibility, facilitating its application in future studies.

DOI: 10.29289/259453942018V28S1044

AN TYROSINE KINASE RECEPTOR AS PROMISING THERAPEUTIC TARGET FOR TRIPLE NEGATIVE BREAST CANCER

Joana D. R. Santos¹, Maria F. S. Cardoso¹, Carlos H. M. Castelletti¹, Nancy C. F. L. Ferreira^{1,3}, Darley L. F. Filho³, José L. L. Filho^{1,2}, Danyelly B. G. Martins^{1,2}

¹Setor de Prospecção Molecular, Laboratório de Imunopatologia Keizo Asami, Universidade Federal de Pernambuco – Recife (PE), Brazil.

²Departamento de Bioquímica, Centro de Biociências, UFPE – Recife (PE), Brazil.

³Hospital Barão de Lucena – Recife (PE), Brazil.

Financial support: CAPES, FACEPE, CNPq

Objective: The aim of this study was to evaluate the expression of PDGFRA (platelet-derived growth factor receptor alpha) in 8 breast cancer tissue samples of which 2 TNBC (triple negative breast cancer), 3 luminal B and 3 luminal A. Clinical, and staging data were considered. **Methods:** The RNA was purified with the *RNeasy Mini Kit* (Qiagen), the cDNA was prepared by RT-PCR with the *QuantiTect Reverse Transcription* (Qiagen). The analysis was performed by qPCR (quantitative polymerase chain reaction) in the real-time PCR systems StepOnePlus™ (Applied Biosystems) with the GoTaq® qPCR Master Mix kit (Promega). β -actin was used as the reference gene to normalize the expression of PDGFRA with specific primers. **Results:** In TNBC patients, show higher gene expression of PDGFRA 2.36 times more than luminal patients. In addition, the PDGFRA overexpression occur even after neoadjuvant chemotherapy. Is associated with. It has also been demonstrated that african ethnicity and use of hormonal contraceptives may be related to the activation of PDGFRA. Both TNBC patients this study were recurrent in breast cancer. TNBC is an aggressive form of breast cancer that differs in epidemiology, risk factors, and prognosis from other types of breast cancer. The challenge in treating TNBC lies in the lack of targeted therapies currently available. Chemotherapy remains the only treatment option presently. Like many receptor tyrosine kinases (RTKs), PDGFRA is involved in the progression of a variety of cancers either by overexpression or by increased activity. **Conclusion:** These findings indicate that PDGFRA as a potential biomarker as well as a therapeutic target for TBNC.

DOI: 10.29289/259453942018V28S1045

EVALUATION OF THE FALSE NEGATIVE RATE IN THE EXAMINATION OF FREEZING OF THE SENTINEL LYMPH NODE IN HBL – RECIFE (PE)

Ana Flavia Morais Leda¹, Darley de Lima Ferreira Filho¹, Nancy Cristina Ferraz de Lucena Ferreira¹¹Hospital Barão de Lucena – Recife (PE) – Brazil.

Introduction: Breast cancer is the most commonly diagnosed cancer in women in the United States and accounts for 26% of all female cancers. In 2015, approximately 40,290 US women died from breast cancer. In our country, an estimated 59,700 new cases of invasive breast cancer. Then, is very important the early diagnosed of breast cancer and the use of sentinel lymph node probably chance this prognosis of this disease. **Objective:** To evaluate the false-negative rate of freezing of sentinel lymph node In breast cancer and the factors associated with it. **Methods:** Twenty-five breast cancer patients who underwent patent blue injection and sentinel lymph node (LS) biopsy between March 2016 and March 2017 in 570 surgery were evaluated for the following variables: age, tumor size, histological characteristics of the tumor (type- and degree of differentiation) and diagnosis of the definitive freezing and histopathological examination of LS. **Results:** The average age was 56,2, the tumor size was 2,3 cm and the histological type was carcinoma ductal invasive in 80%. In a relation the degree of diferenciacion, the G2 appear in 19 patients, G1 in 2 cases and G3 in 3 cases. The false-negative rate in the LS freezing test was 8% (2 cases). The specificity was 100%, and sensitivity was 50%. None of the variables evaluated was associated with increase in the incidence of false-negative. In our research, there was no association between tumor size and incidence of axillary metastasis. **Conclusion:** The false-negative rate of the LS freezing test was 8% and none of the variables evaluated correlated with this event. No anatomopathological variable studied was associated with the occurrence of false negative

DOI: 10.29289/259453942018V28S1046

BREAST CANCER IN AN 18-YEAR-OLD PATIENT: A CASE REPORT

Maria Letícia Moraes¹, Eduardo Resende Sousa e Silva¹, Laura Viana de Lima¹, Carla Paulinelli Seba¹,
Isabela Vieira Bastos¹, Ana Paula Teixeira Leite¹, João Emanuel Meireles Gonçalves¹, Renata Pereira Fontoura¹

¹Brasília (DF) – Brazil.

Objective: The present report aims to describe a case of invasive ductal carcinoma in a young patient. **Methodology:** Active search in medical records. **Case:** B.S.M., female, 18 years old, arrived at the oncology service with diagnosed ductal infiltrative carcinoma, with a tumor measuring 6 x 8 cm, staging T4 Nx M1, with metastasis to lungs, and marker level CA 15-3 was 9,2, 4 months ago. The histopathology presented the carcinoma was moderately differentiated with intraductal component associated and the immunohistochemistry shows 95% for both estrogen and progesterone receptors, negative for HER-2 and level of 20% for Ki-67, both of them from 5 months ago. The thoracic CT from 4 months ago showed many non-calcified nodules sparse in pulmonary parenchyma up to 14 mm, compatible with implants, increased density in retroareolar region measuring 39 x 34 mm with slight thickening of adjacent skin, two nodules of 5 mm each and axillary lymph nodes up to 11mm. At physical examination, the tumor was compromising 70% of the right breast, which was presenting nipple retraction and ulceration. Palliative chemotherapy was initiated with Fluoracil, Doxorubicin and Cyclophosphamide (FAC) once 4 months ago (11/21/17), that was switched to Doxorubicin, Cyclophosphamide and Paclitaxel (AC-T) for unavailability of Fluoracil at the Institution. The patient was submitted to 3 sessions of FAC 3 months ago (01/22/18). At 02/16/18, at physical examination right breast presented a tumor sized 7 x 6 with nipple ulceration, demonstrating fail of treatment, and then was administered the first cycle of Docetaxel. At physical examination at 03/09/18, right breast presented a tumor of 5 x 5 cm with nipple injury on healing phase and the patient was conducted to the second cycle of Docetaxel. **Considerations:** Breast cancer is considered rare before the age of 35, and the risk of having it grows rapidly with age, with its apex appearing between 40 and 60 years. The patient of the case for being 18 years of age demonstrates an exception to the age values in the statistical data. Thus, new studies are needed to understand the appearance of breast cancer in young patients.

DOI: 10.29289/259453942018V28S1047

SYNTHETIC CHALCONES CYTOTOXIC ACTIVITY ON EHRlich ASCITIC TUMOR CELLS (MURINE BREAST CANCER)

Eliane B. Nunes^{1,2*}, Aline Bernardes³, Caridad Noda-Perez³, Stanislaw P. Cardozo¹, Hugo D. Silva², Ingrid O. Travassos², Paula F. F. Silva², Elisângela P. Silveira-Lacerda²

¹Programa de Pós-graduação em Inovação Farmacêutica, Faculdade de Farmácia, Universidade Federal de Goiás (UFG) – Goiânia (GO), Brazil.

²Laboratório de Genética Molecular e Citogenética, Instituto de Ciências Biológicas, UFG – Goiânia (GO), Brazil.

³Instituto de Química, Instituto de Química, UFG – Goiânia (GO), Brazil.

*Corresponding author: ebnunes@gmail.com

Breast cancer is the world leading cause of women death. The chemotherapy has presented several side effects and many cases of chemo-resistance. Thus, research of new antineoplastic molecules with less aggressive effects is necessary. Chalcones have demonstrated extensive pharmacological potential including antineoplastic. **Objective:** The aim of this study was to evaluate *in vitro* cytotoxic effect induced for synthetic chalcones CLF and DMF on Ehrlich Ascitic Tumor (TAE) cells of murine mammary carcinoma, by MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) colorimetric assay as described by Mosman (1983). **Methodology:** The compounds CLF and DMF were solubilized in dimethylsulfoxide 1%. TAE cells were collected from Murine peritoneal cavity, washed with PBS and maintained in 5% CO₂ for 24h at 37°C in humidified atmosphere. After, RPMI-1640 medium was supplemented with 10% FBS, 1% penicillin/streptomycin and 0.3% amphotericin. The 1.0×10⁵ TAE cells were plated in 96-well tissue culture plates and treated with different concentrations of CLF and DMF (0.2, 2.0, 20, 50, 100 and 200 µM) for 48 h. After treatment, 10 µL of MTT (5 mg.mL⁻¹) was added to each well, and the plates were incubated at 37°C for 3 h. The purple formazan crystals were dissolved in 50 µL of SDS (dodecyl sulfate sodium), and the absorbance was determined at 545 nm. The cell viability was calculated: viability (%) = (absorbance of the treated wells)/(absorbance of the control wells) ×100. The tests were performed in triplicates and IC₅₀ (concentration (µM) that results in a 50% reduction in cellular viability) was obtained from sigmoidal dose-response curves (nonlinear regression) using the software GraphPad Prism 5.0 for Windows. **Results:** The chalcones CLF and DMF presented a statistically significant cytotoxic effect inducing cell death in a dose dependent manner. The chalcones inhibit TAE cells viability with an estimated IC₅₀ of 22.30±5.10 µM and 46.30±6.10 µM, respectively at 48 h of treatment, by nonlinear regression curve. **Conclusions:** The chalcones exhibits significant cytotoxicity in 48h against TAE cells. The CLF was more potent than DMF, but both results showed important dose-dependent biological property of chalcones. Therefore, future studies will be necessary to identify the molecular mechanisms where compound operates.

DOI: 10.29289/259453942018V28S1048

PROGNOSTIC EVALUATION OF PATIENTS WITH TRIPLE NEGATIVE IMMUNOPHENOTYPE, SUBMITTED TO NEOADJUVANT CHEMOTHERAPY

Fábio F. O. Rodrigues¹, Etiénne A. Bastos¹, Laís A. Inaba¹, Eveline A. Silva¹, Rodrigo M. Silva¹, Carlos E. Fristachi¹¹Instituto do Câncer Doutor Arnaldo Vieira de Carvalho – São Paulo (SP), Brazil.

Objectives: Evaluate the characteristics of patients with triple negative (TN) submitted to neoadjuvant chemotherapy attended at the Dr Arnaldo Vieira de Carvalho Cancer Institute (ICAVC) from 2006 to 2012. **Methods:** A longitudinal study was performed, descriptive type, retrospective, of patients with invasive breast carcinoma attended at ICAVC in Sao Paulo; data were analyzed on the data base of Microsoft Excel®. **Results:** Were evaluated 267 patients, among them, 24 (9%) presented TN breast neoplasm by the immunohistochemical study; 21% of them underwent neoadjuvant chemotherapy, having as main regimens the association of anthracycline and taxane; 4% presented pathological complete response (pCR) and 29% evolved with distant metastasis; being the disease free survival and mean global survival of 3 years calculated with Kaplan Meier. **Conclusion:** With this study, we observed 27% of triple negative tumors that, differently from the literature, presented the small complete pathological response index of only 4%.

DOI: 10.29289/259453942018V28S1049

IMPACT OF THE NEW STAGING TNM AJCC ON THE EVALUATION OF PATIENTS WITH BREAST CANCER

Fábio F. O. Rodrigues¹, Étienne A. Bastos¹, Laís A. Inaba¹, Eveline A. Silva¹, Rodrigo M. Silva¹, Carlos E. Fristachi¹

¹Instituto do Câncer Doutor Arnaldo Vieira de Carvalho – São Paulo (SP), Brazil.

Objectives: To evaluate the importance of the incorporation of the biological markers on the TNM 8th Edition to improve the staging of patients attended on the Dr Arnaldo Vieira Cancer Institute (ICAVC) from 2006 to 2012. **Methods:** A longitudinal study was performed, descriptive type, retrospective, of patients with breast carcinoma attended at ICAVC in Sao Paulo; data were analyzed on the data base of Microsoft Excel@. **Results:** Were evaluated 119 patients; of wich 58% presented changes related to staging accomplished in 2012, of these 52% presented changes with upstaging, 17.4% downstaging and 30.6% remained with the same clinical staging. **Conclusion:** With that study, we observed that most patients presented modification on the clinical staging, wich shows the importance of the incorporation of the biological markers to improve the related knowledge of the prognoses and treatment of breast cancer.

DOI: 10.29289/259453942018V28S1050

BIOLOGICAL BEHAVIOR OF BREAST CARCINOMA *IN SITU* FROM JANUARY 2006 TO DECEMBER 2012

Fábio F. O. Rodrigues¹, Etiénne A. Bastos¹, Laís A. Inaba¹, Eveline A. Silva¹, Rodrigo M. Silva¹, Carlos E. Fristachi¹

¹Instituto do Câncer Doutor Arnaldo Vieira de Carvalho – São Paulo (SP), Brazil.

Objectives: Evaluate the incidence of breast carcinoma *in situ* on patients attended at the Dr Arnaldo Vieira de Carvalho Cancer Institute from 2006 to 2012. **Methods:** A longitudinal study was performed, descriptive type, retrospective, of patients with breast carcinoma *in situ* attended at ICAVC in Sao Paulo; data were analyzed on the data base of Microsoft Excel®. **Results:** Were evaluated 267 patients attended at the mastology outpatient clinic, of wich 19 (7.1%) presented a diagnosis of breast carcinoma *in situ* by the core needle biopsy, being 13 (5%) histologically confirmed at the biopsy of the surgical part, of these 5% presented lobular breast carcinoma *in situ* (CLIS), two (10%) of the patients with carcinoma *in situ* presented recurrence, both with invasive component tumor, with hormone receptor positive status and HER-2 positive with high rates of KI67, one patient underwent mastectomy and adjuvant chemotherapy and the other one underwent chemotherapy only, since the first surgical approach was mastectomy; none of these patients presented distant-metastasis until now. **Conclusion:** With our study, it was observed that the incidence of breast carcinoma *in situ* at ICAVC was of 7.1%, being compatible with the prevalent incidence of current literature; in addition, it is understood that the presence of breast carcinoma *in situ* is a marker of risk for invasive carcinoma in relapse.

DOI: 10.29289/259453942018V28S1051

PROGNOSTIC EVALUATION OF PATIENTS WITH LUMINAL B IMMUNOPHENOTYPE, SUBMITTED TO NEOADJUVANT CHEMOTHERAPY ATTENDED AT DR ARNALDO VIEIRA DE CARVALHO CANCER INSTITUTE FROM 2006 TO 2012

Fábio F. O. Rodrigues¹, Étienne A. Bastos¹, Laís A. Inaba¹, Eveline A. Silva¹, Rodrigo M. Silva¹, Carlos E. Fristachi¹

¹Instituto do Câncer Doutor Arnaldo Vieira de Carvalho – São Paulo (SP), Brazil.

Objectives: Evaluate the characteristics of the patients with Luminal B breast cancer submitted to neoadjuvant chemotherapy, attended at Dr Arnaldo Vieira de Carvalho Cancer Institute from 2006 to 2012. **Methods:** A longitudinal study was performed, descriptive type, retrospective, of patients with invasive breast carcinoma attended at ICAVC in Sao Paulo; data were analyzed on the data base of Microsoft Excel[®]. **Results:** Were evaluated 267 patients, in which 82 (31%) presented breast neoplasm with compatible immunohistochemistry with luminal B tumors, of these, 19 (7.1%) were submitted to neoadjuvant chemotherapy, 4 (21%) presented pathological complete response (pCR), both mammary and axillary, 1 (5.2%) patient presented distant metastasis for bone and none of them had local recurrence. The overall mean survival was 12 years and the disease-free survival was 4 years calculated with Kaplan Meier. **Conclusion:** The pCR was 21% in patients with luminal B tumors seen in our service, with high overall survival rates due to the possibility of adjuvancy with hormonal block, wich is consistent with the data present in the current literature regarding the prognosis of such patients.

DOI: 10.29289/259453942018V28S1052

EVALUATION OF PATIENTS WITH INVASIVE BREAST CARCINOMA SUBMITTED TO NEOADJUVANT CHEMOTHERAPY ATTENDED AT DR ARNALDO VIEIRA DE CARVALHO CANCER INSTITUTE FORM 2006 TO 2012

Fábio F. O. Rodrigues¹, Etiénne A. Bastos¹, Laís A. Inaba¹, Eveline A. Silva¹, Rodrigo M. Silva¹, Carlos E. Fristachi¹

¹Instituto do Câncer Doutor Arnaldo Vieira de Carvalho – São Paulo (SP), Brazil.

Objectives: Evaluate the characteristics of patients with invasive breast carcinoma submitted to neoadjuvant chemotherapy attended at Dr Arnaldo Vieira de Carvalho Cancer Institute form 2006 to 2012. **Methods:** A longitudinal study was performed, descriptive type, retrospective, of patients with invasive breast carcinoma attended at ICAVC in Sao Paulo; data were analyzed on the data base of Microsoft Excel@. **Results:** Were evaluated 267 patients, of wich, 48 (18%) underwent adjuvant chemotherapy with predominance of patients with tumors T3 (35.4%), DCI (91.6%) and luminal (54.1%); the main regimen used was anthracycline in association with taxane (45.8%); 14.5% patients presented complete clinical-pathological response (pCR); 74.5% underwent conservative surgery; 5.4% presented local recurrence and 23.6% presented distant metastasis in the mean period of 2 years (disease-free survival), the overall survival was 10 years. **Conclusion:** Neoadjuvant chemotherapy was shown in this study as a treatment of choice in locally advanced tumors targeting pCR with increased conservative breast surgeries.

DOI: 10.29289/259453942018V28S1053

COMPARATIVE ANALYSIS OF THE USE OF FINE NEEDLE ASPIRATION IN THE DIFFERENTIAL DIAGNOSIS OF BREAST NODES IN THE STATE OF GOIÁS

Fernando C. Silva¹, Edison B. L. B. Junior¹, Cassiano S. Coutinho¹, Rayssa F. Silva¹, Rayssa O. Glória¹, Fabiana C. Q. S. Anjos¹, Vantuir J. D. Mota¹, Priscila F. Barbosa¹

¹Centro Universitário UNIRG – Gurupi (TO), Brazil.

Objective: To analyze the result of fine needle aspiration (FNA) according to the most prevalent types of nodules in malignant cases, considering the color/race and the age bracket of the patients. **Methodology:** A cross-sectional study based on data collected in the Breast Cancer Information System (SISMAMA) - from Information Technology Department of the Brazilian Public Health Care System (DataSus) based on data from cytopathological breast exams realized by Fine Needle Aspiration (FNA) performed in the State of Goiás in the period from February 2012 to February 2014. The present study comprised the following age groups: 25–29; 30–34; 35–39; 40–44; 45–49; 50–54; 55–59; 60–64; 65–69; over 70 years old; in addition to analyzing the prevalence of white, black and drab colors/races. Solid, cystic and solid-cystic nodules were studied. **Results:** In the analyzed period, 1052 exams were performed that detected breast nodules, of which 76.61% (n=806) were solid, 11.21% (n=118) cystic and 12.16% (n=128) solid-cystic. Of total exams, 615 (58.46%) were submitted to FNA and, through the results, the solid nodule presented the highest percentage of malignancy (5.85%; n=36). In the analysis of the patients diagnosed with this type of alteration, there was a predominance of drab color, totaling 230 cases (21.86%). Taking into consideration their age group, it was observed that women between 40 and 49 years old were associated with a higher percentage of breast nodules (27.33%), confirming statistical significance ($p < 0.05$). **Conclusion:** The results of this study demonstrate that several women (n=1052) in the state of Goiás were diagnosed with breast lumps according to the analyzed period, and more than half of this population (58.46%) was submitted to the FNA. By assessing the types of nodules and those who most frequently show positivity for malignant processes, it was noticed that the solids correspond to this category, being that such alteration guided the present study for comparisons assessed later. The prevalence in terms of age range occurred in the fourth decade, confirming Brazilian society of Mastology statistics by saying that breast cancer occurs mainly after the age of 40. With regard to the color/race, drab patients exhibit higher percentage of solid breast lumps.

DOI: 10.29289/259453942018V28S1054

RUTHENIUM (II)/AMINO ACID/DIPHOSPHINE COMPLEXES AS APOPTOSIS INDUCERS IN BREAST CANCER CELLS AND THEIR TOXIC EFFECTS IN ANIMAL MODELS

Francylli Mello-Andrade^{1,2}, Clever G. Cardoso³, Paulo Melo-Reis⁴, Cesar Grisólia⁵, Carlos Castro⁶, Carlos Menck⁷, Alzir A. Batista⁸, Elisângela Silveira-Lacerda^{1,2}

¹Laboratório de Oncologia Experimental, Instituto de Física, Universidade Federal de Goiás (UFG) – Goiânia (GO), Brazil.

²Departamento de Genética, Instituto de Ciências Biológicas, UFG – Goiânia (GO), Brazil.

³Departamento de Histologia, Embriologia e Biologia Celular, Instituto de Ciências Biológicas, UFG – Goiânia (GO), Brazil.

⁴Departamento de Biomedicina, Pontifícia Universidade Católica de Goiás – Goiânia (GO), Brazil.

⁵Departamento de Genética e Morfologia, Universidade de Brasília – Brasília (DF), Brazil.

⁶Departamento de Fisiologia, Instituto de Ciências Biológicas, UFG – Goiânia (GO), Brazil.

⁷Departamento de Microbiologia, Instituto de Ciências Biomédicas, Universidade de São Paulo – São Paulo (SP), Brazil.

⁸Departamento de Química, Universidade Federal de São Carlos – São Carlos (SP), Brazil.

Cytotoxicity and mechanism of action of ruthenium(II)/amino acids/diphosphine complexes were evaluated against breast cancer (MDA-MB-231 cells), as well as their toxic effects in animal models. The cell death type induced in MDA-MB-231 cells by compounds was evaluated measuring Annexin V-positive number, the activated caspases levels, and by morphological features. In order to clarify which mechanisms are responsible for led breast cancer cells to death, we evaluated whether these compounds can cause DNA damage, changes in cell cycle kinetic, mitochondrial dysfunction, and ultrastructural alterations in MDA-MB-231 cells. As toxicity assessment is a required step for preclinical study of novel metal based compounds, the acute oral toxicity was evaluated, and the genetic toxicity was determined by Micronucleus (MN) and Comet assay protocols on Swiss mice treated with these complexes. Zebrafish model was used to evaluate toxicity of Ru(II)/amino acids complexes during embryonic and larval development, more specifically, the mortality and hatching rates of zebrafish were determined. Ru(II)/amino acids complexes induced apoptosis in MDA-MB-231 cells by increase in number of Annexin V-positive cells, morphological changes, loss of mitochondrial membrane potential, and caspases-3 and 7 activation. Although these compounds have a weak interaction to DNA molecule, it was observed DNA damage, probably due to reactive oxygen species production related to mitochondrial membrane depolarization. As can be seen by ultrastructural analysis, these complexes can reduce mitochondrial amount in MDA-MB-231 cells. Thus, Ru(II)/amino acids complexes are more active for tumor cells, and their mechanism of action are related to induction cell cycle block, DNA damage, and mitochondrial dysfunction, lead to apoptosis involving p53, PARP cleavage and caspases activation. In addition, these compounds were very well tolerated orally, and it was observed lack of micronuclei formation in bone marrow cells and low DNA damage in peripheral blood cells from Swiss mice. As regard to toxicity on zebrafish embryo development, the compounds caused low embryotoxicity, being mainly observed hatching delay and mortality at high concentrations. Altogether our findings suggest that ruthenium (II)/amino acids/diphosphine complexes induce cellular and molecular responses in breast cancer cells leading to mitochondria-mediated apoptosis, and they are not harmful, presenting low systemic toxic effects.

DOI: 10.29289/259453942018V28S1055

A RARE CASE OF METASTATIC MELANOMA IN THE BREAST

Aline R. Nunes¹, Célio S. R. Vidal², Délio M. Conde¹, Erika P. S. Silva^{1,2,3}, Flávia V. Cabero¹, Gustavo L. Q. Lima⁴, Kauana C. S. Nascimento⁴, Rogério B. Ferreira^{1,2}

¹Mastology Service, Hospital Materno Infantil de Goiânia – Goiânia (GO), Brazil.

²Mastology Service, Hospital Geral de Goiânia – Goiânia (GO), Brazil.

³Mastology Department, Hospital Geral de Goiânia – Goiânia (GO), Brazil.

⁴Hospital Geral de Goiânia – Goiânia (GO), Brazil.

Objective: To report a case of metastatic melanoma in the breast. **Case:** woman, 66 years, complaining of mass in the left breast. She reported treatment of melanoma in the gluteal region in 2011 and left breast cancer in 2016 (luminal B (RH+, HER2-), undergoing mastectomy with sentinel lymph node biopsy, chemotherapy and hormone therapy). Physical examination: breast implants; regular, mobile mass, 10h of left breast, measuring about 2.0 cm. Ultrasonography showed a solid-cystic mass, predominantly solid, with a hypoechogenic content, with its largest axis parallel to the skin, with slightly lobed contours and well-defined limits, with an apparent increase in the surrounding echogenicity, at 10 hours of the left breast, measuring 1.9 x 1.7 x 0.9 cm, suggestive of complex cyst, BI-RADS 4. Mammography: mass partially visualized in the medial quadrants of the left breast, visible, only visible in the craniocaudal view. The hypothesis was that of recurrence of breast cancer. Patient underwent extensive excision of the lesion to obtain surgical margins of neoplasia. Histopathological findings, associated with immunohistochemistry (AE1/AE3: negative. S-100 protein, melan A and Hmb-45: positive), were compatible with the diagnosis of metastatic melanoma for the breast. Subsequently, a recurrence of melanoma in the gluteus was diagnosed. Chest CT demonstrated multiple lung lesions. Patient is being treated for melanoma, and using letrozole. **Conclusion:** metastatic melanoma for the breast is uncommon. The breast surgeon should be aware of this possibility, especially in women with a history of melanoma. In addition, there is a need to work together with the pathologist, making diagnosis and treatment as soon as possible.

DOI: 10.29289/259453942018V28S1056

EPIDEMIOLOGICAL ANALYSIS OF GRANULOMATOUS MASTITIS AT THE MASTOLOGY SERVICE AT HOSPITAL MATERNO INFANTIL DE GOIÂNIA

Aline R. Nunes¹, Célio S. R. Vidal², Délio M. Conde², Erika P. S. Silva^{2,3,4}, Flávia V. Cabero¹, Gustavo L. Q. Lima¹, Rogério B. Ferreira^{2,3}, Paola F. Freitas⁵

¹Hospital Geral de Goiânia – Goiânia (GO), Brazil.

²Mastology Service, Hospital Materno Infantil de Goiânia – Goiânia (GO), Brazil.

³Mastology Service, Hospital Geral de Goiânia – Goiânia (GO), Brazil.

⁴Mastology Department, Hospital Geral de Goiânia – Goiânia (GO), Brazil.

⁵Maternidade Municipal Aristina Cândida – Senador Canedo (GO), Brazil.

Objectives: The present study aims to evaluate the patient's profile, along with ways of diagnosing granulomatous mastitis (GM) disease, duration of treatment, remission and relapse time. **Methodology:** A retrospective analysis of physical records of patients treated between January 2010 and October 2017, at Hospital Materno Infantil de Goiânia. **Results:** During this time, ten women were diagnosed and treated for GM. All women underwent clinical breast examination to identify palpable tumors. Ultrasonography was the initial imaging method used for the evaluation, however mammography was performed in three cases due to the screening age range for breast cancer. The definitive diagnosis was obtained by core biopsy or surgical excision, with histopathological result confirming GM. The median age was 36.1 years. As for the use of oral contraceptive, the mean time of use was 5.5 years, and only one patient did not report the use of it. Only one patient was nulliparous, the others had been pregnant at least once, with an average time of breastfeeding of 1.5 years and all had interrupted it for more than two years. Two patients had diabetes mellitus, one of whom was diagnosed during the investigation of the GM. The most common symptoms were pain (90%), breast masses (80%) and association with abscess (60%). Three patients used antibiotics because of the association of abscess infections with the inflammatory process of mastitis. Nine patients have used corticosteroids to treat the disease since diagnosis. The only patient with abscess drainage, quadrantectomy and antibiotic therapy was also the only one that present with local recurrence two months after treatment, and remission was achieved after using corticotherapy. Regarding treatment with corticosteroids, dose reduction was done according to clinical response. The mean time of treatment was 3.1 months [1-7 months], and mean follow-up time of the patients was 24.8 months [9-96 months]. **Conclusion:** The study presents the same epidemiology, follow-up time and treatment response in accordance with international literature. Implementing a flow chart of care, diagnosis, therapy and follow-up of GM improves the quality and assertiveness of early treatment, avoiding misuse of antibiotics.

DOI: 10.29289/259453942018V28S1057

PATTERNS OF POST-OPERATIVE IRRADIATION IN BREAST CANCER PATIENTS SUBMITTED TO NEOADJUVANT CHEMOTHERAPY

Kennya Medeiros L. de B. Lima¹, Thiago B. de Freitas¹, Allan A. Lima Pereira¹, Saulo Brito Silva¹, Heloisa de Andrade Carvalho^{2,3}, José R. Filassi⁴, Max S. Mano^{1,5}, Gustavo Nader Marta^{1,4}

¹Department of Radiology and Oncology, Radiation Oncology Unit, Faculdade de Medicina, Universidade de São Paulo, Instituto do Câncer do Estado de São Paulo – São Paulo (SP), Brazil.

²Department of Radiology and Oncology, Radiation Oncology Unit, Faculdade de Medicina da Universidade de São Paulo, Instituto de Radiologia – São Paulo (SP), Brazil.

³Department of Radiation Oncology, Hospital Sírio-Libanês – São Paulo (SP), Brazil.

⁴Department of Gynecology and Obstetrics, Instituto do Câncer do Estado de São Paulo, Universidade de São Paulo – São Paulo (SP), Brazil.

⁵Department of Clinical Oncology, Hospital Sírio-Libanês – São Paulo (SP), Brazil.

Introduction: The neoadjuvant chemotherapy (NAC) approach can expressively influence radiation therapy indications after breast surgery by theoretically altering the loco-regional relapse risk in accordance with clinical stage features. Precisely, it could reduce this risk for a subgroup of patients with locally advanced breast cancer who achieved a good pathologic response after NAC. Thus, this might raise clinical questions of whether these more favorable subset of patients do benefit from PORT therapy or whether radiation should be performed only for patients with gross residual tumor after neoadjuvant chemotherapy. In addition, there are many uncertainties about the most appropriate radiation therapy fields in patients who received NAC given that all recommendations are usually based on initial clinical and pathological aspects.

Objectives: The aim of the current study was to assess the radiation therapy fields and survival outcomes in breast cancer patients who underwent (NAC) followed by surgery. **Methods:** We performed a retrospective analysis of all non-metastatic breast cancer patients treated between 2008 and 2014 at our institution, who received NAC and post-operative radiation therapy (PORT). **Results:** A total of 528 women were included of whom 396 were submitted to mastectomy or adenomastectomy. Most (92.8%) of the patients had locally advanced disease (clinical stage IIB to IIIC). All patients underwent irradiation for breast or chest wall. Most patients received radiation therapy of the supraclavicular and axillary (levels II and III) nodes (87.1 and 86.4% for breast-conserving surgery and 95.1 and 93.8% for mastectomy / adenomastectomy, respectively). Irradiation of level I axillary and internal mammary nodes was uncommon. The most common radiation therapy schedule was the conventional dose of 50Gy to 50.4Gy in 25 to 28 fractions. The mean overall survival was 66.6 months and the mean disease-free survival was 54.6 months. **Conclusions:** After NAC, most patients received irradiation of the breast/chest wall and axillary and supraclavicular nodes. Indications were based on initial disease presentation associated with relapse risk factors. In this setting, post-operative radiation therapy to breast/chest wall with or without regional nodal irradiation was effective management that is associated with acceptable survival rates.

DOI: 10.29289/259453942018V28S1058

DOES THE ADDITION OF THE NEW PROGNOSTIC BIOMARKERS ON EIGHTH EDITION OF AJCC CHANGE BREAST CARCINOMAS STAGING?

Jéssica E. P. Ramos^{1,2}, Cesar A. S. T. Vilanova-Costa^{1,3}, Juliana F. Paes⁴, Vera A. Saddi^{1,2,4,5}

¹Laboratório de Genética e Biodiversidade, Programa de Pós-graduação em Ciências Ambientais e Saúde, Universidade Católica de Goiás – Goiânia (GO), Brazil.

²Escola de Ciências Médicas, Farmacêuticas e Biomédicas, Pontifícia Universidade Católica de Goiás – Goiânia (GO), Brazil.

³Laboratório de Biologia Tumoral, Hospital Araújo Jorge, Associação de Combate ao Câncer em Goiás – Goiânia (GO), Brazil.

⁴Programa de Pós-graduação em Ciências da Saúde, Universidade Federal de Goiás – Goiânia (GO), Brazil.

⁵Laboratório de Oncogenética e Radiobiologia, Instituto de Ensino e Pesquisa, Associação de Combate ao Câncer em Goiás – Goiânia (GO), Brazil.

Tumor staging plays a pivotal role in the treatment against cancer. The new revision of AJCC included histologic grade, estrogen receptor (ER), progesterone receptor (PgR), HER2 and Ki-67 expression as prognostic biomarkers for breast cancer (BC) staging. This study aimed to evaluate if the new prognostic biomarkers included on the eighth edition of AJCC are sufficient to change BC staging. Clinicopathological and immunohistochemical characteristics of 92 BCs, 36 triple-negative (TNBC) and 56 non-triple-negative (NTNBC) breast tumors, were collected from medical records of the Pathology Department of Araújo Jorge Hospital-ACCG (Goiânia, Goiás, Brazil). The study was approved by the institutional Ethics Committee of Araújo Jorge Hospital (Report n° 948.930, 2015). BCs were evaluated under the recommendations of the seventh edition of AJCC and then compared to new (eighth) edition criteria. Results show that 2.8% of TNBC progressed from stages I-II to III-IV ($p=0.81$) and 5.4% of NTNBC regressed from stages III-IV to I-II ($p=0.57$), revealing that the addition of new prognostic biomarkers abovementioned had no influence on BC staging. It is well known that, compared with other tumor types, TNBCs have high expression of cyclin E, caveolins 1 and 2, P-cadherin; low expression of E-cadherin and cyclin D1. TNBCs also present aberrant expression levels for p53, BRCA1 and PARP. These biomarkers can be important on BC staging and the lack of a more comprehensive assessment of possible BC prognostic biomarkers can be a weakness of new AJCC version.

DOI: 10.29289/259453942018V28S1059

LATISSIMUS DORSI MUSCULOCUTANEOUS FLAP RECONSTRUCTION AFTER NIPPLE SPARING MASTECTOMY COMPLICATION: CASE REPORT

Jorge V Biazus¹, Andrea Damin¹, José A Cavalheiro¹, Angela Zucatto¹, Rodrigo Cericatto¹, Marcia P. Melo¹, Anita S. Cassiano¹, Murilo L. Brazan¹

¹Hospital de Clínicas de Porto Alegre, Universidade Federal do Rio Grande do Sul – Porto Alegre (RS), Brazil.

Introduction: The latissimus dorsi (LD) myocutaneous flap has been an option for partial and total breast reconstruction. The LD muscle is supplied by thoracodorsal artery and vein. These substantial vessels branches into a network of smaller tributaries, which provide a rich blood supply both for the muscle itself and for the overlying skin. Reconstruction with the LD flap is now considered to be the primary procedure of choice for many patients for reasons of robustness and low rates of complications. The indications for autologous reconstruction are variable and include women that do not want prosthetic reconstruction, have had a prior failed prosthetic reconstruction or have had prior radiation to the breast or chest wall. **Case:** A 45-year-old female patient with multicentric breast carcinoma underwent nipple sparing mastectomy on the right breast with prosthesis reconstruction. The patient progressed with lower quadrants skin necrosis of the right breast. Patient had no local or systemic signs of infection and no exposure of the prosthesis. Four weeks later the first surgery, a LD myocutaneous flap was performed after necrosed area debridement with nipple preservation and rescue of the first surgery prosthesis. The prosthesis was covered by the preserved major pectoral muscle and the LD flap. Patient evolved well without complications. **Conclusion:** The use of musculocutaneous flaps in breast reconstructive surgery has a wide variety of applications. The LD flap is robust and reliable with a failure rate of less than 1%. It can be safely utilized even in “high-risk” cases such as heavy smokers and obese patients. It is an especially useful technique for treatment of implant breast reconstruction complications.

DOI: 10.29289/259453942018V28S1060

QUALITY OF LIFE IN PATIENTS WITH LYMPHEDEMA SUBMITTED TO SURGICAL TREATMENT FOR BREAST CÂNCER ATTENDED AT THE PHYSIOTHERAPY AMBULATORY OF A REFERENCE HOSPITAL OF THE FEDERAL DISTRICT

Kalléria W.C. Borges¹, Laércio L. Luz¹¹Instituto Hospital de Base do Distrito Federal – Brasília (DF), Brazil.

Introduction: Lymphedema, secondary to surgical treatment of breast cancer, can result in functional, social and emotional complications that limit the patient's ability to lead a normal life. The impact on the patient's life can be measured through the application of questionnaires that evaluate the quality of life. **Objective:** to evaluate the quality of life in patients with lymphedema secondary to the surgical treatment of breast cancer and how much this is related to the functional limitations caused by this complication. **Method:** descriptive cross-sectional study in patients with lymphedema secondary to surgical treatment for breast cancer, attended at the physiotherapy ambulatory of a reference hospital of the Federal District, from November 2017 to January 2018. DASH instruments were used to evaluate the functionality and EORTC QLQ-C30/BR23 for quality of life assessment. Results: Fifty-one women, with a mean age of 54 years, were included and 76.5% of them presented some type of pain. The mean total functional score was 44.37 ± 20.08 . The functional quality of life scales with the best and worst result were social functioning and sexual functioning, respectively. The scales of symptoms with the worst outcomes were insomnia, financial distress, pain and symptoms in the arm. The relationship between the existence of symptoms and low functionality was identified. **Conclusion:** Symptoms caused by lymphedema secondary to the treatment of breast cancer significantly affect quality of life and functional capacity, including physical, social and emotional aspects.

DOI: 10.29289/259453942018V28S1061

PREVALENCE OF BREAST ABNORMALITIES FOUND IN ULTRASSOUND

Laís L. Neves¹, Fernanda G. Filardi¹, Dânisia S. Carrijo¹, Julia B. Gomes¹, Milena V. M. Moreira¹, Felipe Augusto A. Carvalho¹, Raphael Costa Lima¹, Waldemar Naves Amaral²

¹Faculdade de Medicina, Centro Universitário de Anápolis (UniEVANGÉLICA) – Anápolis (GO), Brazil.

²Faculdade de Medicina, Universidade Federal de Goiás – Goiânia (GO), Brazil.

Introduction: Breast disorders are very frequent in the female population, and can be divided into benign and malignant ones. For its evaluation, ultrasonography (USG) provides important information on breast findings, providing additional data to the other diagnostic methods. Within this, the Breast Imaging Reporting and Data System (BI-RADS) classification universalizes the diagnosis and assists in the standardization of an examination that is a dependent operator. **Objectives:** Quantify the prevalence of normal findings in breast USG, establish the most frequent pathological findings in breast USG and establish the frequency of BI-RADS findings. **Methodology:** This is a cross-sectional retrospective descriptive study, carried out with 209 breast ultrasonography reports from the Center for Fetal Medicine and Human Reproduction of Goiânia - Fértil Diagnósticos, carried out in 2016 and 2017. **Results:** Considering the whole sample, 209 mammary USG reports, of which 143 had unchanged results (68.42%), with 80% of reports classified as BI-RADS 1 (absence of abnormal findings). The second BI-RADS most found was classified as 2, in which was found 10,5% of benign abnormalities. Already 31.5% of the reports presented sonographic alterations, with the main abnormality observed being the benign functional alterations, present in 56% of them. Then, the cystic nodes occupied the position of the second most frequent abnormality (36.6%). It also was analyzed 16 reports of solid nodules (24.4%). **Conclusion:** In this way, the prevalence of normal USG was 68%, whereas among the altered reports, benign functional alterations were the most frequent followed by solid nodules. Regarding BI-RADS, types 1 and 2 were the most found, according to Brazilian literature.

DOI: 10.29289/259453942018V28S1062

MALIGNANT PHYLLODES TUMOR OF THE BREAST: REPORT OF TWO CASES AND A RARE ASSOCIATION

Leandro G. Oliveira¹, Barbara A. Barbosa¹, Elaine X. Machado¹, Thamires S. Lopes¹, Ana Cláudia G. Lima¹, Alexandre M. X. de Jesus¹, Sebastião A. Pinto¹, Juarez A. de Sousa¹

¹Instituto Goiano de Oncologia e Hematologia – Goiânia (GO), Brazil.

Introduction: Phyllodes tumors (PT) represent less than 1% of all breast tumors. Most of them are benign, but up to 30% show malignant stroma. Malignant PT of the breast (MPTB) are characterized by marked stromal cellularity, atypia and high mitotic rate (> 10 per 10 high-power fields). Unlike benign PT, MPTB can be clinically aggressive, with local recurrence rates ranging from 23–30%. Malignant change of the epithelium in MPTB is rare. The most frequent occurrence is infiltrating carcinoma of various types while ductal carcinoma *in situ* (DCIS) is more rare. The purpose of this study was to present two cases of MPTB, one of them with a concomitant DCIS. **Case reports:** Case 1: A 49-year-old female patient presented a history of mass in the right breast with accelerated growth. Core biopsy was compatible with a fusocellular neoplasm. Patient underwent simple mastectomy with surgical margins free. Macroscopy evidenced a 12x11 cm lesion. Immunohistochemistry (IHC) showed a tumor with high mitotic rate (12 mitoses/hpf), ki-67 10%, CD-34 and vimentin +, S-100 protein and AE1/AE3 ceratines -, compatible with malignant phyllodes tumor. Case 2: A 54-year-old female patient presented with a firm mass in the left breast with accelerated growth. Core biopsy was compatible with a low grade MPTB (6 mitoses/hpf). IHC was positive for CD-3 and vimentin and ki-67 15%. Patient underwent left adenomastectomy with sentinel node biopsy. The features were consistent with MPTB of 4.5x3.0cm. The epithelial component showed a 0.7x0.6cm DCIS, nuclear grade 2, solid with hormonal-receptors positives (>90%). Surgical margins were free of tumor. **Discussion:** PT represent a specific subset of breast soft tissue tumors composed of a connective stroma and epithelial elements. These epithelial components may show a range of metaplastic (apocrine, squamous) and proliferative changes. However, DCIS in association with PT is extremely rare, with only few cases reported in the literature. The role of adjuvant radiation or chemotherapy following partial breast surgery (PBS) or mastectomy is still unclear due to the rarity of these tumors. **Conclusion:** In this study, we described a MPTB that harbored an intraductal carcinoma. This result could determine an absolute indication for adjuvant radiotherapy if the patient underwent PBS. Although the stroma in these tumors is commonly the aggressive component, the epithelial component also requires close and careful histological appraisal.

DOI: 10.29289/259453942018V28S1063

PATHOLOGICAL FEATURES AND GENETIC TESTING AMONG BREAST CANCER YOUNG WOMEN: A SINGLE-INSTITUTION EXPERIENCE

Leandro G. Oliveira¹, Ana C. G. Lima¹, Danilo A. Gusmão¹, Deidimar C. B. Abreu¹, Frank L. B. Rodrigues¹, Alexandre Marchiori¹, Juarez A. Sousa¹, Rosemar M. S. Rahal¹

¹Instituto Goiano de Oncologia e Hematologia – Goiânia (GO), Brazil.

Objectives: To describe the anatomopathological characteristics and the access to genetic testing among breast cancer young women attended at a private institution in Goiás-Brazil. **Methodology:** In this descriptive and retrospective study, were reviewed medical records of breast cancer young women (< or = 40 years at the diagnosis) attended at a private cancer institution from March 2016 to March 2018. Data with respect to age at diagnosis, histopathological and immunohistochemical aspects and access to genetic testing were examined. **Results:** A total of 67 women were identified. The median age at the diagnosis was 36 years (range 26–40 years). The most common histological type was invasive ductal carcinoma (98.5%); 45% (n=30) and 41% (n=28) were grade 3 and 2 tumours respectively. 70% (n=47) were estrogen receptor (ER) positive (at least 1% of positivity). 20,8% (n=14) were triple-negative and 2 women had ER positivity between 1–10%. 26% (n=18) were human epidermal growth factor receptor -2 (HER-2) positive (HER-2 3+ or amplified FISH/ISH). The ki-67 index was >14% in 51 women (76%). Overall, 39% (n=26) got a genetic testing. The mean age of tested patients was 33,8 years. Among the women with available genetic results so far (n=19), six were identified (31%) with pathogenic mutations (BRCA-1= 4; BRCA2+MUTYH=1; SKT11=1). Three of BRCA-1 mutated were triple-negative and the fourth had low hormonal receptor positivity (1%). **Conclusion:** In this real-world and single-institution study, the proportion of triple negative and HER2 positive women was discretely higher than that described for older breast cancer patients. As expected, the majority of women had high grade tumours. A hereditary predisposition is more common amongst young women and may impact decisions on local and systemic disease management. Only the minority of this clinically high-risk group got a genetic testing. In this brief analysis, it was observed that almost one in four tested women harbored a pathogenic genetic mutation. Despite of the efforts to reduce the cancer care disparities, including the access to genetic counseling, significant barriers persist even among patients in private health assistance.

Keywords: Breast cancer in young women; genetic testing; BRCA mutation.

DOI: 10.29289/259453942018V28S1064

MOLECULAR ANALYSIS OF TRANSLOCATIONS ASSOCIATED WITH EWING'S SARCOMA IN THE DIFFERENTIAL DIAGNOSIS OF UNDIFFERENTIATED AND UNCLASSIFIED SARCOMAS

Leonardo R. Soares¹, Marise A. R. Moreira¹, Paula O. C. Queiroz¹, Hugo A. Bayeh¹, Izabela C. S. Albuquerque¹, Miguel A. C. Coutinho¹, Débora F. Rodrigues¹, Ruffo Freitas-Junior¹

¹Brazilian Mastology Research Network – Goiânia (GO), Brazil.

Undifferentiated and unclassified sarcomas constitute a heterogeneous group of sarcomas, including the Ewing sarcoma family of tumors (ESFT). However, some tumors with Ewing sarcoma (ES) morphology remain with undifferentiated histogenesis due to negative or non-informative molecular results. A case of 25-year-old female patient with a massive lesion occupying almost all of the mammary parenchyma on the right, with a hardened appearance and measuring 18.0cm was described. Chest tomography and resonance showed extension of the lesion to the chest wall, with no apparent bone involvement. A histology of the material from core biopsy revealed high grade fusocellular neoplasia with areas of necrosis, atypia and mitosis; and absence of epithelial component represented in the sample. An immunohistochemistry of the lesion revealed negative expression of smooth muscle actin, HHF35, desmin, S100, CD34 and EMA. Positive expression of CD99, CD56 and CD57 markers was also observed. The patient underwent radical surgery and adjuvant treatment with chemotherapy and radiotherapy. In view of the immunohistochemically profile and rosette arrangement in a fusocellular lesion, a molecular evaluation of translocations associated with ES was performed. However, there were no translocations commonly associated with this pathology [t(11; 22) (q24; q12)]. The present report may contribute to a diagnostic investigation of similar cases and to the pre-test orientation of molecular evaluation.

DOI: 10.29289/259453942018V28S1065

EPIDEMIOLOGIC ANALYSIS OF MAMMOGRAPHY IN WOMEN WITH LESS THAN 45 YEARS OF AGE IN GOIÁS BETWEEN 2010–2013

Ana C. L. Caixeta¹, Bruna O. Andrade¹, Lucas de S. Steinmetz¹, Lucas N. Silva¹, Lucas V. Peixoto¹, Matheus L. T. Costa¹, Júlia J. Caetano¹, Vitória N. Macedo¹

¹Universidade Federal de Goiás – Goiânia (GO), Brazil.

Objective: Quantify and analyze mammography made in women with less than 45 years of age in Goiás between 2010–2013. **Methodology:** This is a descriptive study using data from the Sistema de Informações do Câncer de Mama (SISMAMA), which has the objective to evaluate and quantify the mammography in women up to the age of 45, as well as its indication, diagnosis and trace results, and distribute these tests within Goiás. **Results:** In the period of 2010 to 2013 were realized, in the state of Goiás, 53763 mammograms in women under the age of 45 years, which 52134 (96.9%) as screening test and 1629 (3.1%) as diagnostic exam. Among of 53763 exams, 32270 (60.025%) were executed in Goiânia and 21493 (39.985%) in the country towns in the state. From the totality, 39 (0,007%) were performed in girls under 10 years old, 21 (0.04%) in women in the age range of 11 to 14 years, 104 (0,2%) between 15 to 19 years, 233 (0.43%) between 20 to 24 years, 572 (1.06%) in women between 25 to 29 years, 1764 (3.28%) between 30 to 34 years, 9801 (18.23%) between 35 to 39 years and 41229 (76.069%) mammograms were performed in women in the age range of 40 to 44 years. As to BI-RADS (Breast Imaging Reporting and Data System) classification, 6716 as BI-RADS 0, 33799 as BI-RADS 1, 11433 as BI-RADS 2, 1207 as BI-RADS 3, 522 as BI-RADS 4, 65 as BI-RADS 5 and 25 as BI-RADS 6. **Conclusion:** Most of the women under the age of 45 did screening tests (96.9%), therefore, were asymptomatic, and most of them were between the ages of 40–44 years. Among the exams, 62.86% were classified as BI-RADS 1, indicating negative detection, 12,49% were BI-RADS 0, meaning additional imaging is needed, 21.26% were BI-RADS 2, indicating benign finding, 2.24% were BI-RADS 3, showing a probable benign finding, usually followed by a 6-month follow-up mammography, 0,97% were BI-RADS 4, indicating a suspicious abnormality in which a biopsy is considered, 0.12% were BI-RADS 5, being highly suggestive of malignancy and typically evaluated with biopsy, and 0.05% were classified as BI-RADS 6, showing known cancer in which appropriate action should be taken. This results indicates that most of the women under the age of 45 at the state of Goiás do not show malignant or benign signs of breast cancer. However, the early detection of any sign of breast cancer is crucial for its treatment, which brings the discussion if the early screening (under 45 years old) actually changes the overall prognosis of this disease.

DOI: 10.29289/259453942018V28S1066

PREVALENCE OF TYPES OF BENIGN BREAST LESIONS, BY AGE GROUP, IN GOIÁS

Ana C. L. Caixeta¹, Bruna O. Andrade¹, Lucas de S. Steinmetz¹, Lucas N. Silva¹, Lucas V. Peixoto¹, Matheus L. T. Costa¹, Júlia J. Caetano¹, Vitória N. Macedo¹

¹Universidade Federal de Goiás – Goiânia (GO), Brazil.

Objective: To carry out a study of the prevalence of benign breast lesions, by age group, in Goiás, establishing the types that most affect certain ages. **Methodology:** This is a descriptive study of DATASUS data regarding histopathological examination of the breast in the years 2010 to 2013, in Goiás. **Results:** Fibroadenoma with 652 cases is the most prevalent benign lesion, with the highest age group of 15-19 years, with a small decrease of 20-24 years, remaining stable until the age of 49 and regressing thereafter. There was a decrease from 2011, with 197 cases, to 111 cases in 2013. Fibrous condyloma with 235 cases, with an age group with major involvement between 45-54 years (37.44%), is stable between 30-44 years and increase to 45 years with a subsequent decrease after 54 years. There is stability between the number of cases in 2010 (61 cases) and 2011 (68 cases), leading to a small decrease until 2013, with 43 cases. Adenosis WOS (without other specifications) type lesions, with 75 cases, were more prevalent in the 40-54 age group (52%), with small involvement among the young, observing an increase in adult cases and regression in the elderly. There is a decrease in cases between 2010 (27 cases) and 2011 (15 cases), remaining stable until 2013 (15 cases). The ductal hyperplastic lesion without atypia, with 62 cases reported, showed a stability in the number of cases between 30-49 years, with increase between 50-54 years (14 cases) and subsequent decrease between 55-69 years. There is a considerable decrease between 2010 and 2011 (from 23 to 13 cases), with a stabilization until 2013, where 11 cases were reported. **Conclusion:** It is possible to note that the cases that were most reported in Goiás between 2010 and 2013 were, in order of higher prevalence, fibroadenoma, fibrous condyloma, adenosis WOS and ductal hyperplastic lesion without atypia. The most prevalent age group in the observed cases was 40-54 years, except in fibroadenoma, the highest prevalence in adolescents (15-19 years). In the period between 2010 and 2011, there was an increase in cases of fibrous condyloma and a decrease in the cases of adenosis WOS and ductal hyperplastic lesion, in the same period. As of 2011, there was a decrease in cases of fibroadenoma. In 2013 there was a reduction in cases of fibrous condyloma. The number of cases of adenosis WOS and ductal hyperplastic lesion without atypia remained stable in the same period.

DOI: 10.29289/259453942018V28S1067

IMPACT OF TAMs IN TUMOUR MICROENVIRONMENT AS PROGNOSTIC MARKERS IN BREAST CANCER

Luciana V Q Labre¹, Vera A Saddi^{2,3,4}, Aline C Batista⁵, Juliana F Paes², Jessica E P Ramos^{2,3,4}, Cesar A S T Vilanova-Costa⁶, Erika C Aquino¹, Silvia H Rabelo-Santos^{1,7}

¹Instituto de Patologia Tropical e Saúde Pública - Universidade Federal de Goiás – Goiânia (GO), Brazil.

²Programa de Pós-graduação em Ciências da Saúde, Universidade Federal de Goiás – Goiânia (GO), Brazil.

³Laboratório de Diversidade Genética, Pontifícia Universidade Católica de Goiás – Goiânia (GO), Brazil.

⁴Laboratório de Oncogenética e Radiobiologia, Hospital Araújo Jorge, Associação de Combate ao Câncer em Goiás – Goiânia (GO), Brazil.

⁵Faculdade de Odontologia, Universidade Federal de Goiás – Goiânia (GO), Brazil.

⁶Laboratório de Biologia Tumoral e transplante de Medula Óssea, Hospital Araújo Jorge, Associação de Combate ao Câncer em Goiás – Goiânia (GO), Brazil.

⁷Faculdade de Farmácia, Universidade Federal de Goiás – Goiânia (GO), Brazil.

The objective of this study was to relate the presence of tumor-associated macrophages (TAMs) from 82 breast cancer cases with clinico-pathological characteristics of breast cancer cases and to assess the impact of TAMs on the tumor as a possible prognostic marker. The selection of the cases was made from the search of anatomopathological records in a reference center for the treatment of cancer. We included confirmed cases of diagnosis of invasive ductal breast cancer, with clinical follow-up for five years. The laminas were subjected to immunohistochemical analysis with monoclonal antibody TAMs (CD68). In the statistical analyzes, the cases were classified according to the semi-quantitative evaluation of cell marking as low infiltration or high infiltration and according to the mean. Elevated levels of TAMs (CD68+) were significantly correlated with patient age, presence of distant metastasis, more advanced tumor staging, absence of estrogen receptors (ER) and progesterone (PR), and triple negative breast cancer (TNBC). The survival of the cases of breast cancer with larger TAM infiltrate was reduced corroborating with the other results found. The number of CD68 positive cells represented a strong independent prognostic factor. The presence of TAMs in this study served as an indicator of a worse prognosis among women with the TN phenotype. The high infiltration of TAMs significantly impacted prognostic factors relevant to cases of women with breast cancer.

DOI: 10.29289/259453942018V28S1068

M2-TYPE MACROPHAGES IN TUMOUR MICROENVIRONMENT AS PROGNOSTIC MARKERS IN WOMEN WITH BREAST CANCER

Luciana V Q Labre¹, Vera A Saddi^{2,3,4}, Megmar A S Carneiro¹, Aline C Batista⁵, Diego A C Arantes⁵, Jessica E P Ramos^{2,3,4}, Erika C Aquino¹, Silvia H Rabelo-Santos^{1,6}

¹Instituto de Patologia Tropical e Saúde Pública, Universidade Federal de Goiás – Goiânia (GO), Brazil.

²Programa de Pós-graduação em Ciências da Saúde, Universidade Federal de Goiás – Goiânia (GO), Brazil.

³Laboratório de Diversidade Genética, Pontifícia Universidade Católica de Goiás – Goiânia (GO), Brazil.

⁴Laboratório de Oncogenética e Radiobiologia, Hospital Araújo Jorge, Associação de Combate ao Câncer em Goiás – Goiânia (GO), Brazil.

⁵Faculdade de Odontologia, Universidade Federal de Goiás – Goiânia (GO), Brazil.

⁶Faculdade de Farmácia, Universidade Federal de Goiás – Goiânia (GO), Brazil.

The objective of this study was to assess the prognostic value of tumour-associated macrophages (TAMs) with a possible M2-type macrophage phenotype (CD163+) in women with breast cancer. Cases selected among the records of anatomic-pathological examinations carried out at a reference center for cancer treatment. Inclusion of confirmed cases of invasive ductal carcinoma with clinical follow-up for 5 years. The laminas were subjected to immunohistochemical analysis with monoclonal antibody TAMs like M2-type (CD163). For the statistical analyses, the cases were classified according to the mean value of cell tagging as low infiltration or high infiltration. High levels of TAMs (CD163+) were significantly correlated with distant metastases, lack of receptors estrogen (ER) or progesterone receptors (PR) and triple-negative breast cancer (TNBC). A high number of CD163+ cells was a strong independent prognostic factor. High infiltration of CD163+ emerged as a strong independent prognostic factor. Additional markers able to identify patients with more aggressive types of breast cancer may help predict a poorer prognosis.

DOI: 10.29289/259453942018V28S1069

IN VIVO IMAGING USING BIOLUMINESCENCE FOR EVALUATION OF ANTITUMORAL EFFICACY IN A BREAST CANCER MODEL

Raquel S. Faria¹, Luiza I. de Lima¹, Marina S. Franco², Marjorie C. Roque², Mônica C. Oliveira², Ricardo B. Azevedo¹, João Paulo F. Longo¹

¹Genetics and Morphology Department, Institute of Biological Science, Universidade de Brasília – Brasília (DF), Brazil.

²Department of Pharmaceutical Products, Faculty of Pharmacy, Universidade Federal de Minas Gerais – Belo Horizonte (MG), Brazil.

Objective: To evaluate the Antitumor activity of a liposomal nanosystem containing Paclitaxel and Doxorubicina (pHSL-PTX/DXR), by monitoring the tumor cells bioluminescence *in vivo*. **Methodology:** The tumor cell line 4T1 (adenocarcinoma mammary murine – ATCC CRL-2539) was modified using lentivirus particles Lentifact™ (GeneCopoeia, MD, EUA), according to the manufacturer's protocol. The cells were cultivated in DMEM medium, with fetal bovine serum and antibiotics, 10% and 1%, respectively. The orthotopic implants of 4T1-luc (1×10^5) was performed in the left fifth breast of female Balb/c mouse. Throughout the procedure, the animals were kept under inhalation anesthesia with isoflurane (2.5%). The treatment started after 14 days of tumor induction, in the following groups: *pHSL-PTX/DXR*, *pHSL-DXR*; *pHSL-PTX*, *free DXR*, *free PTX* e *free PTX/free DXR*. The intravenous treatments were performed in 4 doses of 100 µL, three days interval. For the bioluminescence images, 100 µL of D-Luciferin (concentration 150 mg.kg⁻¹) (Sigma) was used. The animals were kept under anesthesia in the Lumina XR IVIS equipment, with 1 minute exposure time. The quantification of the photon signal was processed in the Living Image 3.0 program, Caliper Life Sciences, Alameda, CA. **Results:** The *pHSL-PTX/DXR* treated group system showed the smallest sign of bioluminescence during and after treatment, showing a reduction in tumor mass, and consequently better efficacy against primary breast tumor. There was also a reduction of the light signal in the lung region, indicating a decrease in the metastasis process caused by this type of tumor cell. **Conclusion:** using the *in vivo* bioluminescence method it was possible to evaluate the antitumor efficacy of the liposomal nanosystem *pHSL-PTX/DXR*, and the non-invasive monitoring of pulmonary metastases.

DOI: 10.29289/259453942018V28S1070

PAIN IN WOMEN WITH BREAST CANCER

Magno B. Cirqueira¹, Carolina R. de Mendonça¹, Leonardo R. Soares¹, Marise A. R. Moreira², Ruffo Freitas-Júnior^{3,4}¹Postgraduate Program in Health Sciences, Universidade Federal de Goiás– Goiânia (GO), Brazil.²Department of Pathology, Universidade Federal de Goiás – Goiânia (GO), Brazil.³Universidade Federal de Goiás, Mastology Program, Clinics Hospital – Goiânia (GO), Brazil.⁴Goiás Association to Cancer Combat, Araújo Jorge Hospital – Goiânia (GO), Brazil.

Objective: To describe the prevalence of pain in women with breast cancer. **Methodology:** The search for scientific articles was carried out in the databases: PubMed, LILACS and SciELO. Articles published between 2000 and 2018 were selected in English, Portuguese or Spanish. The searches in the databases were performed with the following combinations: (*“Pain”*[MeSH Terms] OR *“Chronic Pain”*[MeSH Terms] AND *“Breast Neoplasms”*[MeSH Terms] OR *“Prevalence”*[MeSH Terms]).

Results: Painful perception in breast cancer patients is multifactorial and presents different prevalence rates. Persistent pain in the breast (duration ≥ 6 months) was reported by 46.5% of breast cancer survivors and 12.7% of women without a history of breast cancer. In Denmark, it was observed that the prevalence of chronic pain after breast cancer treatment ranged from 22% to 53%. In patients submitted to mastectomy, the prevalence of chronic pain may vary between 20 and 68%. Among the justifications for the variation in the prevalence rates of pain in women with breast cancer, the different criteria of pain measurement and the heterogeneity of this population stand out. The occurrence of preoperative breast pain was 27.1%. In the postoperative period, there is also a high prevalence of breast pain. In a prospective study involving 398 women with breast cancer, it was observed that approximately 25% of the sample had significant and persistent levels of breast pain in the first six months after surgery. In another similar study, breast pain after six months of surgery was described by 52.9% of the patients. In another study, three months after surgery, 60 patients (40%) reported persistent pain in the breast, for which 62% used analgesics. It should be noted that this painful perception is usually of musculoskeletal origin and may suffer variations due to the surgical technique used and the length of each procedure. In a study comparing quality of life after axillary lymphadenectomies and sentinel lymph node biopsies in breast cancer patients, ipsilateral shoulder and upper limb pain was observed in 68% and 36% of patients, respectively. **Conclusion:** Breast cancer represents a public health problem, with a high incidence of pain related to the disease or the treatment performed. Strategies to improve persistent breast pain should be explored for incorporation into the standard treatment of breast cancer survivors.

DOI: 10.29289/259453942018V28S1071

IS THERE AGREEMENT BETWEEN EVALUATORS THAT USED TWO SCORING SYSTEMS TO MEASURE ACUTE RADIATION DERMATITIS?

Marceila A. Fuzissaki¹, Carlos E. Paiva¹, Marcelo A. Maia¹, Mariana T. M. Lima¹, Paula P. L. Canto¹, Yara C. P. Maia¹¹Universidade Federal de Uberlândia – Uberlândia (MG), Brazil.

Objective: To analyze the correlation between the Radiation Therapy Oncology Group (RTOG) and WHO (World Health Organization) scales and the agreement between evaluators. **Methodology:** A prospective and longitudinal study conducted during the period of 2016-2017, in a university hospital, including 834 images of irradiated sites of breast cancer (BC) women during radiotherapy (RT). The images were evaluated by three experienced nurses. In order to evaluate the correlation between the RTOG and WHO scales and the agreement between the evaluators, the Spearman correlation *rho* coefficient (ρ) and the *Kripp* alpha ordinal agreement coefficient were used, respectively. The statistical software's used were SPSS version 20.0 and R version 3.2.5. **Results:** Moderate to strong correlations were identified between the two scales (correlation coefficients 0.4 - 0.9). When analyzing the agreement among the evaluators, moderate agreements (RTOG scale: 0.409, 95%CI 0.369–0.438; WHO scale: 0.550, 95%CI 0.509–0.593) were identified and general agreement rates of 10.2% and 28.8%, respectively, were obtained. **Conclusions:** The RTOG and WHO scales should be used with caution in the clinical practice to identify radiodermatitis prevalence and severity. An illustrative scale containing typical cases of radiodermatitis is being designed by our group aiming to improve accuracy and agreement between evaluators that will be tested in subsequent clinical studies.

DOI: 10.29289/259453942018V28S1072

PHYSICAL ACTIVITY AND HEALTH RELATED QUALITY OF LIFE OF BREAST CANCER WOMEN SUBMITTED TO NEOADJUVANT CHEMOTHERAPY

Julia de M. R. Medina¹, Suzana S. de Aguiar¹, Luiz Claudio S. Thuler¹, Marcelo Adeodato Bello¹, Anke Bergmann¹¹Hospital do Câncer III, Instituto Nacional de Câncer – Rio de Janeiro (RJ), Brazil.

Objective: To evaluate the influence of the level of Physical Activity (PA) in the Health Related Quality of Life (HRQOL) of women submitted to neoadjuvant chemotherapy for the treatment of breast cancer (BC). **Methods:** Prospective cohort study in women diagnosed with BC, with neoadjuvant chemotherapy with adriamycin and cyclophosphamide (AC), from April 4, 2016 to August 2, 2017, at Hospital do Cancer III (HCIII / INCA). The patients were submitted to the interview, physical examination and application of questionnaires at the first consultation with the oncologist and after 21 days of the end of the 4th cycle of neoadjuvant chemotherapy with AC. This project was submitted and approved by the Research Ethics Committee (CAAE 51100615.7.0000.5274). The analysis of the population and the simple and adjusted linear regression analysis between the levels of PA and the domains of HRQoL were performed, considering the 95% confidence interval. **Results:** A total of 253 women were included, with a mean age of 50.87 years (\pm SD 10.62). After adjusting, the women with higher levels of PA have a better overall QoL (Beta 4.06, $p=0.040$) after neoadjuvant chemotherapy. There was no statistically significant difference between the AF levels after a neoadjuvant chemotherapy and the HRQoL domains. **Conclusion:** Patients who had higher levels of pre-diagnosis PA had better overall QoL after neoadjuvant chemotherapy and there was no difference between groups when assessing the levels of PA and HRQoL after the AC neoadjuvant chemotherapy regimen.

DOI: 10.29289/259453942018V28S1073

IMPACT OF NUMBER OF POSITIVE LYMPH NODES AND LYMPH NODE RATIO ON SURVIVAL OF WOMEN WITH NODE-POSITIVE BREAST CANCER

Fabiana Tonello¹, Anke Bergmann¹, Karende Souza Abrahão¹, Suzana Sales de Aguiar¹, Marcelo Adeodato Bello¹, Luiz Claudio Santos Thuler¹

¹Hospital do Câncer III, Instituto Nacional de Câncer – Rio de Janeiro (RJ), Brazil.

Purpose: This study aimed to evaluate the association of axillary lymph node ratio (LNR) and number of positive lymph nodes (pN) with the risk of breast cancer recurrence and death. **Methods:** A retrospective cohort study of node-positive stage II e III breast cancer patients diagnosed and treated between 2008 and 2009 at the Brazilian National Cancer Institute (INCA), Brazil. Overall and disease-free survival curves for number of positive lymph nodes (pN) and lymph node ratio (LNR) risk groups were constructed using the Kaplan-Meier method and compared by the log-rank test. Multivariate analysis was performed using stepwise forward Cox regression models. **Results:** In total, 628 women with node-positive breast cancer were included. Most patients (69.5%) had advanced clinical stage tumors (\geq IIB). The median follow-up was 58 months (range: 3-92 months). The adjusted recurrence hazard of pN2 and pN3 patients was 2.47 (95% Confidence Interval [CI] 1.72–3.56) and 2.42 (1.62–3.60), respectively, compared to pN1 patients ($p < 0.001$), while the hazard of intermediate (0.21-0.65) and high-risk (>0.65) LNR was 2.11 (1.49–3.00) and 3.19 (2.12–4.80), respectively, compared to low-risk LNR (≤ 0.20) patients ($p < 0.001$). On the other hand, the hazard of death of pN2 and pN3 patients was 2.17 (1.42–3.30) and 2.41 (1.53–3.78), respectively ($p < 0.001$), and the hazard of intermediate (0.21-0.65) and high-risk (>0.65) LNR patients was 1.70 (1.13–2.56) and 2.74 (1.75–4.28), respectively ($p < 0.001$). **Conclusion:** Higher pN and LNR were associated with shorter disease-free survival and overall survival times.

DOI: 10.29289/259453942018V28S1074

SENTINEL LYMPH NODE BIOPSY AFTER NEOADJUVANT CHEMOTHERAPY IN WOMEN WITH BREAST CANCER: CLINICAL PROFILE AND PROGNOSIS

Marcelo Bello¹, Anke Bergmann¹, Marcelo Barbosa¹, Suzana Aguiar¹, Fabiana Tonello¹, Sergio Monteiro¹, Patricia Juca¹, Luiz Claudio Santos Thuler¹

¹National Cancer Institute – Rio de Janeiro (RJ), Brazil.

Objectives: Describe the frequency and factors associated with the indication of Sentinel lymph node biopsy (SLNB) after neoadjuvant chemotherapy (NACT) and the impact of SLNB on prognosis. **Methods:** Retrospective cohort study of women with breast cancer submitted to NACT from January 2013 to July 2015 at Hospital do Cancer III of the National Cancer Institute (HCIII/INCA). The data were collected in electronic and physical records. This study was approved by INCA's ethics and research committee. **Results:** A total of 783 women with mean age of 52 years (± 11) were included. The majority were diagnosed with clinical stage IIIA (23%) and IIIB (33%), and with a Luminal A subtype (ER+ and / or PR+, HER2-) (52%). After NACT, complete response was observed in 15% and partial in 58%. Mastectomy was performed in 85% of the women, and 25% were submitted to SLNB (alone 14% and SLNB + Lymphadenectomy 11%). The factors associated with SLNB were: Initial clinical stage (OR = 6.22 95%CI 4.28–9.06) and complete response to NACT (OR=1.96 95%CI 1.29–2.97). In the follow-up, were observed 23% of recurrence /metastasis and 13% of death. After adjustment (cT, cN and NACT response), there was no association between the performance of SLNB with recurrence/metastasis (HR=1.51 95%CI 0.93–2.46) and death (HR=1.48 95%CI 0.76–2.90). **Conclusion:** The indication of SLNB after NACT was associated with initial clinical stage and complete response to NACT. After adjustment, SLNB was not associated with prognosis.

DOI: 10.29289/259453942018V28S1075

SURVIVAL IN PATIENTS WITH STAGE IV HER2 POSITIVE BREAST CANCER

Marcelo A. Bello¹, Marcelo M. Barbosa¹, Marcela E. Carvalho¹, Sergio Monteiro¹, Patricia Juca¹, Anke Bergmann¹

¹Hospital do Câncer III, Instituto Nacional de Câncer – Rio de Janeiro (RJ), Brazil.

Objective: To evaluate the survival of patients with metastatic breast tumors, with human epidermal growth factor (HER2) receptors, present and treated in Unit 3 of the National Cancer Institute. **Methodology:** Retrospective cohort study with women diagnosed and treated in Unit 3 of the National Cancer Institute, in the years 2008 and 2009 (n=2035). For this analysis, patients with HER2 positive at diagnosis, clinical staging I to III (n=412) and who progressed with metastasis in the follow-up period (n=92) were selected. In this group of patients, 15.2% did herceptin palliative and 84.8% did not. **Results:** Among the 92 positive Herb 2 patients who progressed with distant metastasis, 32.6% had bone or soft tissue, 62% visceral and 5.4% locoregional metastasis, respectively. The mean survival of the patients was 38.4 months. Among those taking Herceptin, the mean survival was 49.7 months, while those who did not were 35.9 months. Among 31 patients with bone metastasis, enrolled between 2007 and 2009, 5 (16%) did Herceptin and 31 (84%) did not. The mean survival was 51.3 months and 31.7 months, respectively. In general, those patients who did not take Herceptin had a 2.35-fold higher risk of death compared to those who did (HR=2.35 95%CI 1.12–4.95, p=0.024). **Conclusion:** The use of trastuzumab in the palliative treatment of Herb2 positive patients had an impact on survival in the present study, including isolated bone metastases.

DOI: 10.29289/259453942018V28S1076

PROFILE OF MAMMOGRAPHIES REALIZATION IN BRAZIL BETWEEN 2013 AND 2014

Lara C. R. Alvarenga¹, Luís M. M. Medeiros¹, Mateus F. O. Vilela¹, Bráulio B. Rodrigues¹, Isadora G. C. K. Severino¹, Júlia R. Moraes¹, Mirian P. Silva¹

¹Centro Universitário de Anápolis (UniEVANGÉLICA) – Anápolis (GO), Brazil.

Objective: The objective of this study is to outline the profile of patients who undergo mammography in the Brazilian scenario between the years of 2013 and 2014. **Methodology:** It is a retrospective and quantitative study presenting a cross-sectional approach carried out in Brazil between the years of 2013 to 2014. It was based on the general female population who underwent mammography in this selected period. Data were obtained from the Department of Informatics of the Unified Health System (DATASUS), of secondary order, in the database category of the Breast Cancer Information System (SISMAMA). The age of the women studied, their region of origin, the results found in this exam, as well as their final conduct according to each case were also used as variables. **Results:** During the analysis period of the study, 4,041,101 mammograms were performed in women aged less than 10 years older to 70 in Brazil. Of these, the prevalence among those aged between 50 and 54 years is responsible for approximately 20% of the total number of exams. In addition, most mammograms were performed in the Southeast region, followed by the Northeast, South, Central - West and North regions of the country. According to the results obtained in the total number of mammograms performed, 1,930,433 were classified as Bi-RADS 2, and 50,921 were suspected or highly suggestive of malignant breast neoplasm. Among the latter, the prevalent age group was those older than 70 years. Regarding the conducts for mammograms performed, 3,442,629 women were instructed to repeat the examination every 2 years, 50,921 were biopsied and only 2,689 were referred for specific therapy. **Conclusion:** It is known that breast cancer is the most prevalent cancer in women all over the world and thus its screening and early diagnosis is necessary. One of the main risk factors is age, justifying the highest number of mammograms among women over 50 years of age, as well as their highest occurrence in the elderly. Also, expanding screening in regions such as the Midwest and North regions to women with risk factors and clinical symptoms is essential to know the neoplasia in its earliest stages and to obtain better prognosis in the ducts.

DOI: 10.29289/259453942018V28S1077

RESOURCES USED BY THE SUS WITH RESECTION OF NON - PALPABLE BREAST CODES FROM 2015 TO 2017

Luís M. M. Medeiros¹, Mateus F. O. Vilela¹, Bráulio B. Rodrigues¹, Isadora G. C. K. Severino¹, Júlia R. Moraes¹, Lara C. R. Alvarenga¹, Mirian P. Silva¹

¹Centro Universitário de Anápolis (UniEVANGÉLICA) – Anápolis (GO), Brazil.

Objectives: To evaluate the expenditures to the health system with the non-palpable breast node resection procedure in the Unified Health System (SUS) between the years 2015 and 2017. **Methodology:** This is a retrospective and quantitative study with an approach by cross-sectional study carried out in Brazil between the years 2015 and 2017. The sample population was the Hospital Information System (SIH) of SUS with data obtained from the DATASUS platform, of secondary order. The year of the procedure, the emergency or elective care, the region of the country and the average hospitalization value, the total value per year, the days of hospitalization and the public or private nature were used with variables. **Results:** Between 2015 and 2017 a total of 17,703,833 million reais were spent on resection of non-palpable breast lesions with oncology marking, and the year 2017 represented the highest percentage of expenditures, with 34 % of total value. The years 2015 and 2016 represented similar but smaller expenditures, with 32.5% and 33.4%, respectively. In addition to being the year in excess of expenses, 2017 was also the year in which there was a higher hospital admission authorization (AIH), with 4,880, representing 33.8%, while the other years were 32.5% in 2015, and 33.6% in 2016. According to the AIH, the cost for each hospitalization has an average value of 1878.15 reais per patient who needs the procedure. Among the regions of Brazil, the highest number of AIH approved in the Southeast Region is 44.11%, followed by the Northeast Region, with 30.9% and the South Region, with 22.11%. The Midwest and North regions totaled 1.6% and 1.1%, respectively. Most of the procedures were carried out on an elective basis and little in an emergency, representing 81.8% for the first option and 18.2% for the second. Regarding the payment scheme, 73.5% of the procedures are in the category ignored, while a small amount, 3.4%, were in the private regime and 23% in the public regime. **Conclusion:** It is noticed that breast cancer is still a neoplasia present in the population due to the large number of procedures that are performed for the reversal and improvement of the patient's condition. Still, it is a pathology that demands large expenditures to the Brazilian public and private health system, with coverage in all regions of the country, however, with emphasis on the Southeast Region, where there was a greater quantity of AIH approved, while the North Region exhibited the less quantity.

DOI: 10.29289/259453942018V28S1078

ANALYSIS OF THE CLINICAL INDICATIONS FOR MAMMOGRAMS IN BRAZIL FROM 2013 TO 2014 BY THE UNIFIED HEALTH SYSTEM

Diogo T. de Lima¹, Gabriela F. Araújo¹, Fábio F. Marques¹, Guilherme N. Correa¹, Mateus F. O. Vilela¹, Bráulio B. Rodrigues¹, Mirian P. Silva¹

¹Centro Universitário de Anápolis (UniEVANGÉLICA) – Anápolis (GO), Brazil.

Objective: To elaborate an analysis of the character of mammograms performed in Brazil between 2013 and 2014 by the Unified Health System (SUS) in relation to the indication profile and the patients submitted to the examination in this period. **Methodology:** The data from the study originated from the Department of Informatics of the Unified Health System (DATASUS), in the database category of the Breast Cancer Information System (SISMAMA). It is based on a retrospective and quantitative cross-sectional study done in Brazil between 2013 and 2014 with a population composed of women of all age groups who underwent mammography during the evaluation period. Was used as analysis variables: the clinical indications of the exam, the age groups of the patients involved and their declared race. **Results:** In the study period, 4,041,101 mammograms were performed, of which 2,315,907 were in women aged 50–54 years. Of the races analyzed in the survey, only 642 women declare themselves indigenous, in contrast, 2,304,282 have relinquished this information. Of the total number of tests performed, 97.44% had a screening character, while 2.56% were intended for diagnosis. Women between the ages of 45 and 50 predominated with regard to diagnostic mammograms in the year in question, while those with 50-54 years of age were the majority on screening mammograms. Regarding the clinical indication of the test, 931,994 women had no prior clinical examination with a complementary indication, and 1,857,189 had never performed previous mammograms. **Conclusion:** Strategies for the early detection of breast cancer are based on premature diagnosis in symptomatic patients or screening in those who do not present symptoms. Routine physical examination by a health care professional as well as a detailed medical history are essential to perform the correct indications for complementary examinations and careful evaluation of the risk factors related to the neoplasia, and thus to reduce their false-positive and iatrogenic results caused by them. Under SUS, therefore, it is imperative to perform the basic physical examination of the breast to avoid unnecessary expenses with other methods of tracking.

DOI: 10.29289/259453942018V28S1079

THE PREVALENCE OF MALIGNANT AND BENIGN BREAST NEOPLASMS IN BRAZIL BETWEEN 2010 AND 2014

Fábio F. Marques¹, Guilherme N. Correa¹, Mateus F. O. Vilela¹, Bráulio B. Rodrigues¹, Diogo T. de Lima¹, Gabriela F. Araújo¹, Mirian P. Silva¹

¹Centro Universitário de Anápolis (UniEVANGÉLICA) – Anápolis (GO), Brazil.

Objective: To compare the prevalence of benign and malignant breast neoplasms in Brazil between 2010 and 2014. **Methodology:** This is a quantitative, population-based, cross-sectional study in Brazil between 2010 and 2014. A study population composed of cases of mammary neoplasia was used. Data were obtained from the DATASUS system, of secondary order, in the category of database in the Hospital Information System of SUS (SIH/SUS). It was used as variable year, sex, age, type and character of care. **Results:** In Brazil, between 2010 and 2014, benign and malignant breast neoplasms total 267,038 cases, with a progressive increase between the years. Half of the cases are concentrated in the southeast region, with 50.12%, followed by the northeast region, 21.70%, southern region, 19.84%, center-west, 5.41% and north, 2.89%. Of the total, 92.25% are malignant neoplasms of the breast. The female sex prevails over the male in the number of cases, with 98.34% versus 1.65%. In relation to age, there is a distribution in all age groups, prevailing in the larger ranges, such as 50 to 59 years, 26.55%, 40 to 49 years, 24.86%, 60 to 69 years, 18.95%, 30 to 39 years, 10.41%, 70 to 79 years, 9.91%. Many of the attendances are elective, 66.07%, and others as a matter of urgency, 33.92% and 64.89% are from the private network versus 35.10% of the public network. **Conclusion:** Breast neoplasms are very prevalent in Brazil and most of them are malignant. The southeastern region of Brazil presents the largest number of cases together with the female sex, but it is emphasized that they exist in the male sex. The most prevalent age groups are those over 30 years of age, but this does not prevent them from being found in the lower ranges, such as 20 to 29 years. Therefore, it is necessary to invest more in medical qualification programs, to rank hierarchically and to guarantee quick access to the secondary and tertiary centers for resolute care, providing the tracking of the disease, aiming in the medium term a greater reduction of cancer mortality.

DOI: 10.29289/259453942018V28S1080

HORMONE THERAPY IN TRANSSEXUALS AND THE RISK OF BREAST CANCER

Gabriela F. Araújo¹, Fábio F. Marques¹, Guilherme N. Corrêa¹, Mateus F. O. Vilela¹, Bráulio B. Rodrigues¹, Diogo T. de Lima¹, Mirian P. Silva¹

¹Centro Universitário de Anápolis (UniEVANGÉLICA) – Anápolis (GO), Brazil.

Objective: To carry out a systematic review of the literature on the relation of breast cancer in transsexual women using hormone therapy. **Methods:** Systematic review of the literature, using the PubMed databases, between 2003 and 2016, using the key words “breast cancer”, “female-to-male transsexuals”, “male-to-female-transsexuals”, “Hormone therapy”. From the total of 22 articles found, a total of 15 articles were selected in the last 13 years. Six studies were excluded because they were not the morphological alterations of the use of hormonal therapy in transsexuals, as well as the possibility and occurrences of breast cancer in these. **Results:** The nine articles selected show the relationship between hormone therapy and breast cancer in transsexuals, taking into account that in all the cases reported, the patients had firstdegree relatives with a history of breast, ovary or prostate cancer. However there are differences between studies. *Lancet Oncol.* 2013 September sees strong association of breast cancer with serum concentration of testosterone. It discusses the action of testosterone to be aromatized in estrogen and stimulate the direct activation of estrogen receptors, exposed in two cases reported by Shao T, et al. in which in both the expression of estrogen receptors in transsexuals using testosterone presented in 90%. While in a study by SLAGTER, Margarita H. et al. it is stated that testosterone promotes apoptotic effects in cancerous cells of breast cancer. Therefore, the simultaneous presence of high circulating testosterone and estrogens in these subjects could have prevented the biological action of E2, causing a histological image of involution. **Conclusion:** Although not yet studied, when referring to transsexuals from woman to man in use hormone therapy with high testosterone levels, strongly associated with positive family history, risks can not be excluded. As well as the presence of the BRCA2 mutation in transsexuals from man to woman using antiandrogen and estrogen may lead to breast cancer. A full family history is important in evaluating transgender people seeking hormone treatment.

DOI: 10.29289/259453942018V28S1081

NEW LABORATORY TESTS FOR DIAGNOSIS AND PROGNOSIS OF BREAST CANCER

Guilherme N. Corrêa¹, Mateus F. O. Vilela¹, Bráulio B. Rodrigues¹, Diogo T. de Lima¹, Gabriela F. Araújo¹, Fábio F. Marques¹, Mirian P. Silva¹

¹Centro Universitário de Anápolis (UniEVANGÉLICA) – Anápolis (GO), Brazil.

Objectives: To identify new laboratory tests for prognosis and diagnosis of breast cancer. **Methodology:** Literature review, with search of articles in the databases PubMed and Medline, using the descriptors: new laboratory tests, breast cancer, prognosis, diagnosis. A total of 75 articles were selected, of which 41 were adequate for the review research. The inclusion criteria were more recent articles and of high theoretical value for this review. **Results:** The laboratory tests identified can be divided into three groups: the first is those that use computer engineering to form images, three-dimensional models of the breast and to track tumor cells. In the second are those that investigate gene and epigenic expressions. In the third group we have histochemical tests of specific substances associated with breast cancer. In the first group, there are several applications of computer technology, such as CellSearch System software and EPISPOT, which use specialized microscopy to identify tumor cells in breast tissue biopsies. Another example of joint work between pathology and computation are high-resolution imaging exams such as the US-CNBs, a highly-defined ultrasound examination, the ED-CSCT, an examination that uses X-ray scatter to calculate the type and the size of microcalcifications that can occur in breast cancer. Computer technology also enabled the creation of a three-dimensional model of the breast, enabling the physician to accurately visualize the tumor in the midst of tissue layers. In the second group, gene expression screening focuses on joint monitoring, observing the traditional genes, such as BRCA1/BRCA2 and Ki67, and other associated ones, such as the CA72-4 carcinogen antigen and the GRP94 and FN14 biomarkers, since their expression is directly associated with breast cancer. In addition to the genetic markers, epigenetic studies revealed a correlation between methylation of the RASSF1A and CDH1 gene promoters and the prognosis of metastasis in breast cancer. In the third group are improved histochemical tests, which guarantee greater accuracy and specificity, such as the use of chromogenic and fluorescent methods in the evaluation of the estrogen receptor, the automated use of the FISH protocol to reduce tumor cell damage, avoiding sample loss, the analysis of the organizing regions of the nucleus of a tumor cell, the change in the technique of histochemical staining for a better visualization of the endothelial growth factors 2. It is also worth noting the elaboration of new protocols for exams, such as for the analysis of the factor gene of endothelial growth 2. **Conclusion:** The evolution of technology allowed not only the creation of more specific exams but also the improvement of old techniques. In this way, there are now tools that help the physician in the accurate diagnosis and prognosis of breast cancer, allowing a more specific visualization of the health-disease process, that is, the tests allow to identify the uniqueness of the pathology that affects the individual, either by more accurate imaging tests, either by enhancement and identification of new biomarkers, or by enhancement of old histochemistry techniques. These are factors that allow identifying the characteristics of the pathology that affects the individual, increasing the possibilities of diagnosis, treatment and prognosis by the physician.

DOI: 10.29289/259453942018V28S1082

INCIDENCE AND MORTALITY OF BREAST CANCER IN WOMEN BY AGE GROUP IN BRAZIL AND LATIN AMERICA

Adila C. S. de Sousa¹, Carolina B. dos S. Azevedo¹, Lázaro L. M. de Oliveira¹, Matheus L. T. Costa¹, Sarah R. Santos¹, Thaynara Mercadante¹

¹Universidade Federal de Goiás – Goiânia (GO), Brazil.

Objective: Analyse the Latin America data on female breast cancer, comparing by age group the general rates of incidence and mortality with the Brazilian ones. **Methodology:** epidemiological analytic study carried out about the interpretation of data available in the GLOBOCAM stand, referring to the incidence and mortality of breast cancer in South America compared to Brazil. **Results:** There were 67.316 cases of breast cancer in the age group 0 to 74 years, of which 16.412 died, representing a 24% mortality rate. It was possible to identify through data analysis that the highest incidence of breast cancer is in the age group of 50–59 years, where it presented an incidence of 26%, however the highest mortality rate was seen in the age group over 70 years, with a percentage of 29%. In South America, excluding Brazil, it was possible to perceive that the highest incidence of breast cancer is in the age group above 70 years of age, presenting a percentage of 25% of the 48.398 cases. **Conclusion:** From these data it is possible to conclude that the mortality rate does not follow the same age pattern of the incidence rate, that is, there are variations according to age. Such variations reveal that the higher the age group, the higher the mortality rate. This can be caused by the fact that cancer, in this age group, presents a histological grade of higher malignancy, either by a late detection or by the patient's advanced age. Thus, early diagnosis is of great importance for reducing mortality rates and intervening in factors that influence detection is fundamental, such as: accessibility to the health service, coverage of mammography and quality of treatment.

DOI: 10.29289/259453942018V28S1083

BREAST PAIN/MASTALGIA DUE TO A FOREIGN BODY IN A PSYCHIATRIC PATIENT: CASE REPORT

Lucas N. da Silva¹, Luis F. P. Oliveira¹, Matheus L. T. Costa¹, Taynara Mercadante¹¹Universidade Federal de Goiás – Goiânia (GO), Brazil.

Objectives: To report a case of indication of bilateral mastectomy in a psychiatric patient with breast pain/mastalgia due to a foreign body. **Methodology:** Case report obtained at HC-UFG. The data were collected from medical records and the findings. **Results:** In 02/12/17, during an opportunistic screening campaign for breast cancer, executed by the Mastology Program from Advanced Center for Breast Diagnosis and Academic League of Breast, conjoined with the Municipal Health Secretary from Rubiataba, the patient M.V.S., female, 43 years, birthplace Rubiataba, was examined and referred to the HC-UFG Mastology Department to evaluate the indication of total adenomastectomy. The patient refer that 3 years ago, after her husband death, she developed a severe depressive disorder, which is currently in remission due the treatment. The patient, M.V.S, reports that 3 months later she suffered from sexual abuse that evolved to a bilateral breast pain/mastalgia. During the screening campaign the mammography performed on 29/11/17 was analyzed and evidenced 22 needles in the right breast and 24 in the left one. The patient says that the needles were inserted by the abuser, however the family claims that the patient inserted by herself. In 18/12/17 The Mastology Department requested a psychiatric evaluation, which of was diagnosed with slight intellectual disability, limit intelligence, depression in remission, and humor disorder, without psychiatric contraindication to the chirurgic procedure. Then, the Mastology Department discussed the therapeutic proposal of bilateral adenomastectomy for the excision of the injured tissue and foreign body and the possibility of reconstruction with saline or silicone prostheses. The patient is following up with Tamoxifen 10mg for relieve of breast pain/mastalgia and the service still analyzing the viability of the reconstruction with the breast prosthesis, considering new insertions of foreign body can generate even more damages.

DOI: 10.29289/259453942018V28S1084

SEXUAL BEHAVIOR OF WOMEN WITH BREAST CANCER AND IMPACT ON THE COUPLE'S LIFE

Mayara M. P. Carvalho¹, Ruffo de Freitas Júnior^{1,2}¹Universidade Federal de Goiás (UFG) – Goiânia (GO), Brazil.²Centro Avançado de Diagnóstico da Mama, UFG – Goiânia (GO), Brazil.

Sexuality is experienced by individuals from their birth until old age, and is related to their history and culture. Sex, a constituent factor of human physical and emotional health, derives from an intrinsic characteristic of humans regardless of their social status, and changes according to the cultural and historical context. Sexual activity is one of the main pillars of individuals' sexuality and well-being, and not only associated with reproduction. The emotional aspects related to sexual dysfunction in people with cancer are common and frequent, and may occur since the diagnosis. Breast cancer is the second most common type of cancer worldwide, the most frequent in women, and the leading cause of death in patients with malignant neoplasms. In the state of Goiás (Brazil), the estimated number of new breast cancer cases in 2017–2018 is 49.51 per 100,000 women. In general, breast cancer tumors present morphological, clinical, and biological heterogeneity, due to different histological grades as well as molecular differences. Breast cancer may have multifactorial impact on the quality of life, sexual relationships, and, consequently, marital relationships of patients. Furthermore, the disease can interfere with patients adherence to endocrine therapy. Breast cancer staging may determine the level of sexual dysfunction in women. At early stages, the most reported symptoms are loss of libido and lack of sexual desire, and at later stages, loss of interest in sex and pain during sexual intercourse. Partners' support and understanding are fundamental for women's psychological acceptance of the disease. However, the partners of women with breast cancer also have psychological impacts, which can lead to premature ejaculation, erectile dysfunction, and depression and anxiety disorders. **Objective:** Verify whether the sexuality of both women diagnosed with breast cancer and their partners is affected and the possible causes. **Methodology:** This is a narrative review. The literature review was based on the strategic planning for the search that included the following steps: 1. Constructing a guiding question that explains the interest of the research; 2. Identifying the elements of the research (variables, studied populations, comparisons); 3. Constructing search strategies using keywords (descriptors); 4. Applying the search strategies developed for this study to the articles found in the PubMed Central (PMC) at the U.S. National Institutes of Health's National Library of Medicine (NIH/NLM); 5. Defining search filters such as articles published in the last 5 years, research carried out with humans, with adults; 6. Exclusion criteria: homosexual couples, quality of life, breast cancer in men, and studies that assessed sexuality in single women. The search strategy applied was: cancer treatment OR female breast cancer AND sexuality OR sexual health AND partner OR couple. The filters were: last 5 years and studies with humans. After applying the search strategy and identifying the studies of interest, the first selection of the material was performed analyzing the titles, followed by the selection of the abstracts, and only at the last level of selection the complete articles were identified and read to compose this narrative review. **Results:** In this narrative review, 150 articles were initially collected and examined. After applying the exclusion criteria, 54 were selected. The authors of the articles included in this review unanimously reported that women diagnosed with breast cancer had impairments regarding their sexual performance, due to physiological, therapeutical, and/or emotional changes associated with the neoplasia since the diagnosis. Breast cancer treatment may cause a decline in couples' quality of sexual life. Depression, anxiety, conflicts in the relationship, and low self-esteem can intensify sexual dysfunction in women treated for breast cancer, and this impairment may last many years after the treatment. Mastectomy may substantially aggravate sexual intercourse during and after treatment due to sequels, which may affect couples' relationships. **Conclusion:** Current therapeutic approaches for breast cancer can considerably affect women's sexual performance due to their adverse effects. Therefore, the negative aspects of dealing with psychological and emotional suffering can be potentially aggravated in women diagnosed with breast cancer. Aiming to minimize this suffering, we propose that psychological and medical interventions be offered to women with breast cancer and their partners, both individually or to the couple, during and after the treatment. However, it is of paramount importance that healthcare professionals be attentive to the slightest signs of sexual dysfunction and recommend psychological follow-up.

DOI: 10.29289/259453942018V28S1085

BREAST CANCER SCREENING AMONG BRAZILIAN WOMEN: AN INTEGRATIVE REVIEW

Mônica O. B. Oriá¹, Camila T. M. Vasconcelos¹, Marli T. G. Galvão¹, Aubrey L. Doede², Dan Wilson², Reanna Panagides², Emma M. Mitchell²

¹Universidade Federal do Ceará – Fortaleza (CE), Brazil.

²University of Virginia – Charlottesville (VA), EUA.

Objectives: The aim of this integrative literature review was to explore empirical literature surrounding breast cancer screening practice among Brazilian women. **Methodology:** OVID Medline, CINAHL, Lilacs, Scopus and Web of Science/SciELO were used to identify articles in English, Portuguese and Spanish languages. Twenty eight original articles were included using inclusion and exclusion criteria. **Results:** Most of the studies were developed in the Southeast (n=14) and South (n=7). No studies were found involving populations from North Brazil. The most studies enrolled populations from exclusively urban areas (n=21). Of the total of 91,207 women involved in the 28 studies, 15,801 never had mammograms and 48,944 had mammograms in their lifetime; 555 had never undergone breast clinical examination, and 8,114 had done it at least once in their lives. To screen breast cancer the studies used mammograms (n=21), breast clinical examination (n=11) and mobile mammograms (n=1). White women, with higher schooling, better socioeconomic status have access to better ways of breast cancer screening. **Conclusion:** The gaps in breast cancer screening are related to social and racial inequities and problems about spatial distribution of specialized equipment resources and professional.

DOI: 10.29289/259453942018V28S1086

MICROINVASIVE CARCINOMA INITIALLY LABELED AS *IN SITU*

Ruffo Freitas-Junior¹, Nayara A. F. Lemos², Nilceana M. A. Freitas³, Marise A. R. Moreira⁴, Edésio Martins², José Carlos de Oliveira⁵, Carleane M. B. Silva⁵

¹Breast Program, Universidade Federal de Goiás (UFG) – Goiânia (GO), Brazil.

²Health Science Post-graduate Program of the Medical College, UFG – Goiânia (GO), Brazil.

³Department of Radiotherapy, Araújo-Jorge Hospital, Association for the Combat of Cancer – Goiânia (GO), Brazil.

⁴Department of Medicine, UFG – Goiânia (GO), Brazil.

⁵Goiânia Population-Based Cancer Registry – Goiânia (GO), Brazil.

Carcinoma microinvasors are frequently found in the records of ductal carcinoma *in situ* (DCIS) cases, a factor that may favor non-effective treatment. Survival differs in cases of microinvasion due to the possibility of metastasis. **Purpose:** was to characterize cases of microinvasive carcinoma transcribed as *in situ* and subsequent comparison of survival of these cases in relation to cases of DCIS. **Methods:** We used a secondary database of the Registry of Cancer Population Base of Goiânia and data of the city hall of that city. **Results:** The mean size of the microinvasive lesions associated with DCIS was 1.3 cm, with foci of invasion <1.0 mm, the majority of the positivity for estrogen and progesterone receptors corresponded to 28.6% and for HER2 14, 3%. In addition, 33% of these cases had impaired margins. **Conclusion:** attention different from DCIS should be given to microinvasive carcinoma with regard to registration and diagnosis.

DOI: 10.29289/259453942018V28S1087

PHOTOBIO-MODULATION IN THE SCARIFICATION OF DEHISCENCE AFTER MASTECTOMY: CASE REPORT

Nayara A. F. Lemos^{1,2}, Ruffo Freitas-Junior¹, Fernanda D. Moraes^{2,3}, Arethuzza A. Moreira⁴, Karla F. Carmo⁵, Alexandre L. Cardoso⁵, Eliceia M. Batista⁶

¹Universidade Federal de Goiás – Goiânia (GO), Brazil.

²Universidade Paulista – Goiânia (GO), Brazil.

³Faculdades Objetivo – Goiânia (GO), Brazil.

⁴Associação de Combate ao Câncer de Goiás, Hospital Araújo Jorge – Goiânia (GO), Brazil.

⁵Faculdade Cambury – Goiânia (GO), Brazil.

⁶Hospital das Clínicas de Goiânia – Goiânia (GO), Brazil.

The effects of photobiomodulation on healing occur through the influence of chemical mediators, migration of inflammatory cells and increase of growth factors. Studies emphasize the efficacy in different animal models, however, dosimetric parameters are discrepant. Therefore, the objective of the study is to show a therapeutic option for the treatment of this wound to optimize the other stages of the treatment. This case report refers to a 40-year-old female patient diagnosed with mammary carcinoma followed by confirmation of gestation. The patient had gestational diabetes and on the day of delivery underwent modified radical mastectomy and right axillary lymphadenectomy that evolved with a dehiscence near the axillary cavus. In the evaluation, the patient's main complaint was related to poor healing of the surgical wound and impediment to the start of radiotherapy. A photographic record of the region was carried out, besides a qualitative evaluation of the dehiscence that presented hyperemic, edemaciated and retracted borders. In the main complaint, two physical therapy sessions were performed with a light emitting diode (LEDs) at two wavelengths 470 nm (blue) and 630 nm (red) at 5J / cm² and the 19 seconds and 1:30 minutes respectively. Resolution was observed in the surgical dehiscence. Photobiomodulation proves to be an important resource for the treatment of dehiscence after mastectomy.

DOI: 10.29289/259453942018V28S1088

ANALYSIS OF THE DETECTION OF MALIGNANT BREAST CANCER LESIONS IN WOMEN IN THE STATE OF GOIÁS IN THE PERIOD 2010-2013

Paula de O. C. Queiroz¹, Nayara P. A. Oliveira¹, Fernando C. R. Pinheiro¹, Vitória N. Macedo¹, Hygor A. P. Dutra¹

¹Universidade Federal de Goiás – Goiânia (GO), Brazil.

Objective: To analyze the detection of malignant lesions of breast cancer, through the following indicators: clinical breast examination and mammography, in women in the state of Goiás from 2010 to 2013. **Methodology:** This is a descriptive study using data from the Breast Cancer Information System (SISMAMA). **Results:** It was observed that in the period of 2010 the detection of malignant lesions by clinical examination overlapped the imaging method (mammography) in 17.88%. In 2011, this difference increased to 24.23%. Following the same trend, this difference increased to 53.53% and 59.83%, in the years of 2012 and 2013, respectively. **Conclusion:** The results demonstrate that the detection of malignant lesions by clinical examination in the breast overlaps mammography, evidencing a failure in the early diagnosis of breast cancer in Goiás. Some limitations in adherence to mammography screening are misinformation about the examination by the target public, the lack of training of health teams and the lack of machines in Brazil associated with a poor distribution of machinery. Since mammography is the only method capable of reducing breast cancer mortality in an absolute way, it is fundamental to implement strategies to cope with the low adherence to this imaging examination.

DOI: 10.29289/259453942018V28S1089

COMPARATIVE ANALYSIS OF THE INCIDENCE AND MORTALITY OF BREAST CANCER IN EUROPE AND LATIN AMERICA AND THE CARIBBEAN IN 2012

Paula O. C. Queiroz¹, Larissa S. Valadares¹, Pedro R. Soares¹, Luisa R. Barros¹

¹Universidade Federal de Goiás – Goiânia (GO), Brazil.

Objective: To compare the incidence and mortality of breast cancer in Europe and the countries of Latin America and the Caribbean in 2012. **Methodology:** An ecological analytical study carried out by the interpretation of the data available on the GLOBOCAN platform, concerning the incidence and mortality of breast cancer in Europe and in the countries of Latin America and the Caribbean in 2012. **Results:** In Latin America and the Caribbean, the incidence of breast cancer, in absolute numbers, is 152,059, while the mortality rate is 43,208. In Europe, the incidence is higher, 458,718, with a mortality of 131,347. Incidence and mortality rates are higher in Europe than in Latin America and the Caribbean: the incidence rate in the former is 119.5 / 100.000 inhabitants and in the second it is 49.8 / 100.000 inhabitants, while the mortality rate in the former is 34.2 / 100,000 inhabitants, while in the latter it is only 14.1 / 100,000 inhabitants. If, on the other hand, we compare the incidence / mortality in women, by age group, in both regions, we can conclude that these data are always higher in the group aged 40 years or older. In Europe, the incidence of breast cancer in women aged 40 years or older is 94.55%, while mortality in the same group represents 97.8% of the total. On the other hand, the incidence of breast cancer in Latin American and Caribbean women aged 40 years or older is 89.4%, while mortality in this age group corresponds to 92.9% of the total. **Conclusion:** Breast cancer accounts for 25 percent of all breast cancer cases in the world and is also the leading cause of cancer-related death in this population, according to the WHO. In Latin America, it is the second leading cause of death among women. Although developed regions have higher incidence rates, an overall increase in the incidence of breast cancer is seen each year as a result of encouraging screening and early diagnosis, two strategies that seek to reduce the mortality rates of the disease. However, less developed countries have lower survival rates due to late diagnosis, as well as lack of adequate diagnosis and treatment. Thus, despite the highest incidence rates in developed countries, almost 50% of all cases are in less developed countries, which account for about 58% of deaths (GLOBOCAN 2008).

DOI: 10.29289/259453942018V28S1090

EPIDEMIOLOGICAL ANALYSIS OF THE DISTRIBUTION OF MAMMOGRAPHY PERFORMED IN GOIÁS, MATO GROSSO, MATO GROSSO DO SUL AND FEDERAL DISTRICT FROM 2010 TO 2013

Paula de O. C. Queiroz¹, Giselle M. Guimarães¹, Murilo H. C. e Silva¹, Isadora R. Silva¹, Túlio P. A. Basílio¹, Laine R. Martins¹

¹Universidade Federal de Goiás – Goiânia (GO), Brazil.

Objective: To evaluate the spatial distribution of mammograms performed in the central-west region, comparing the results obtained from the State of Goiás to the other States of the region (Mato Grosso, Mato Grosso do Sul and Federal District). **Methodology:** This is a descriptive study, quantitative analysis, based on data collected from any mammography type, from January 2010 to December 2013, available in the Breast Cancer Information System (SISCAM) of the Ministry of Health. **Results:** Organizing into categories, the Federal District was the state that most increased the number of mammograms in absolute and relative numbers compared to the states of the center-west region. This data is very relevant, since it indicates a greater coverage of the health system and investment when compared with the others. In addition, a very important indicator is the capital/state ratio since it shows whether the coverage of mammograms is concentrated only in the metropolitan region or if it can give access to a large part of the population of the state in a uniform way. Mato Grosso is the state with the largest distribution of mammography coverage, compared to Goiás and Mato Grosso do Sul. Only 33.54% of mammograms were performed in the capital of Mato Grosso, while in Goiás (56.53%) and in Mato Grosso do Sul (51.79%) in their respective capitals. It should be noted that the study period was from 2010 to 2013, during which data were collected. **Conclusion:** The results show significant growth of mammograms performed in the center-west region. This shows that coverage and access to this exam is being expanded. However, it also shows that the central regions performs larger numbers of mammograms in relation to the other regions of the states. This corroborates to an unequal coverage estimate for the population of the states. In addition to making access difficult due to patient locomotion problems to go to the capitals / central regions to perform the exams, this concentration of exams in the metropolitan regions can cause a crowding in the services, making access to the breast cancer screening even more difficult.

DOI: 10.29289/259453942018V28S1091

LIPOFILLING – DIFFERENT TECHNIQUES, SIMILAR RESULTS – EXPERIENCE OF THE ONCOPLASTIC COURSE OF SBM DE JAÚ - SP

Paulus F. M. Ramos¹, Maurício de A. Resende¹, Ailton Joioso¹, João R. A. Paloschi¹, Alice J. Pereira¹¹Hospital Amaral Carvalho – Jaú (SP), Brazil.

Objectives: To analyze the experience with the surgical technique of lipoenxertia of the Hospital Amaral Carvalho in Jaú - SP, during the first course of Oncoplastica by the Brazilian Society of Mastology. **Methodology:** Retrospective analysis of a series of cases, in which the lipofilling technique was used, carried out in the 1st Oncoplastic course of SBM of Jaú - SP in the year 2016–2017. Eight patients were identified using this technique. Their indications, techniques used and results were evaluated by pre and post operative photographic comparison. **Results:** During the course, eight cases were performed with the lipofilling technique in the 12 modules, of a total of 103 patients operated and 198 procedures performed. The mean age of patients submitted to lipofilling was 48.8 years (range 31–66 years). The median volume of injected fat was 232.5 mL (range 40–500 mL). The surgeries were indicated with three specific purposes: one (01) case for preparation of the recipient bed of the reconstruction that had been irradiated and the skin was not of good quality; two (02) cases of breast defect repair after conservative surgeries in irradiated breasts; and five (05) cases to improve breast contours in reconstructed breasts (4 cases of autologous tissue and 1 case of reconstruction with prosthesis). Two techniques were used, Coleman and Coleman modified, the latter with acquisition of larger volumes of fat. The pre and post operative photographic studies did not show important aesthetic differences with the two techniques. **Conclusions:** It is an important refinement technique, technically and oncologically safe, which improves the initial surgeries. Very common in oncoplastic is the need for a second intervention to correct defects, whether large or small, and this technique should be included in the arsenal of breast surgeons. When we think of patients previously submitted to radiotherapy and need new surgical intervention, this procedure becomes almost mandatory.

DOI: 10.29289/259453942018V28S1092

TATTOO – FORMAL INDICATION OF RADIOISOTOPE USE TO LOCATE SENTINEL LYMPH NODE? REPORT OF TWO CLINICAL CASES

Paulus F. M. Ramos¹, Claudia A. A. Góes¹, Tatiana P. B. B. Figueiredo¹, Rafael A. M. Hassan¹, Naiana S. C. Andrade¹, Rodrigo S. C. Guindalini¹

¹Clínica CAM/CLION – Salvador (BA), Brazil.

Introduction: Tattoos have gained popularity in recent years and have become socially accepted between men and women. It is estimated that 14% of the American population has at least one. The process of skin pigmentation creates an inflammatory reaction and in a few days the macrophages encompass the injected pigments. The carbon particles migrate through the lymphatics and within 3 days the pigments can already be seen in the lymph nodes. Tattoo ink and inflammation can block lymphatic chains, influencing detection and sensitivity, causing false-negatives in sentinel lymph node rates. **Case-report:** Two women aged 46 years, with breast cancer. The first one with left breast breast lesion and submitted to quadrantectomy with sentinel lymph node biopsy. The second had bilateral breast cancer and underwent skin-spared adenomastectomy. Both had a tattoo on the back ipsilateral to the tumor and underwent sentinel lymph node biopsy with the radiopharmaceutical technique alone. The lymph nodes were detected with the radioisotope and a parasitizing lymph node of blackened aspect that did not capture. The pathological anatomical study confirmed the presence of the tattoo pigment in the lymph nodes stained. **Discussion:** Axillary staging continues to be an important prognostic factor in breast cancer in the last decade, lymphadenectomy has been replaced by the search for sentinel lymph nodes in clinically no axilla without compromising. The combined technique using radioisotope and patent blue results in higher rates of sentinel node identification. The popularization of the skin tattoo may bring a new paradigm for the surgical staging of the armpit. The pigments can drain to the locoregional lymph nodes causing inflammation and consequent obstruction of the lymphatic channels, compromising their biopsy. The pigment may further hide the presence of metastatic cells or promote detection of the false sentinel if used only with the use of patent blue. Due to the pigmentation of regional lymph nodes by tattoo ink, there may be a failure to identify the true sentinel, and a “pseudosentinella” is identified. When they are located superficially in the armpit they can mimic the lymph node dyed blue, causing the true sentinel to be lost and the patient to undergo staging. **Conclusion:** Breast surgeons should be aware of the presence of tattoos in patients who will do sentinel lymph node screening, especially those near the drainage area to be screened. This may lead to the choice of technique and may be an indication of the use of the radioisotope as a method of choice or at least a combined technique. The use of only the blue patent may confuse the surgeon and remove only the lymph node stained by the tattoo, generating a false negative in the assessment of the axilla and impairing oncologic staging.

DOI: 10.29289/259453942018V28S1093

USE OF ASPIRATIONAL PUNCTUATION OF TUMORS WITH FINE NEEDLE (FNAB) FOR THE DIAGNOSIS OF DUCTAL CARCINOMA: ANALYSIS OF CYTOPATHOLOGICAL EXAMINATIONS DONE IN BRAZIL BETWEEN JANUARY 2012 TO JANUARY 2014

Rafael S. de Lima¹, Cassiano de S. Coutinho¹, Fabiana C. Q. S. Anjos¹, Fernando C. da Silva¹, Ildelbrando I. A. C. Sobrinho¹, Priscila F. Barbosa¹, Rayssa de O. Glória¹, Vantuir J. D. da Mota¹

¹Centro Universitário de Gurupi – Gurupi (TO), Brazil.

Objective: To analyze the number of exams performed through FNAB, with emphasis on positive results for malignancy, compatible with ductal carcinoma. **Method:** Cross - sectional study based on data from 6,433 fine - needle aspiration of breast nodules performed between January 2012 and January 2014, registered in Brazil according to the data from SisMama- DataSus. The age range, the states of the country with the highest prevalence and the cytological analysis were evaluated. The nodules were classified as benign, of indeterminate malignancy, suspected for malignancy or malignant. The chi-square test was performed with 5% significance. **Results:** Were performed 6433 FNA tests in women aged 50-64 years; being 5560 benign cases (86.43%); 111 undetermined malignancy (1.73%); 347 suspect for malignancy (5.39%); 415 positive for malignancy (6.45%). Of these, 319 exams (76.87%) were identified in FNAB as Ductal Carcinoma, being 111 between 50 and 54 years, 108 between 55 and 59 years, and 100 between 60 and 64 years. Statistically, there was an association between the FNA score and the age group, where 10.42% of the women between 60 and 64 years were diagnosed with malignant nodules ($p < 0.05$); in the age group between 55 and 59 years had 132 positive results for malignancy of 1911 exams performed (6.91%); and in the age group of 50 to 54 years had 143 positive results for malignancy of 3178 tests performed (4.50%). The states with the highest prevalence of Ductal Carcinoma cases are the states of São Paulo and Minas Gerais, with 75 and 61 cases (23.51% and 19.12%), respectively. **Conclusion:** It was observed that the majority of the cytopathological examinations carried out resulted in benign processes, but a significant number of malignant processes still exist, and of these a predominance of ductal carcinoma was observed, in which the women between 60-64 years present a higher prevalence. Corroborating with the literature, data showed that the prevalence of positive results for malignancy increased with age. According to the Ministry of Health, this neoplasm is observed in about 80–90% of the cases, and in the present study a rate of approximately 76.87% was observed.

DOI: 10.29289/259453942018V28S1094

ANALYSIS OF THE MAIN RADIOLOGICAL FINDINGS OF MAMMOGRAPHIES DONE IN THE BRAZILIAN UNIFIED HEALTH SYSTEM IN THE DISTRITO FEDERAL BETWEEN 2013 AND 2015

Brenda B. M. Mendes¹, Higor M. M. Melo¹, Lindainez A. de Souza¹, Matheus F. M. Maia¹, Wglaison P. A. Sobral¹, Fabiana C. Q. S. Anjos¹, Rafael S. Lima¹, Priscila F. Barbosa¹

¹Centro Universitário de Gurupi – Gurupi (TO), Brazil.

Objective: To analyze the main radiological findings of mammograms performed in the Unified Health System in the Distrito Federal, correlating them with the age group. **Methods:** Cross-sectional study, in which the statistical analysis of variance was analyzed from 96,857 mammographic examinations performed in the Distrito Federal female patients of different age, race/color, from January 2013 to July 2015, being the data obtained from the Information System of Breast Cancer (SisMama). The chi-square test was performed with 5% significance. **Results:** In the analyzed period, 96,857 mammograms were performed in women in the Distrito Federal, a predominant age group age 40–59 years (71.96%). No radiological findings were found in 81.54% of the breasts. When comparing the laterality of the breast regarding the radiological findings, it cannot be compared with the significant statistic ($p=0.75$). The nodules that appear discretely predominate in the left breast, where as the microcalcifications were more found in the right breast. A total of 16,981 nodules, mostly between 11–20 mm and 1,352 microcalcifications were counted. Of the axillary lymph nodes found, 91.33% were normal. Regarding BI-RADS, it was observed that 15.61% of mammographies were inconclusive (BI-RADS 0). Benign findings were found in 70% ($n = 67,834$) of mammograms (BI-RADS 2). A total of 2,621 women (2.70%) presented BI-RADS 4, 5 or 6. Analyzing the age group, 57.21% of the women diagnosed with BI-RADS 5 had between 50–69 years ($p<0,05$), and 18.71% were between 40–49 years old. **Conclusion:** The close relationship between the radiological findings of the breasts and the age group of the patients leads us to recognize the importance of this examination for the early diagnosis and effective treatment. It was observed that in the great majority of mammograms performed no radiological findings were found. However, it is worth mentioning that the rate of women between 40–49 years diagnosed with BI-RADS 5 is considerable and worrying. This fact evidences the importance of the early screening of the malignant mammary disease through annual mammography from the age of 40, as recommended by the Brazilian Society of Mastology.

DOI: 10.29289/259453942018V28S1095

ANALYSIS OF THEIR DELETIONS AND DUPLICATIONS IN THE *BRCA1* AND *BRCA2* GENES IN PATIENTS FROM GOIÁS-BRAZIL WITH SUSPECTED OF HEREDITARY BREAST AND OVARIAN CANCER SYNDROME

Rebeca M. Goveia¹, Paula F. F. Silva¹, Thais B. Teixeira², Bruno F. Gamba¹, Aliny P. Lima¹, Ruffo Freitas-Junior², Elisângela P. Silveira-Lacerda¹

¹Laboratório de Genética Molecular e Citogenética, Instituto de Ciências Biológicas, Universidade Federal de Goiás (UFG) – Goiânia (GO), Brazil.

²Departamento de Mastologia, Hospital das Clínicas, UFG – Goiânia (GO), Brazil.

Breast cancer is the second most common cancer in the world, and the most common among women population, about 5% to 10% of cases are hereditary and half of them are caused by hereditary breast and ovarian cancer (HBOC), caused by variations in *BRCA1* and *BRCA2* genes. The present study aimed to identify deletions and duplications prevalence in *BRCA1* and *BRCA2* genes on breast cancer patients in Goiás, Brazil, and was approved by the National Research Ethics Committee according with number of CAAE 50626315.6.0000.5078. Were evaluated 46 breast cancer's patients who fulfilled National Comprehensive Cancer Network (NCCN) criteria for HBOC syndrome screening. About 4 mL blood sample was collected from each patient for DNA extraction using commercial kit and the Multiplex Ligation Dependent Probe Amplification (MLPA) technique was performed using the SALSA MLPA P002 *BRCA1* and SALSA MLPA P045 *BRCA2/CHECK2* kits. The majority of the patients were female (97.83%) and the mean age of the patients was 37.52 years. In this group, 43.47% of the patients were younger than 35 years at the time of diagnosis and 35% of them were diagnosed with triple negative tumors. The most common molecular subtype was luminal A (46.2%) followed by triple negative tumors (28.20%). Four family history profiles were observed, which 45.65% had a family history of breast cancer and other cancers, 33% had a family history of cancer without cases of breast cancer, 17.39% had only a personal history of cancer in the family and 4.34% of the patients had only a family history of breast cancer. No patient was found with rearrangements in the *BRCA1* gene. In the *BRCA2* gene, one patient (2.12%) presented a false positive result for the heterozygous deletion of the 27 exon, which may have been caused by the presence of a small change in the probe binding region. This was the first study performed to analyze large deletions and duplications in patients from the central-western region of Brazil. Can be conclude that the frequency of large deletions and duplications in the *BRCA1* and *BRCA2* genes is low in the population of Goiás. **Acknowledgments:** LGMC, CORA and CAPES.

DOI: 10.29289/259453942018V28S1096

A FUNCTIONAL *IN VIVO* SHRNA SCREEN FOR THE IDENTIFICATION OF TRIPLE-NEGATIVE BREAST CANCER METASTASIS SUPPRESSOR GENES

Rebeka Tomasin¹, Ana Carolina Baptista Moreno Martin¹, Márcia Regina Cominetti¹, Kaylene Simpson², Robin Anderson³, Richard Redvers³

¹Laboratory of Biology of Aging, Department of Gerontology, Universidade Federal de São Carlos – São Carlos (SP), Brazil.

²Peter MacCallum Cancer Centre, Victorian Comprehensive Cancer Centre – Australia.

³Olivia Newton John Cancer Research Institute – Australia.

Objective: When diagnosed early, breast cancer can be treated with a great rate of success. However, once the disease reaches metastatic stage, spreading to organs such as lungs and bones, the available treatments are extremely limited and prognosis is poor. This is particularly true for the triple-negative subtype. By identifying genes that regulate metastasis, it would be possible to predict the patients whose disease is more likely to spread, and it would create an opportunity to develop more effective therapies. **Methods:** Using a genome-wide lentiviral shRNA library, we initiated an *in vivo* functional screen aiming to identify genes whose reduction in expression levels, in non/poorly metastatic murine triple-negative breast cancer cells (67NRch1/66cl4ch14, respectively), would lead to spontaneous metastasis *in vivo*. **Results:** Dozens of putative metastasis suppressors were identified, in a wide range of organs, by conventional and next generation sequencing. These candidates were further subjected to an *in vitro* RNAi screen in 66cl4ch14 cells, involving viability, wound healing and adhesion assays. **Conclusion:** Integrative analysis of both screens (*in vivo* and *in vitro*), literature review, RNA-seq and bioinformatics analysis of gene expression and clinical outcome (OncoPrint/Breastmark) indicated highly-confidence candidates to be taken for further analysis and validation *in vivo*. Additional *in vivo* and *in vitro* experiments, aiming to validate the obtained candidates as *bona fide* metastasis suppressor genes, as well as elucidate their possible mechanisms of action, are ongoing.

DOI: 10.29289/259453942018V28S1097

AESTHETIC OUTCOMES AND PATIENTS SATISFACTION AFTER DELAY/IMMEDIATE AUTOLOGOUS BREAST RECONSTRUCTION

Renata S. Brondi¹, Vilmar M. Oliveira¹, Fabio Bagnoli¹, José Francisco Rinaldi¹, Evandro Mateus Falacci¹, Fabrício P. Brenelli¹

¹Faculdade de Ciências Médicas, Santa Casa de São Paulo, Universidade Estadual de Campinas – Campinas (SP), Brazil.

Aim: The timing and aesthetic outcome of the breast reconstruction may be affected by the need for radiation therapy (RT). Autologous reconstruction is a novel surgical approach for patients in whom the need for adjuvant RT after mastectomy is preoperatively unknown. The aim of this study was to evaluate the outcomes and satisfaction of breast reconstruction using a fat-grafted latissimus dorsi flap in immediate and late reconstruction. **Method:** A total of 18 patients were submitted to autologous latissimus dorsi flap, with 11 immediate and 7 late reconstructions. Patient demographics, surgical characteristics, and complications were recorded. During outpatient follow-up, the reconstruction was evaluated and compared to the contralateral breast in relation to shape, volume and symmetry. Patients scored their satisfaction using a like scale between 12–18 months after surgery, and two surgeons performed their evaluations in that same time interval. **Results:** Of the eleven patients submitted to immediate reconstruction, ten performed neoadjuvant chemotherapy, and 9 performed adjuvant radiotherapy. All patients who underwent late reconstruction had undergone radiotherapy prior to reconstruction. All the flaps received autologous fat only in a surgical time prior to the confection of the flap. All flaps had 100% viability. In the subjective evaluation through the like scale of the shape, volume and breast symmetry, it was evidenced that the patients were more satisfied than the physicians in all parameters analyzed, with good and excellent evaluation of greater than 80% in relation to the shape and symmetry, this evaluation proved to be in agreement with the medical evaluation, and by the physicians the evaluation was better in relation to the symmetry, 64% classified between good and excellent. In the separate assessment of patient groups, we can note that even after adjuvant radiotherapy patients who underwent immediate reconstruction were more satisfied than patients submitted to late reconstruction, concentrating the evaluation in excellent in 77,8% in relation to the form, 55,6% volume and 44,4% in symmetry. Radiotherapy does not seem to cause great damage to the breast reconstructed by this technique. We conclude that the autologous latissimus dorsi flap is an accessible procedure, with long-term and lowcost results to the health system, and can be performed in small centers and with a high patient satisfaction rate

DOI: 10.29289/259453942018V28S1098

THERAPY OF DIGNITY: STRATEGY FOR THE CONSTRUCTION OF A LEGACY AT THE END OF LIFE

Natália Novaes Pavani Araújo¹, Marcella Tardeli Esteves Angelloni Santana¹, Ricardo Caponero¹, Joice Aline Klein¹, Ana Lúcia Coradazzi¹, Danilo Augusto Mendonça Faleiros¹

¹Universidade de São Paulo – São Paulo (SP), Brazil.

Introduction: Dignity Therapy is a tool proposed by Chochinov et al. (2005) that aims to build a legacy through a letter written with the patient in palliative care, which can be delivered to loved ones in life or after his death. The multidisciplinary team can use this technique in their practice, not only with the purpose of promoting patient care while preserving their dignity, but also offering a holistic attention to the patient / family care unit. **Objective:** to report a patient experience based on Dignity Therapy intervention. **Method:** This is a descriptive, case study. The activity was developed in a large private hospital, in São Paulo, in February 2018. The sample consisted of a 53-year-old patient with advanced malignant neoplasm at the end of life. The intervention was conducted in two meetings with the formulation of 17 letters dictated and signed by the patient and written by the chaplain of the service in the period that preceded her palliative sedation. All the letters were destined to the daughter in order to open them over the next few years. **Result:** After the preparation of the letters, the patient's psychosocial suffering was improved, which consisted of no longer participating in the main moments of the daughter's life due to her impending death, as well as the possibility of retrieving memories and projecting expectations, manifesting hope and become, symbolically, a participant in the daughter's future. **Considerations:** It is considered that the team's performance in an interdisciplinary approach, and based on the therapy of dignity, contributed with the care of the intense psychosocial suffering experienced by the patient in the terminal phase of the disease.

DOI: 10.29289/259453942018V28S1099

PROGRAM OF QUALITY CONTROL IN MAMMOGRAPHY IN THE STATE OF GOIÁS: EXPERIENCE OF 11 YEARS

Rosangela da Silveira Corrêa¹, Ruffo Freitas-Junior¹, Suzana Alves Bastos², Lucy Aparecida Parreira Marins³, Rodrigo Massakatsu Nishiharu Tanaka³, Dagoberto Luiz Suzana Costa¹

¹Centro Avançado de Diagnóstico da Mama, Universidade Federal de Goiás, Rede Brasileira de Pesquisa em Mastologia – Goiânia (GO), Brazil.

²Superintendência de Vigilância em Saúde do Estado de Goiás – Goiânia (GO), Brazil.

³Departamento de Vigilância Sanitária de Goiânia – Goiânia (GO), Brazil.

Objective: To evaluate the results of the last ten years of a Mammography Quality Control Program in Goiás (PCQMGO).

Methods: The unit of observation was the Diagnostic Centers (CD), which performed mammography in the state of Goiás, from 2007 to 2017. The structure data were collected and tests were performed to evaluate the image quality, the equipment performance parameters and the radiation dose applied in the examinations. **Results:** In 2007, PCQM-GO 103 equipment was registered, 24% of which were CR-type digital and 76% were conventional. In 2017, 197 CDs were registered, of which 160 were in operation, with 164 equipment, 87% of which were digital CR type and 3% DR type, and 10% were conventional. It was observed that, during the period evaluated, 37 CD discontinued mammography. Initially, the technical parameters that presented improvement due to the quality control action were related to the adjustments of the breast compression force, the Automatic Exposure Control device and the alignment of the compression tray. In the last evaluation, 87.3% of the CDs showed compliance in image quality, 71.8% in the Simulator Input Dose and 72.2% in the tension accuracy. **Conclusion:** Experience in the execution of the Program showed that it is possible to maintain greater quality and risk control in mammography.

DOI: 10.29289/259453942018V28S1100

“O TOQUE PELA VIDA”: THE IMPORTANCE OF THE PINK OCTOBER CAMPAIGNS FOR WOMEN HEALTH IN JATAÍ, GO

Sabrina R. J. Pimenta^{1,2}, Thaynara B. C. Souza^{1,2}, Izabella C. Amaral^{1,2}, Thayanne S. Ramos^{1,2}, Carlos A. L. Silva², Carolina A. E. Terceiro², Bianca D. Vieira², Ana Paula S. Perez²

¹Department of Extension and Culture, Universidade Federal de Goiás – Cidade (GO), Brazil.

²Universidade Federal de Jataí – Jataí (GO), Brazil.

Objectives: The extension project “O Toque pela Vida” was created in 2016 by teachers, administrative technicians and Medicine students from the Federal University of Jataí (UFJ), in partnership with the Jataí’s Municipal Secretariat of Health (MSH). It aims to inform and guide the population regarding etiology, risk factors and early detection of Breast Cancer (BC), in addition to providing preventive examinations to women. **Methodology:** For the project execution, study groups were realized out monthly for the discussion of articles and clinical cases about BC. The information obtained during the groups discussions was used to design pamphlets, banners, posters, T-shirts and ribbons, symbols of the October Pink campaign. Extension activities and educational materials were disseminated on social networks, local radio and television programs. The activities were developed in the main Basic Health Units (BHUs) and in the Raízen Company of Jataí. Among these activities, on October 18, 2017, project students lecture on BC in the Raízen Company. On October 6, 2017, project participants went to BHUs to distribute information materials about BC and to disclose the D-Day Campaign. This campaign took place at BHU James Phillipe Minelli on October 28, and in the external area of the Unit, passwords were given to women for they could receive medical attention and care. Meanwhile, the project members measured the patients’ pressure and glycemia. In addition, patients received guidance and clarification about BC, regarding risk factors, symptoms and early detection of cancer. The orientation activities were performed using anatomical breast molds, posters and ribbons distribution. Inside the unit, the patients updated the Unified Health System (UHS) card and filled in the necessary forms for medical care. The medical service were conducted by physicians, along with medical students. During the examinations, population screening for BC was verified, and women over 50 years old and/or presenting changes in clinical examination were scheduled for mammography. **Results:** During the activities held in the company, around 100 women employees, attended the educational lecture and, at the end, asked questions about BC. The project “O Toque pela Vida” was more widely accepted by the population, compared to the year 2016, which was the beginning of the project. In this year, the actions were focused only on the BC guidelines. However, in 2017, the partnership with MHS and the wide dissemination in the BHUs and media, allowed the project growth. Thereby, the D-Day brought around 200 women to the BHU James Phillipés Minelli. Of the 200 women, 152 were scheduled for mammograms performed by professionals from the Jataí’s MHS, 09 were unilateral for diagnosis and 143 were bilateral for tracking the BC. Mammographies were scheduled between October 31 and December 1, 2017. **Conclusion:** The partnership established between the project “O Toque pela Vida” and the MHS shorted mammography waiting process in the Unified Health System (UHS). Thereby, the scheduling was quick and accessible to a great number of women, during the D-Day in October Pink Campaign. In addition to the activities developed during October, the study groups qualified the Medicine students and they were able to transmit the information to women population in a clear and correct way, taking into account that the project is composed by fresh and senior students. Therefore, the project proved itself of great relevance for the population, by providing more information and effecting the prevention and early diagnosis of breast cancer.

DOI: 10.29289/259453942018V28S1101

RETROSPECTIVE COHORT STUDY WITH SURVEY ANALYSIS OF BREAST CANCER SUBMITTED RADIOSURGERY BY CEREBRAL METASTASIS IN 10 YEARS

Nilceana Maya Aires Freitas¹, Thais Franco Simionatto¹, Nathalya Ala Yagi¹, Jean Teixeira de Paiva¹, Thais de Toledo Lima Santana¹, Sebastião Berquó Peleja¹, Mattheus Humberto do Vale¹, Silvia Sousa Pires¹

¹Hospital Araújo Jorge – Goiânia (GO), Brazil.

Objectives: Breast cancer is the second most common type of cancer associated with brain metastasis. Symptomatic brain metastasis are diagnosed up to 16% of patients with metastatic breast cancer. The risk of brain metastasis in ten years depends on the phenotypic subtypes of breast cancer. Luminal A with a risk of 0.7%; Luminal B, 12%; Luminal HER2 Positive, 8%; HER2, 12%; and Triple Negative, 7%. Stereotactic Radiosurgery (SRS) is one therapeutic modalities used to treat of brain metastases, which offers local control and survival comparable to neurosurgery. This study aims to evaluate the epidemiological profile of patients submitted to radiosurgery, correlate survival, and time tumor progression in the brain. **Methods:** Data were collected from patients with diagnosis of breast cancer brain metastasis treated with SRS from 2007 to 2017 at Hospital Araújo Jorge/Goiânia/Brazil. It was analyzed immunohistochemistry pattern, number of lesions treated with SRS, neurosurgery, whole brain radiotherapy (WBRT). Statistical analysis was performed using Fisher's test and Kaplan-Meier curve, considering 5 criteria separately: overall survival (OS), progression-free survival (PFS) for the central nervous system (CNS) after radiosurgery, brain disease free survival (DFS) after diagnose and survival after radiosurgery for brain metastasis (SRC). **Results:** In a time 10 years, total 78 patients with brain metastasis underwent to SRS, 71 (91%) were treated with a single dose procedure and six (8%) with a five fractions procedure. Among these, 59 (76%) were submitted to SRS in a single, 15 (19%) in a double, and four (5%) in three procedure. The number of lesions treated ranged from one to six metastases. Most patients were triple negative (30%) and HER2 positive (30%). Twenty patients (26%) underwent to WBRT, 60% after SRS. 34 patients (44%) presented brain disease progression after the first SRS, in the mean time of 9 months and 17 patients (22%) died after 19 months, and OS 78% The mean of follow-up was 12 months, with the maximum of 60 months. **Conclusion:** SRS is a therapeutic tool that improves survival, providing better quality of life and lower neurocognitive deficit when compared with WBRT in the treatment of brain metastases. WBRT can be avoided up to 3/4 of the population that was treated by SRS as a primary treatment for breast cancer brains metastasis and more than a half presented with stable disease control at CNS after radiosurgery.

DOI: 10.29289/259453942018V28S1102

THE INFLUENCE OF THE BREAST TUMOR BED CLIPPED IN THE HEART AND LUNG IRRADIATED VOLUME DURING THE BOOST RADIATION PLANNING

Thaís Franco Simionatto¹, Nathalya Ala Yagi¹, Ana Flávia de Paula Guerra Campedelli¹, Nilceana Maya Aires Freitas¹, Mattheus Humberto do Vale¹, Silvia Sousa Pires¹, Jean Teixeira de Paiva¹, Carolina Martinelli Bezerril¹

¹Hospital Araújo Jorge – Goiânia (GO), Brazil.

Objetives: Adjuvant radiotherapy is a standard procedure particularly in women following breast conserving surgery (BCS). Markers may be employed for accurate localization of the boost volume during treatment planning. The implanted clips are typically stable in the tumor bed throughout the course of radiation therapy. Brazil is one of the most countries that the oncoplastic surgery is employed in breast conservative treatment, and so the importance to clip tumor bed to guide the radiation boost planning volume. The purpose of this study was to evaluate the heart and lung irradiated volume considering the presence of surgical clips in breast tumor bed during the radiation treatment planning. **Methods:** A retrospective descriptive study of 103 post-surgical patients who underwent breast-conserving treatment in early breast cancer. Statistical analysis was performed using Fisher's test, with a 95% confidence interval (CI, $p < 0.05$). It was considered the irradiated volume of lung and heart relative to 40% of prescribed dose in the boost radiation planning (V40) that was compared in patients with or without surgical clips. Using the dose-volume histogram (DVH). **Results:** This study evaluated 103 from from 2011 to 2018, and the average age was 57 years, varying from 32 to 78 years; the predominant histological type was invasive ductal carcinoma, 90%; 76% of the patients were luminal type, which 53% were Luminal A and 23% were Luminal B. Among the patients, 62% had left breast carcinoma and 38% right breast tumor. The mean heart V40 of the right breast was 2.0 cm³ (with clip) and 0 cm³ (without clip); and the left breast was 3.4 cm³ (with clip) and 0.9 cm³ (without clip). The mean lung V40 of the right breast was 60,86 cm³ (with clip) and 54.81 cm³ (without clip); and the left breast was 49.43 cm³ (with clip) and 43.71 cm³ (without clip). **Conclusion:** It is known that the use of the clip increases the local control. It was observed that the pulmonary and heart volume irradiated in the boost planning was higher in the group of clipped tumor bed in the tumor bed. Despite the probability of a geographic error, due to the clip use, it can lead to a better local control. In contrast, a increase of irradiated volume in normal tissue, like the heart and lung, can increase the possibility of side effects.

DOI: 10.29289/259453942018V28S1103

PREVALENCE ANALYSIS OF DUCTAL CARCINOMA *IN SITU* OF BREAST IN WOMEN POPULATION OF GOIÁS BETWEEN 2009-2014

Lucas N. Silva¹, Richard R. Santos¹, Julio M. B. Garcia¹, Thaynara de M. Pacheco¹¹Universidade Federal de Goiás – Goiânia (GO), Brazil.

Objective: Compare the prevalence of ductal carcinoma *in situ* of the breast and the prevalence between age groups in the period from 2009 to 2014 in the state of Goiás. **Methodology:** This is a descriptive study, using the Information System for Breast Cancer Control (SISMAMA). The analyzed data correspond to the reported cases of ductal carcinoma *in situ* in the female population in the years 2009 to 2014 in Goiás. **Results:** We identified 59 cases of ductal carcinoma *in situ* (DCIS) and lobular carcinoma *in situ* from September 2009 to February 2014 in the female population in Goiás. In 2009, only 2 cases of DCIS were reported, with intermediate histological grade, affecting the population between 20 and 39 years. In 2010, 16 cases were identified, 56% of intermediate grade DCIS. The age group most affected this year was between 50 and 59 years. The population most affected in 2011 was over 60 years. Furthermore, 11 cases were identified, 55% of intermediate grade DCIS. The most affected population in 2012 was between 40 and 49 years. There were a total of 10 cases this year, 50% of them high-grade DCIS. Eighteen cases were identified in 2013, and 67% of them were of intermediate grade DCIS. Two groups were equally affected: between 40 and 49 years and over 60 years. In 2014, 2 cases were identified, one of intermediate and one high-grade DCIS, at the extremes of the age groups. **Conclusion:** It is conceived that, in Goiás, DCIS is the main cause of breast cancer in the female population. About the age group, there is an alternation, with predominance over 40 years. The year with the highest number of cancer was 2013.

DOI: 10.29289/259453942018V28S1104

ANALYSIS OF THE INCIDENCE OF PRIMARY BREAST CANCER IN RELATION TO THE TOTAL CASES IN THE FEMALE POPULATION OF THE CENTRAL-WEST SCENARIO

Paula O. C. Queiroz¹, Nayara P. A. Oliveira¹, Hygor A. P. Dutra¹, Thaynara P.¹

¹Universidade Federal de Goiás – Goiânia (GO), Brazil.

Objective: Compare the incidence of primary breast cancer in relation to the total cases in the female population of the Central-West scenario and capitals, except Federal District. **Methodology:** Cancer Incidence Estimates for the biennium 2018-2019 produced by the Division of Surveillance and Situation Analysis of the Coordination of Prevention and Surveillance (Conprev) of INCA / MS were used. **Results:** According to INCA data, breast cancer will account for 19.24% of all cancer cases expected in Brazil in 2018 and 2019. For the Center-West region, this percentage is estimated to be 29%. The state in which breast cancer has a higher incidence compared to the other neoplasms is Mato Grosso do Sul (60.90%), in which the capital (Campo Grande) reaches 81.05%. Goiás presents the second highest percentage (48.68%), being in the capital Goiânia 72.17%. Finally, in Mato Grosso, the lowest value is expected (41.32%) and in its capital (Cuiabá), 59.03%. **Conclusion:** Regional disparities in access to early detection of breast cancer could explain a higher or lower incidence of breast cancer in the different states that make up the Central-West. It is known that, with the increase of the access by the population to mammography, the incidence of breast cancer tends to increase, reflecting greater access to early diagnosis. In this sense, a lower mortality rate can be expected in these regions, due to an early diagnosis.

DOI: 10.29289/259453942018V28S1105

QUANTITATIVE EVOLUTION OF MAMMOGRAPHY IN COUNTIES OF GOIÁS BETWEEN 2010 AND 2013

Thaynara Mercadante¹, Guilherme A. Sampaio¹, Maria L. de Oliveira¹, Natália S. D. Mendonça¹, Caio Matheus Fonseca de Brito¹, Jonatha Fonseca Lopes¹, Letícia de Souza Cainelli¹, Pedro Rodrigues Soares¹

¹Universidade Federal de Goiás – Goiânia (GO), Brazil.

Objectives: Analysing the quantitative evolution of mammograms in counties in the state of Goiás between the years 2010 to 2013. **Methodology:** this is a descriptive study using data from the Information System of breast cancer (SISMAMA). The reports reviewed were mammograms of women between 40 to 69 years in Goiás, during the years 2010 to 2013. **Results:** in the year 2010, the total number of tests was of 47,542. The distribution was: Goiânia with 28,903, Anápolis with 8,194 and Catalão with 1052. Right behind, the counties with the highest number of mammograms was Rio Verde, Goiatuba, Porangatu and Jataí. In 2011, the total amount of mammography per year increased. However, there was drop in the number of tests in the city of Goiânia and increased in other cities - mainly in Catalão and Rio Verde. In 2012, the highlight was for the county of Goiânia which increased considerably (more than 4000 examinations). In the year 2013 there was drop in the total number of mammograms performed comparing to the previous year (51,797 mammograms in 2013 and of 54,312 in 2012). In general, it was observed an increase of 4,255 mammograms carried out during the period from 2010 to 2013. Specifically, among the cities that were included in the 10 with the highest number of tests carried out in 2010, 6 remained during the period (Goiânia, Anápolis, Catalão, Rio Verde, Jataí and Aparecida de Goiânia) and four varied, with the municipalities of Goiânia, Anápolis and Catalão remained in firsts positions respectively. **Conclusion:** the cities with more features like Goiânia, Anápolis and Catalão have remained in top positions, showing the concentration of mammographers in the most developed cities and the difficulty of access to examination by women from smaller towns. The Brazilian Society of Mastology recommends that mammography is performed from women over 40 years annually for women with no risk factors. Among women who have access to preventive mammography, the number of deaths caused by breast cancer decreases considerably due to early diagnosis. Therefore, it is important to encourage practices such as “pink october” awareness campaigns so that it is easier to carry out the examination and the decentralization of large centers in order to reach a larger number of women occurs.

DOI: 10.29289/259453942018V28S1106

THE MASTOLOGIST PROFILE IN BRAZIL

Thaynara Mercadante¹, Guilherme A. Sampaio¹, Maria L. de Oliveira¹, Natália S. D. Mendonça¹, Caio Matheus Fonseca de Brito¹, Jonatha Fonseca Lopes¹, Letícia de Souza Cainelli¹, Pedro Rodrigues Soares¹

¹Universidade Federal de Goiás – Goiânia (GO), Brazil.

To analyze in a comparative way the number of mastologist doctors in Brazil who attended in the year of 2015. Data were analyzed from the study “Medical Demography in Brazil 2018” performed by the Federal Council of Medicine (CFM) and the Regional Council of Medicine of São Paulo (CREMESP), comparing them with the same study carried out in previous years (2011, 2013, 2015). In 2017, in Brazil, 0.6% of medical specialists were mastologists, totaling 2,219 professionals corresponding to 1.07 specialists per hundred thousand inhabitants. The mean age of the mastologists was 45.5 years and 50.9% of them are men. Among the mastologists, according to the Brazilian macro regions, 52.7% work in the Southeast, 20.9% in the Northeast, 13.8% in the South, 8.5% in the Midwest and 4.1% in the North. In addition to mastologists, 1,710 are obstetrician gynecologists, 320 are general surgeons, 90 are clinical oncologists and 78 oncology surgeons. From the data presented, we can observe the expansion of the specialty in Brazil. In 2011, the number of mastologists in the country was 669, corresponding to 0.33% of medical specialists at the time. In six years, this percentage gradually increased to 0.6% (2,219 professionals), from 0.75 mastologists per 100 thousand inhabitants in 2013 to 1.07 in 2017. The mean age during the analyzed periods was oscillating 44, 3 years in 2011, 45.41 in 2013, 44.8 in 2015 while that of the percentage by sex that went from 60.4% of males in 2013 to 50.9% in 2017, a trend that accompanies the feminization of medicine in Brazil. It is also noticeable the heterogeneous distribution of this specialty in the Brazilian regions, with more than half of the mastologists (52.7%) concentrating only in the southeast region. The main access route for mastology continues to be obstetrical gynecology (choice of 77% of mastologists working in 2017). In addition, a significant number of mastologists are also specialists in clinical or surgical oncology, both currently (168 professionals) and in the last two studies (117 in 2013 and 146 in 2015).

DOI: 10.29289/259453942018V28S1107

NARRATIVE REVIEW ON THE CONSUMPTION OF SOY AND DERIVATIVES RELATED TO THE RISK OF BREAST CANCER

Fernando C. Ramos¹, Nayara P. Almeida¹, Thaynara Mercadante¹

¹Universidade Federal de Goiás – Goiânia (GO), Brazil.

Objective: The objective of this work is to know the studies developed about the correlation of soy food intake and its relation with the risk of breast cancer, synthesizing the most recent recommendations. **Methodology:** In the preparation of this narrative review some search terms were used: Breast cancer, soy food intake and risk. These terms were introduced in Brazilian and international bibliographic databases, such as SciELO - Scientific Library Online. In addition, articles were searched in international journals such as: The American Journal of Clinical Nutrition and British Journal of Cancer. **Results:** In the analyzed articles, what is observed is an inverse relationship between the consumption of soy and its derivatives and the risk of breast cancer. The consumption of soy and its derivatives, in a moderate amount, decreases the risk of breast cancer and, also, the percentage of recurrence of the disease, once diagnosed. The effect of isoflavones as an antiestrogen and the anticancer effects of multiple soy components such as DNA topoisomerase inhibition, immunoinflammatory modulation and angiogenesis blockers have been pointed out in most studies as responsible for protective action. Some studies in Asian women have come to estimate 15% percent reduction in the risk of developing breast cancer, every 10 mg of isoflavone ingested per day. In addition, studies also showed that isoflavones had no interaction with tamoxifen in patients taking this therapy. **Conclusion:** This review allowed us to analyze the correlation between soy consumption and its derivatives with breast cancer risk and mortality, suggesting that moderate soy food intake is safe and potentially beneficial for women with breast cancer. This review can highlight the subject for future works.

DOI: 10.29289/259453942018V28S1108

DOUBLE MUTATION CARRIER IN BRCA2 AND MUTYH GENES: A CASE REPORT IN A YOUNG BREAST CANCER PATIENT

Deidimar C.B. Abreu¹, Leandro G. Oliveira¹, Pedro P. B. Abreu¹, Vera A. Saddi¹

¹Instituto Goiano de Oncologia e Hematologia, Pontifícia Universidade Católica de Goiás, Associação de Combate ao Câncer em Goiás – Goiânia (GO), Brazil.

Objectives: To describe a case of a young female patient (32 years old) diagnosed with breast cancer and submitted to a genetic testing with an unusual result. **Case description:** The tumor was an invasive ductal carcinoma, estrogen and progesterone receptors positive and HER2 negative. The family history revealed a paternal aunt with bilateral breast cancer at about 60 years old, a paternal grand uncle with a central nervous system cancer, and another paternal granduncle with leukemia, who had three daughters with breast cancer at young age. During the treatment period, the father of the patient was also diagnosed with breast cancer. The genetic test revealed a concomitant double pathogenic variant, namely c.9382C>T (p.Arg3128*) in *BRCA2* and c.1187G>A (p.Gly396Asp) in the *MUTYH* gene. **Discussion:** A pathogenic mutation in *BRCA2* gene is associated to increased risk of developing breast and ovary cancer in young age, compared to general population. The risk of pancreatic cancer and melanoma are also increased by this mutation. A pathogenic mutation in the monoallelic *MUTYH* gene seems to slightly increase the risk of developing colorectal cancer, compared to average woman. Although the cancer risk in individuals who carry multiple pathogenic variants has not been established for combined *MUTYH* and *BRCA2* pathogenic variants, the identification of multiple pathogenic variants does allow for screening for cancers associated with both syndromes and has implications for cancer risk for family members. More information is needed on the frequency of multiple pathogenic variants, as well as the phenotypic spectrum when multiple pathogenic variants are present.