

MASTOLOGY

Official Journal of the Brazilian Society of Mastology

Volume 35, 2025

ISSN 2594-5394



MASTOLOGY

Official Journal of the Brazilian Society of Mastology

Volume 35, 2025

EDITOR-IN-CHIEF

Rafael Henrique Szymanski Machado (Rio de Janeiro; RJ; Brazil)

CO-EDITORS

Gil Facina (Sao Paulo; SP; Brazil)

René Aloisio da Costa Vieira (Barretos; SP; Brazil)

Ruffo de Freitas Júnior (Goiania; GO; Brazil)

SPECIALTY EDITORS

MASTOLOGY

André Mattar (Sao Paulo; SP; Brazil)

Alfredo Carlos Simões Dornellas de Barros (Sao Paulo; SP; Brazil)

Antonio Luiz Frasson (Porto Alegre; RS; Brazil)

Cássio Cardoso Filho (Campinas; SP; Brazil)

Cícero de Andrade Urban (Curitiba; PR; Brazil)

Daniel de Araújo Brito Buttros (Rio Claro; SP; Brazil)

Délio Marques Conde (Goiania; GO; Brazil)

Fabiana Baroni Makdissi (Sao Paulo; SP; Brazil)

Fábio Bagnoli (Sao Paulo; SP; Brazil)

Fabio Postiglione Mansani (Ponta Grossa; PR; Brazil)

Fabrizio Palermo Brenelli (Campinas; SP; Brazil)

Felipe Pereira Zerwes (Porto Alegre; RS; Brazil)

Flavia Maria de Souza Clímaco (Rio de Janeiro; RJ; Brazil)

Gustavo Antonio de Souza (Campinas; SP; Brazil)

Gustavo Zucca-Matthes (Barretos; SP; Brazil)

Joaquim Teodoro de Araújo Neto (Sao Paulo; SP; Brazil)

Jordana de Faria Bessa (Sao Paulo; SP; Brazil)

José Luiz Pedrini (Porto Alegre; RS; Brazil)

Jose Roberto Filassi (Sao Paulo; SP; Brazil)

Jurandyr Moreira de Andrade (Ribeirão Preto; SP; Brazil)

Luiz Henrique Gebrim (Sao Paulo; SP; Brazil)

Marcelo Madeira (Sao Paulo; SP; Brazil)

Maria Júlia Gregório Calas (Rio de Janeiro; RJ; Brazil)

Renato Zocchio Torresan (Campinas; SP; Brazil)

Roberto José S. Vieira (Rio de Janeiro; RJ; Brazil)

Rodrigo Gonçalves (Sao Paulo; SP; Brazil)

Rogério Fenile (Sao Paulo; SP; Brazil)

Rosemar Macedo Sousa Rahal (Goiania; GO; Brazil)

Vinícius Milani Budel (Curitiba; PR; Brazil)

INTERNATIONAL ADVISORY BOARD

Eduardo González (Buenos Aires; Argentina)

Gail Lebovic (Dallas; Texas; Estados Unidos)

Jaime Letzkus Berríos (Santiago; Chile)

Luciane Cavalli (Washington DC; DC; Estados Unidos)

Mahmoud El-Tamer (Nova Yorque; Estados Unidos)

Marcelo Cruz (Chicago; Illinois; Estados Unidos)

Rui Manoel dos Reis (Lisboa; Portugal)

GENETICS

Maria Isabel Achatz (Sao Paulo; SP; Brazil)

PATHOLOGY

Ângela Flávia Logullo Waitzberg (Sao Paulo; SP; Brazil)

Helenice Gobbi (Belo Horizonte; MG; Brazil)

PHYSIOTHERAPY

Anke Bergmann (Rio de Janeiro; RJ; Brazil)

Samantha Karla Lopes de Almeida Rizzi (Sao Paulo; SP; Brazil)

TRANSLATIONAL RESEARCH

Tatiana Carvalho de Souza Bonetti (Sao Paulo; SP; Brazil)

MEDICAL ONCOLOGY

Max Mano (Sao Paulo; SP; Brazil)

Sérgio Simon (Sao Paulo; SP; Brazil)

RADIOTHERAPY

Nilceana Maya Aires Freitas (Goiania; GO; Brazil)

Rodrigo Souza Dias (Sao Paulo; SP; Brazil)

Samir Abdallah Hanna (Sao Paulo; SP; Brazil)

RADIOLOGY

Helio Amâncio Camargo (Sao Paulo; SP; Brazil)

Simone Elias (Sao Paulo; SP; Brazil)

FORMER PRESIDENTS

Alberto Lima de Moraes Coutinho (1959–1961)
Jorge de Marsillac (1962–1963)
Eduardo Santos Machado (1964–1965)
Carlos A. M. Zanotta (1966–1967)
Alberto Lima de Moraes Coutinho (1968–1969)
Adayr Eiras de Araújo (1970–1971)
João Luiz Campos Soares (1972–1973)
Jorge de Marsillac (1974–1975)
Alberto Lima de Moraes Coutinho (1976–1977)
João Sampaio Góis Jr. (1978–1982)
Hiram Silveira Lucas (1983–1986)
José Antonio Ribeiro Filho (1987–1989)
Antônio S. S. Figueira Filho (1990–1992)
Marconi Menezes Luna (1993–1995)
Henrique Moraes Salvador Silva (1996–1998)
Alfredo Carlos S. D. Barros (1999–2001)
Ezio Novais Dias (2002–2004)
Diógenes Luiz Basegio (2005–2007)
Carlos Ricardo Chagas (2008–2010)
Carlos Alberto Ruiz (2011–2013)
Ruffo de Freitas Júnior (2014–2016)
Antonio Luiz Frasson (2017–2019)
Vilmar Marques de Oliveira (2020–2022)



BRAZILIAN SOCIETY OF MASTOLOGY

Praça Floriano, 55, sala 801, Centro – 20031-050 – Rio de Janeiro (RJ)
Phone number: (21) 97271-0192
E-mail: secretaria@sbmastologia.com.br

ABOUT

Mastology is a continue publication of the Brazilian Society of Mastology. The responsibility for concepts emitted in the articles is exclusive of its authors. The total or partial reproduction of the articles is allowed, provided the source is mentioned.

Founder: Antônio Figueira Filho

Submissions - mailing address: Praça Floriano, 55, sala 801, Centro – Rio de Janeiro (RJ) – 20031-050

National and international subscription and advertising: Sociedade Brasileira de Mastologia - Phone number: (21) 97271-0192

NATIONAL BOARD OF DIRECTORS OF SOCIEDADE BRASILEIRA DE MASTOLOGIA

Triennium 2023-2025

Founder: Alberto Lima de Moraes Coutinho

President	Augusto Tufi Hassan
National Vice-President	Cícero de Andrade Urban
General Secretary	Roberto Kepler da Cunha Amaral
Assistant Secretary	Annamaria Massahud Rodrigues dos Santos
General Treasurer	Rosemar Macedo Sousa Rahal
Assistant Treasurer	André Mattar
North Region Vice-President	Ewaldo Lúzio Fôro de Oliveira
Northeast Region Vice-President	Maciel de Oliveira Matias
South Region Vice-President	Felipe Pereira Zerwes
Southeast Region Vice-President	Mônica Vieira M. Travassos Jourdan
Midwest Region Vice-President	Rodrigo Pepe Costa
Mastology Editor	Rafael Henrique Szymanski Machado
Escola Brasileira de Mastologia Director	Guilherme Novita Garcia
Deliberative Council President	Vilmar Marques de Oliveira
Mastology Specialist Title (TEMa) Committee	Francisco Pimentel Cavalcante
Ethics Committee	Clécio Ênio Murta de Lucena
Scientific Commission	Gil Facina

EDITORIAL PRODUCTION



ZEPPELINI
PUBLISHERS

Epidemiological and clinicopathological parameters related to the neoadjuvant chemotherapy for breast cancer during the COVID-19 crisis

Maria Fernanda da Motta Sperotto Valadares Gontijo^{1*}, Letícia Martins de Araújo Campos Linhares¹,
Luisa Lazarino de Souza Campos¹, Marcela de Oliveira Sá¹, Caroline Cançado Avelar¹,
João Pedro Costa Apolinário¹, Paulo Henrique Costa Diniz¹, José Tadeu Campos de Avelar¹

ABSTRACT

Introduction: Breast cancer, the second-leading cause of cancer-related deaths among women worldwide, is a complex and heterogeneous disease. Its socioeconomic aspects are recognized as determinants of clinical outcomes. The COVID-19 crisis negatively affected millions, particularly in impoverished macroregions like Brazil. Thus, influences on breast cancer patients' journey may occur, particularly in the neoadjuvant settings, in which a coordinated and multidisciplinary approach is mandatory. The present study aimed to analyze the epidemiological and clinicopathological profile of breast cancer patients who underwent neoadjuvant chemotherapy during the pandemic in Brazil. **Methods:** This is a unicentric, retrospective, and descriptive cross-sectional study conducted by analyzing data obtained from electronic medical records of breast cancer patients who underwent neoadjuvant chemotherapy. **Results:** From March 2020 to December 2022, 55 patients underwent neoadjuvant chemotherapy. They presented an average age of 50.0 years (range 43.9–47.6). About 83.6% of the tumors were invasive ductal carcinomas, and the most prevalent molecular subtype was hormone receptor-positive. T2 tumors accounted for 50.9%, while compromised N1 axillary lymph nodes represented 52.7%. The most commonly used neoadjuvant chemotherapy combined anthracyclines, cyclophosphamide, and sequential taxane. Regarding postoperative pathological response, 42.2% showed a partial response after neoadjuvant treatment, and a complete pathological response of as high as 40.0% occurred. The luminal and hybrid luminal subtypes were those that achieved the greatest response to neoadjuvant therapy. The lack of pathological response was only found in the luminal molecular subtype. **Conclusions:** This study demonstrated the impacts of the COVID-19 pandemic on breast cancer patients' journey. During this period of disruption in healthcare assistance, the disease presented at more advanced stages, but the pathologic complete response was higher than expected, and influences on chemotherapy decisions were not relevant. Overall, there were efforts to keep patients in the best breast cancer care.

KEYWORDS: breast neoplasms; breast; neoadjuvant therapy; COVID-19.

INTRODUCTION

Breast cancer, the second-leading cause of cancer-related deaths among women worldwide, is a major public health issue. Considering the complexity of this malignancy, socioeconomic inequalities may impact its relevant clinical outcomes. During the COVID-19 pandemic, whose beginning was recognized by the World Health Organization on March 20, 2020, approximately 90% of countries experienced setbacks in healthcare services, particularly in elective outpatient care, routine examinations, and complementary propaedeutics, focusing on emergency

care¹. It was even more evident in the most vulnerable populations, which includes Brazil, a continental developing country².

As a result of delays in the cancer patients' journey, from screening to treatment, it was reported an increase in the diagnosis of breast cancer in locally advanced stages, which requires multimodal therapies, notably the combination of neoadjuvant chemotherapy, surgery, and radiation therapy³.

By addressing micrometastases early, neoadjuvant chemotherapy aims to downstage the tumor to enable conservative surgery, evaluate *in vivo* response to systemic therapy, obtain

¹Hospital Mater Dei – Belo Horizonte (MG), Brazil.

*Corresponding author: mariafernandasperotto@gmail.com

Conflict of interests: nothing to declare. Funding: none.

Received on: 02/22/2024. Accepted on: 03/21/2025.

prognostic information, reduce the need for axillary dissection in cases of clinically positive axilla at the initial diagnosis, and provide time for surgical planning and genetic counseling⁴. Randomized prospective studies have shown the benefits of neoadjuvant systemic therapy compared to adjuvant therapy in operable tumors (T1–T3, N0–N1, M0), but with no differences in overall survival.

In the decision-making process involved in breast cancer, phenotypic subtype, histopathological aspects, and epidemiological data such as age and menopausal status should be considered. As these data might have changed significantly during the pandemic, this study analyzes clinical, epidemiological, and histopathological aspects of breast cancer patients treated with neoadjuvant chemotherapy in the COVID-19 pandemic period in Brazil.

METHODS

Patients undergoing neoadjuvant chemotherapy for breast cancer treatment at Mater Dei Hospital, a private referral center in Belo Horizonte, Brazil, between March 2020 and December 2022, were selected. Demographic and clinical data, such as age, menopausal status, family history of cancer, axillary staging, chemotherapy regimen, and pathological response to chemotherapy were gathered from electronic medical records. Histopathological data were obtained, including information on histology subtype, histological grade, tumor size, expression of hormonal receptors, human epidermal growth factor receptor-type 2 (HER2) status, Ki67 protein, molecular subtype, and degree of histological response to neoadjuvant treatment. No pathological review was performed for this analysis.

Inclusion criteria were female gender, age 18 and above, histologically confirmed diagnosis of breast cancer, multifocal disease, availability of data in electronic medical records, and neoadjuvant chemotherapy. Synchronic carcinomas were excluded, according to Figure 1.

Descriptive statistics were used to summarize the data. Normality tests (Shapiro-Wilk) were performed for each continuous

variable. Categorical variables were presented as numbers and percentages, and continuous variables as medians and interquartile ranges. Statistical analysis was conducted using Statistical Package for Social Sciences - SPSS® software, version 20 (SPSS, Chicago, IL).

The study was approved by an independent Ethics Committee (CAAE 73246223.8.0000.5128), and the protocols followed the 1975 Helsinki Declaration ethical guidelines. Due to the retrospective nature of this study, the local Human Subjects Committee approved the waiver of participants' free and informed consent.

RESULTS

Demographic data, clinical characteristics, and histopathological parameters of patients and their respective tumors were summarized in Table 1. Among the patients included in the study (n=237), 55 (22.0%) underwent neoadjuvant chemotherapy, with an average age of 50 years (range 43.9–47.6). Approximately 60% of these patients did not have a relevant family oncologic history, such as a history in first-degree relatives with breast, ovarian, or intestinal cancer. High penetrance gene mutations were found in two patients (TP53 and BRCA 2), and one had a variant of uncertain significance in the POLD1 gene.

Regarding histological aspects, approximately 83.6% of the tumors were invasive ductal carcinomas, and the most prevalent molecular subtype was hormone receptor-positive tumors. The luminal subtype (either A or B) comprised about 45.4% of the analyzed cases. T2 tumors (> 2 to 5 cm) accounted for 50.9%, followed by T3 tumors (larger than 5 cm) at 29.0%. Axillary involvement was found in 52.7% of patients, with mobile and fixed lymph nodes in the axilla ipsilateral to the tumor (respectively, N1 and N2), the majority classified as clinical staging II B. The most commonly used neoadjuvant regimen was a combination of anthracyclines, cyclophosphamide, and sequentially taxane (52.7%).

Partial pathological response after neoadjuvant treatment was seen in 42.2%, and complete pathological response (pCR) in 40.0%. When analyzing molecular subtypes, HER2-positive and hybrid luminal, patients had the highest complete response rates (80% and 50%, respectively), as show in Figure 2. The absence of pathological response to chemotherapy was found only in patients with the luminal molecular subtype, accounting for 20% of all analyzed luminal subtype patients.

DISCUSSION

The present study shows that, during the COVID-19 pandemic period, roughly 22.0% of all breast cancer patients underwent neoadjuvant therapy. Most (83.6%) presented as T2 or above, and clinical axillary involvement was detected in approximately 63.7%. Despite the challenges of keeping the patients at home in this period, the most used chemotherapy regimen




Figure 1. Trial profile.

Table 1. Patients with analyzed outcomes, in which the information was available, expressed as absolute numbers and percentage.

Characteristic	Number (n=55)	%
Age		
<40	11	20.0
40–49	18	32.7
50–64	19	34.5
≥65	7	12.7
Family History		
Positive	22	40.0
Negative	33	60.0
Histology		
Ductal	46	83.6
Lobular	5	9.0
Ductal and Lobular	3	5.4
Others	1	1.8
Molecular Subtype		
Luminal	25	45.4
HER2	5	9.0
Triple Negative	10	18.1
Hybrid Luminal	15	27.2
Tumor Size		
T1	9	16.3
T2	28	50.9
T3	16	29.0
T4	2	3.6
Lymph Node Status		
N0	20	36.3
N1	29	52.7
N2	6	10.9
N3	0	0
Therapy		
ddAC-T	29	52.7
ddAC-THP	10	18.1
ddAC	7	12.7
AC-TC	4	7.2
THP	3	5.4
TCHP	2	3.6
Pathological Response		
Complete	22	40
Luminal	7	27.2
HER2	4	18.1
Triple Negative	3	13.6
Hybrid Luminal	8	36.3
Partial	26	47.2
Luminal	13	50
HER2	1	3.8
Triple Negative	5	19.2
Hybrid Luminal	7	29.9
Absent		9.0
Unknown		3.6

HER2: human epidermal growth factor receptor-type 2; ddAC-T: dose-dense anthracycline plus cyclophosphamide and sequential paclitaxel; ddAC-THP: dose-dense anthracycline plus cyclophosphamide and sequential paclitaxel plus double blockade of trastuzumab and pertuzumab; AC-TC: anthracycline plus cyclophosphamide and sequential paclitaxel plus carboplatin; THP: paclitaxel plus double blockade of trastuzumab and pertuzumab; TCHP: paclitaxel plus carboplatin plus double blockade of trastuzumab and pertuzumab,

was ddAC-T, instead of those that required fewer days for infusions, such as a combination of taxanes and carboplatin or cyclophosphamide. Moreover, the frequency of HER2 tumors was higher than usual, and pathological response rates (partial or complete) in this subgroup were more common than the other molecular subtypes.

Breast cancer is the leading cause of cancer-related deaths in Brazil, except in the Northern, considered a socioeconomic less favorable geographic region⁵. Many epidemiological and clinicopathologic characteristics are associated with relevant clinical outcomes of this malignancy and need to be pointed out.

First, breast cancer incidence rises with age, thus being less common among younger women. Most cases are diagnosed in women aged 50–64, consistent with the predominant age group in our study (34.54% of patients)⁶.

Approximately 10–15% of breast cancers are associated with genetic alterations⁷. The Breast Cancer Association Consortium publication demonstrated an association between nine genes and breast cancer risk. Genes considered high-risk include BRCA1, BRCA2, PALB2, and TP53⁸. In our cohort, 40% had a family history of cancer in first-degree relatives, but only three had genetic mutations, two in high-risk genes (TP53 and BRCA2).

Over the last years, with a greater understanding of tumor molecular biology, breast cancer treatment has become increasingly complex, primarily guided by the subtype. A multidisciplinary approach becomes fundamental for treatment decisions for locally advanced cancer cases, defined as a tumor measuring over 2 cm (T2) and involving lymph nodes (N+). Almost 23% of patients underwent neoadjuvant therapy. Of those, 84.43% had T2, T3, or T4 tumors, and approximately 64% had clinically positive axilla. Due to screening failure or delay in searching for non-COVID-19-related medical assistance, we would expect a higher number of locally advanced tumors under neoadjuvant treatment. However, it is essential to mention the existence of different waves of COVID-19 cases⁹. This profile would probably be worse before vaccinations or when more new cases were reported. The resistance of patients and doctors to undergo chemotherapy during uncontrolled phases of the pandemic may also explain these findings.

In the past 40 years, medications and therapies have been developed to improve the quality of life and long-lasting outcomes for breast cancer patients. In this context, neoadjuvant treatment has emerged as a therapeutic strategy for surgical downstaging, *in vivo* assessment of systemic therapy response, and prognostic evaluation¹⁰. A study at the Memorial Sloan Kettering Cancer Center, between 2013 and 2019, revealed that of breast cancer patients with clinical stage I to III undergoing neoadjuvant chemotherapy, 75% presented a conversion from infeasible to feasible conservative surgery¹¹.

Subsequently, two major studies conducted in the United States in the 1990s demonstrated the non-inferiority of

neoadjuvant compared to adjuvant therapy regarding overall survival and progression-free survival. It was shown that patients achieving pCR had a better prognosis than those with residual disease⁴. The evaluation of this surrogate outcome as a reliable parameter was conducted in a meta-analysis published in 2014, correlating pCR with increased overall survival. This association became even more statistically evident in HER2+ patients, regardless of hormonal status, and triple-negative cases, confirmed in our study, where this group represents 84.4% of patients achieving pCR¹². Another critical point to be explored is the high frequency of HER2+ patients in our cohort. Possibly, these patients were more often referred to neoadjuvant treatment due to advances in antiHER2 treatment in this scenario.

Today, the standard therapy for initial HER2+ subtype breast cancer patients is neoadjuvant therapy, comprising different chemotherapy regimens associated with trastuzumab with or without pertuzumab¹³⁻¹⁵. In our institution's study, 36.6% of patients were hybrid luminal or HER2+ types. Of these patients, 50%

underwent neoadjuvant treatment with ddAC-THP, while the others were treated with de-escalated neoadjuvant protocols such as THP or TCHP, with 54.4% achieving pCR. Interestingly, these data point to the fact that the COVID-19 period did not interfere with medical decisions for regimens requiring fewer infusions despite the attempts to keep patients at home.

For triple-negative breast cancer (TNBC), current evidence indicates that treatments used in adjuvant therapy are also suitable for neoadjuvant settings. Based on the Anthracycline in Breast Cancer study, neoadjuvant treatment in TNBC patients is recommended for at least T1c and N+ using anthracycline and taxane-based chemotherapy^{16,17}. In our study, ten patients (18.8%) had TNBC in our institution.

None of the TNBC patients received neoadjuvant immunotherapy despite the data from the KEYNOTE-522 study. Published in 2020, this phase 3 trial demonstrated that combining pembrolizumab with carboplatin and paclitaxel followed by anthracycline in stage II or III TNBC patients resulted in higher pCR rates (64.8% vs. 51.2%)¹⁸. However, pembrolizumab in this scenario




Figure 2. Analysis of pathological response according to molecular subtypes.

was only approved in May 2022 by regulatory agencies in Brazil. It needs to be clarified whether the pandemic delayed our country's approval process.

Hormone receptor-positive (HR+) and HER2- tumors, despite their high prevalence, have more restricted indications for neoadjuvant therapy compared to other histological subtypes. However, it was a useful strategy worldwide during the COVID-19 period, when temporary contraindications for surgery were required. Despite that, we expected more patients with this condition at our center. Concerns regarding virus exposition during chemotherapy may have been balanced.

It is known that neoadjuvant chemotherapy is less effective in achieving pCR in luminal tumors, especially in the luminal A subtype, compared to more aggressive histologies¹⁹. However, our data showed that a pCR rate of around 25% was observed, surpassing global data of around 6–11% in the literature²⁰. Another intriguing finding is the absence of neoadjuvant endocrine therapy among our cohort, despite data showing pCR at least equivalent to chemotherapy ones²¹.

Our study has some limitations. It was retrospective and performed in a single center, not reflecting our population's sociodemographic and genetic diversity. Moreover, information and selection bias may have occurred. However, it was an important study to assess the impact of the COVID-19 crisis

among our patients, which analyzed epidemiologic and clinical pathological aspects.

CONCLUSION

The study contributed to a better understanding of the epidemiological profile of breast cancer patients who underwent neoadjuvant chemotherapy during the COVID-19 crisis when there was a disruption in healthcare assistance. Despite concerns regarding the pandemic itself, it was shown the effort to keep patients on the best assistance directed to breast cancer.

AUTHORS' CONTRIBUTION

MFMSVG: Conceptualization, Investigation, Methodology, Project administration, Validation, Visualization, Writing – original draft, Writing – review & editing. LMACL: Data curation, Writing – original draft, Writing – review & editing. LLSC: Data curation, Formal analysis, Writing – original draft, Writing – review & editing. MOS: Formal analysis, Investigation, Writing – review & editing. CCA: Formal analysis, Investigation. JPCA: Data curation, Validation. PHCD: Formal analysis, Investigation, Methodology, Project administration, Visualization. JTCA: Conceptualization, Methodology, Project administration, Visualization.









REFERENCES

1. Rocha R, Atun R, Massuda A, Rache B, Spinola P, Nunes L, et al. Effect of socioeconomic inequalities and vulnerabilities on health- system preparedness and response to COVID-19 in Brazil: a comprehensive analysis. *Lancet Glob Health*. 2021;9(6):e782-e792. [https://doi.org/10.1016/S2214-109X\(21\)00081-4](https://doi.org/10.1016/S2214-109X(21)00081-4)
2. Antonini M, Pinheiro DJPC, Matos ABTMB, Ferrado O, Mattar A, Okumua LM, et al. Impact of the COVID-19 pandemic on the breast cancer early diagnosis program in Brazil. *Prev Med Rep*. 2023;32:102157. <https://doi.org/10.1016/j.pmedr.2023.102157>
3. Korde LA, Somerfield MR, Carey La, Crews JR, Denduluri N, Hwang ES, et al. Neoadjuvant Chemotherapy, Endocrine Therapy and Targeted Therapy for Breast Cancer: ASCO Guideline. *J Clin Oncol*. 2021;39(13):1485-505. <https://doi.org/10.1200/JCO.20.03399>
4. Rastogi P, Anderson SJ, Bear HD, Geyer CE, Kahlenberg MS, Robidoux A, et al. Preoperative chemotherapy: updates of National Surgical Adjuvant Breast and Bowel Project protocols B-18 and B-27. *J Clin Oncol*. 2008;26(5):778-85. <https://doi.org/10.1200/JCO.2007.15.0235>
5. Perou CM, Sørle T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA, et al. Molecular portraits of human breast tumours. *Nature*. 2000;406(6797):747-52. <https://doi.org/10.1038/35021093>
6. Ministério da Saúde. Instituto Nacional de Câncer. Estimativa | 2023: incidência de câncer no Brasil. Rio de Janeiro: INCA, 2022.
7. Liu L, Hao X, Song Z, Zhi X, Zhang S, Zhang J. Correlation between family history and characteristics of breast cancer. *Sci Rep*. 2021;11(1):6360. <https://doi.org/10.1038/s41598-021-85899-8>
8. Breast Cancer Association Consortium; Dorling L, Carvalho S, Allen J, González-Neira A, Luccarini C, Wahlström C, et al. Breast Cancer Risk Genes — Association Analysis in More than 113,000 Women. *N Engl J Med*. 2021;384(5):428-39. <https://doi.org/10.1056/NEJMoa1913948>
9. Moura EC, Cortez-Escalante J, Cavalcante FV, Barreto ICHC, Sanchez MN, Santos LMP. Covid-19: evolução temporal e imunização nas três ondas epidemiológicas, Brasil, 2020-2022. *Rev Saúde Pública*. 2022;56:105. <https://doi.org/10.11606/s1518-8787.2022056004907>
10. Rubens RD, Sexton S, Tong D, Winter PJ, Knight RK, Hayward JL. Combined chemotherapy and radiotherapy for locally advanced breast cancer. *Eur J Cancer* 1980;16(3):351-6. [https://doi.org/10.1016/0014-2964\(80\)90352-7](https://doi.org/10.1016/0014-2964(80)90352-7)
11. Petruolo O, Sevilimedu V, Montagna G, Le T, Morrow M, Barrio AV. How Often Does Modern Neoadjuvant Chemotherapy Downstage Patients to Breast-Conserving Surgery? *Ann Surg Oncol*. 2021;28(1):287-94. <https://doi.org/10.1245/s10434-020-08593-5>
12. Cortazar P, Zhang L, Untch M, Mehta K, Costantino JP, Wolmark N, et al. Pathological complete response and

- long-term clinical benefit in breast cancer: the CTNeoBC pooled analysis. *Lancet*. 2014;384(9938):164-72. [https://doi.org/10.1016/S0140-6736\(13\)62422-8](https://doi.org/10.1016/S0140-6736(13)62422-8)
13. Gianni L, Eiermann W, Semiglazov V, Manikhas A, Lluch A, Tjulandin S, et al. Neoadjuvant chemotherapy with trastuzumab followed by adjuvant trastuzumab versus neoadjuvant chemotherapy alone, in patients with HER2-positive locally advanced breast cancer (the NOAH trial): a randomised controlled superiority trial with a parallel HER2-negative cohort. *Lancet*. 2010;375(9712):377-84. [https://doi.org/10.1016/S0140-6736\(09\)61964-4](https://doi.org/10.1016/S0140-6736(09)61964-4)
 14. Schneeweiss A, Chia S, Hickish T, Harvey V, Eniu A, Hegg R, et al. Pertuzumab plus trastuzumab in combination with standard neoadjuvant anthracycline-containing and anthracycline-free chemotherapy regimens in patients with HER2-positive early breast cancer: a randomized phase II cardiac safety study (TRYPHAEANA). *Ann Oncol*. 2013;24(9):2278-84. <https://doi.org/10.1093/annonc/mdt182>
 15. Gianni L, Pienkowski T, Im YH, Roman L, Tseng LM, Liu MC, et al. Efficacy and safety of neoadjuvant pertuzumab and trastuzumab in women with locally advanced, inflammatory, or early HER2-positive breast cancer (NeoSphere): a randomised multicentre, open-label, phase 2 trial. *Lancet Oncol*. 2012;13(1):25-32. [https://doi.org/10.1016/S1470-2045\(11\)70336-9](https://doi.org/10.1016/S1470-2045(11)70336-9)
 16. Von Minckwitz G, Schneeweiss A, Loibl S, Salat C, Denkert C, Rezai M, et al. Neoadjuvant carboplatin in patients with triple-negative and HER2-positive early breast cancer (GeparSixto; GBG 66): a randomised phase 2 trial. *Lancet Oncol*. 2014;15(7):747-56. [https://doi.org/10.1016/S1470-2045\(14\)70160-3](https://doi.org/10.1016/S1470-2045(14)70160-3)
 17. Early Breast Cancer Trialists' Collaborative Group (EBCTCG). Long-term outcomes for neoadjuvant versus adjuvant chemotherapy in early breast cancer: meta-analysis of individual patient data from ten randomised trials. *Lancet Oncol*. 2018;19(1):27-39. [https://doi.org/10.1016/S1470-2045\(17\)30777-5](https://doi.org/10.1016/S1470-2045(17)30777-5)
 18. Schmid P, Cortes J, Pusztai L, McArthur H, Kümmel S, Bergh J, et al. Pembrolizumab for Early Triple-Negative Breast Cancer. *N Engl J Med*. 2020;382(9):810-21. <https://doi.org/10.1056/NEJMoa1910549>
 19. Kim HS, Yoo TK, Park WC, Chae BJ. Potential Benefits of Neoadjuvant Chemotherapy in Clinically Node-Positive Luminal Subtype - Breast Cancer. *J Breast Cancer*. 2019;22(3):412-24. <https://doi.org/10.4048/jbc.2019.22.e35>
 20. Kim HJ, Noh WC, Lee ES, Jung YS, Kim LS, Han W, et al. Efficacy of neoadjuvant endocrine therapy compared with neoadjuvant chemotherapy in pre-menopausal patients with oestrogen receptor-positive and HER2-negative, lymph node-positive breast cancer. *Breast Cancer Res*. 2020;22(1):54. <https://doi.org/10.1186/s13058-020-01288-5>
 21. Semiglazov VF, Semiglazov VV, Dashyan GA, Ziltsova EK, Ivanov VG, Bozhok AA, et al. Phase 2 randomized trial of primary endocrine therapy versus chemotherapy in postmenopausal patients with estrogen receptor-positive breast cancer. *Cancer*. 2007;110(2):244-54. <https://doi.org/10.1002/cncr.22789>



Telehealth after one year of Breast Cancer Surgery as a Physical Therapy Follow-up Strategy

Josélia Fiorin Casagrande¹ , Flávia Oliveira Macedo¹ , Erica Alves Nogueira Fabro¹ ,
Flávia Orind Ferreira¹ , Daniele Medeiros Torres¹ , Juliana Flávia de Oliveira Tavares de Oliveira^{1,2} ,
Simone Abrantes Saraiva¹ , Rejane Medeiros Costa^{1*} 

ABSTRACT

Introduction: Breast cancer is the most prevalent type of invasive cancer in the female population. The surgical procedure is one of the aspects of oncological treatment. However, there are several postoperative complications resulting from this process, in which physical therapists work from prevention to treatment. During the COVID-19 pandemic, the physical therapy service implemented teleconsultations for the six-month and one-year postoperative follow-ups. The purpose of this article was to evaluate the prevalence of complications identified in physical therapy follow-up, through teleconsultation, and to describe the frequency of face-to-face evaluations to confirm the physical therapy diagnosis. **Methods:** This is a cross-sectional study, including patients submitted to surgical treatment of breast cancer, from January 2019 to September 2021, and who were seen at the one-year postoperative teleconsultation. **Results:** We included 362 patients, with a mean age of 58.17 (± 12.16) years. Among the reported complications, the most frequent was paresthesia in the intercostobrachial nerve (87.1%); 23.8% of the patients reported pain; and 22.1%, phantom breast sensation. 11.9% ($n=43$) of the patients were referred for face-to-face evaluation, being 58.1% ($n=25$) due to the perception of lymphedema as the main reason for these referrals. **Conclusions:** The most frequent complications reported by patients in one-year teleconsultation were paresthesia, followed by pain and sensation of phantom breast. The greatest reason for referrals to face-to-face consultation was lymphedema. With such findings, this modality of care shows a possibility of effective follow-up in the postoperative period of breast cancer.

KEYWORDS: teleconsultation; breast cancer; physical therapy.

INTRODUCTION

According to the Brazilian National Cancer Institute (*Instituto Nacional de Câncer* – INCA), there are, for each year of the triennium 2023–2025, 73,610 thousand new estimated cases of breast cancer in Brazil, this being the most frequent type of invasive cancer in the female population¹. The surgical procedure consists of the pillar of oncological treatment for breast cancer, and among the various complications resulting from this process are pain, sensitivity change, reduced range of motion (ROM), scar complications, sensation and pain in the phantom breast, axillary web syndrome (AWS), and lymphedema in the homolateral upper limb (HUL) in relation to surgical treatment. The physical therapist intervenes in the prevention, early detection, and treatment of these complications².

Researchers reinforce that the guidelines for the practice of free active movement and strengthening of the upper limbs are

indispensable for kinetic-functional recovery and pain control after oncological breast surgery^{3–6}. In addition, early postoperative physical therapy with exercises for the shoulder joint showed both pain reduction and improvement of ROM, and recovery of upper limb functionality for daily and labor activities⁷.

Social distancing, in early 2020, was recommended due to the advance of the SARS-CoV-2 virus pandemic, in such a way it was necessary to reduce face-to-face consultations and care⁸. In order to guide healthcare professionals, the Brazilian Federal Council of Physical Therapy and Occupational Therapy (*Conselho Federal de Fisioterapia e Terapia Ocupacional* – COFFITO), through Resolution 516 of March 20, 2020, allowed the online care provided in the modalities of teleconsultation, teleconsulting, and telemonitoring by physical therapists and occupational therapists⁹.

¹Instituto Nacional de Câncer – Rio de Janeiro (RJ), Brazil.

²Hospital Federal de Bonsucesso – Rio de Janeiro (RJ), Brazil.

*Corresponding author: rmcosta2@gmail.com

Funding: none.

Conflicts of interest: nothing to declare.

Received on: 06/26/2024. **Accepted on:** 03/21/2025.

According to Federal Law No. 14,510/2022, which authorizes the practice of telehealth throughout the Brazilian territory, telehealth consists in the provision of distance healthcare services, using communication technologies through texts, sounds, and/or images for health information, and ensures that every healthcare professional has the autonomy to decide whether or not to use telehealth whenever deemed necessary¹⁰.

The physical therapy service of the Hospital do Câncer III [Cancer Hospital III] of the Instituto Nacional de Câncer (HC III/INCA), a hospital unit specialized in the treatment of breast cancer, has established in its routine the following physical therapeutic evaluations: first time (before starting the oncological treatment), on the first day, in 30 days, six months, and one year after surgery¹¹. The physical therapy consultation of one year after breast cancer surgery is the time to evaluate the possibility of discharge of outpatient physical therapy follow-up in HC III/INCA.

In the initial period of the COVID-19 pandemic, the physical therapy service implemented the practice of telehealth, and consultations of 30 days, six months, and one year after the surgery were carried out in the form of teleconsultation. If face-to-face care was necessary, the recommendations of the Brazilian Association of Physical Therapy in Oncology (*Associação Brasileira de Fisioterapia em Oncologia* – ABFO) for prevention and biosafety against the spread of SARS-CoV-2 were followed¹². With the control of the pandemic and the return of face-to-face care, the teleconsultation was maintained for the follow-ups of six months and one year after surgery because it allows more comfort and economy to patients¹³.

In this study, our objective was to describe the prevalence of complications evaluated in physical therapy follow-up through one-year post-operative teleconsultation of breast cancer and to analyze the frequency and need for face-to-face evaluations to confirm the physical therapy diagnosis.

METHODS

This is a cross-sectional study with patients submitted to surgical treatment of breast cancer in HCIII/INCA, from January 2019 to September 2021, and who were seen via teleconsultation for evaluation and physical therapy follow-up in the one-year postoperative period.

The eligible patients were preselected through active search of the evolutions of consultations in the Care Control System of the Physical Therapy Service (*Sistema de Controle do Atendimento do Serviço de Fisioterapia* – SISCASF). Sociodemographic, clinical, and treatment data were collected from paper and electronic medical records. In addition, the following data were collected from: the physical therapeutic evolution record of the one-year postoperative teleconsultation, the postoperative complications reported, such as pain, sensation and/or pain in the phantom breast, paresthesia in the cutaneous region innervated by the intercostobrachial nerve (ICBN), intercostobrachial neuralgia (ICB), axillary web syndrome (AWS), reduced range of motion (ROM), subjective sensation of

lymphedema (sensation of weight and swelling), and lymphedema (patients' perception of change in the volume of HUL).

To evaluate life habits, information regarding personal care and household chores were collected from the physical therapeutic evaluation form, classified as fully carried out, partially or not carried out, practice of home exercises with upper limbs (UL) and the frequency of these exercises, classified as regular, irregular, or not performing it, and practice of leisure physical exercises.

For certain types of complaints reported in teleconsultation, such as lymphedema in HUL, severe pain, reduced ROM, and AWS, the patient was referred to face-to-face consultation. The reason for referral to face-to-face evaluation was collected from physical therapeutic evolutions.

Of the patients referred to face-to-face evaluation, the same complications evaluated in the teleconsultation were collected, and the assessment of lymphedema was verified through perimetry of the upper limbs. Lymphedema was considered when the difference between HUL and the contralateral upper limb was ≥ 2.0 cm in at least one reference point.

To evaluate the perimetry of the UL, a tape measure was used and the circumference of the limb was measured at six specific points, demarcating the first point in the region of the lateral epicondyle and the others every 7-cm distance to the arm and forearm.

Descriptive analyses of measures of central tendency and dispersion were performed for continuous variables and absolute frequency for categorical variables. The Statistical Package for Social Sciences (SPSS), version 21.0, was used for data analysis.

This study was approved by the Research Ethics Committee (*Comitê de Ética e Pesquisa* – CEP) of INCA under number 4.702.209.

RESULTS

We included 362 patients, with a mean age of 58.17 (± 12.16) years. Most were women (98.9%), self-reported to be Black and/or mixed-race (63.0%), did not live with a partner (57.7%), had level of education equal to or greater than eight years of formal education (69.0%), had household chores as the main occupation (69.3%), and resided in the "Metropolitana I" region of the state of Rio de Janeiro (RJ), Brazil (91.4%)¹⁴ (Table 1).

At the time of breast cancer diagnosis, 73.7% of patients were classified with locally advanced clinical staging (\geq IIB), but at the time of one-year postoperative teleconsultation, only 2.8% presented disease progression. Regarding surgical treatment, 82.3% of the patients underwent mastectomy and 88.1% underwent axillary lymphadenectomy. Only 3.1% of women submitted to mastectomy underwent immediate breast reconstruction with silicone prosthesis or tissue expander. Most were submitted to chemotherapy (89.0%), radiotherapy (88.7%), and hormone therapy (86.7%); and only 15.8%, to molecularly targeted therapy (Table 1).

At the time of the one-year teleconsultation, 82.3% of the patients were under treatment with hormone therapy; 3.0%,

radiotherapy; 3.0%, chemotherapy; and 4.1%, molecularly targeted therapy (Table 2).

Among the complications reported in the one-year physical therapy teleconsultation, the most frequent was paresthesia in the ICBN (87.1%); 23.8% of the patients reported pain and 22.1%, phantom breast sensation. The subjective sensation of lymphedema was reported by 4.1% of the patients and 6.9% noticed changes in the volume of HUL, characterizing lymphedema (Table 3).

As for life habits, most patients reported to be independent for personal care (99.7%), of which 79.3% carried out household chores alone, without assistance. Home exercises with upper limbs (UL), according to postoperative guidelines, were present in 83.1% of the population, with regular frequency of

50.1%. As for physical exercises, only 28.7% of the patients practiced them (Table 4).

During the teleconsultation, 43 women (11.9%) were referred to face-to-face evaluation at the HCIII/INCA physical therapy outpatient clinic (non-tabulated data). Regarding the reason for referral, 58.1% (n=25) reported lymphedema, due to the perception of alteration in the volume of HUL. Other complications confirmed in the face-to-face evaluation are shown in Table 5.

DISCUSSION

In this study, we identified that the main symptom reported by patients in the one-year teleconsultation after breast cancer surgery is paresthesia in the ICBN (87.1%), characterized by alteration of sensitivity in the medial region of the arm, axilla, and lateral trunk resulting from nerve injury^{15,16}.

As ICBN has close connection with axillary lymph nodes, the risk of injury is high during lymph node dissection¹⁶. The diagnosis of ICBN injury is clinical and, although numbness is very common, several events may occur in the distribution of the nerve due to changes in its function, such as tingling, burning and electrical sensations, in addition to pain^{15,16}.

According to Andersen *et al.*¹⁷, hypoesthesia was the main sensory dysfunction one year after surgery, affecting 85.0% of patients. The authors also sought a relationship between surgical treatment and sensory dysfunction, and observed that the areas of hypoesthesia were significantly greater for patients treated with mastectomy (p<0.0001) and axillary lymphadenectomy (AL; p<0.0001) compared to those treated with conservative surgery and sentinel lymph node biopsy (SLNB).

Table 1. Sociodemographic, clinical, and treatment data of the study population (n=362).

Variables	n (%) ^a	Variables	n (%)
Age (mean and SD)	58.17 (±12.16)	Disease progression*	
		No	352 (97.2)
		Local recurrence	3 (0.8)
		Distant metastasis	7 (2.0)
Sex		Breast surgery	
Women	358 (98.9)	Conservative	63 (17.4)
Men	4 (1.1)	Mastectomy	298 (82.3)
		No approach**	1 (0.3)
Skin color		Axillary approach	
Black and/or mixed-race	228 (63.0)	SLNB	43 (11.9)
White	134 (37.0)	AL	319 (88.1)
Marital status		Immediate breast reconstruction***	
Have a partner	153 (42.3)	No	285 (96.9)
No partner	209 (57.7)	Yes	9 (3.1)
Level of education*		Chemotherapy	
<8 years	110 (31.0)	No	40 (11.0)
≥8 years	247 (69.0)	Yes	322 (89.0)
Occupation		Radiotherapy	
Housewife	251 (69.3)	No	41 (11.3)
Outside job	111 (30.7)	Yes	321 (88.7)
Residence region (RJ)		Hormone therapy	
Metropolitana I	331 (91.4)	No	48 (13.3)
Metropolitana II	19 (5.2)	Yes	314 (86.7)
Serrana	7 (2.0)		
Baixada litorânea	4 (1.1)		
Médio Paraíba	1 (0.3)		
Clinical staging		Targeted therapy	
0-IIA	95 (26.3)	No	305 (84.2)
≥IIB	266 (73.7)	Yes	57 (15.8)

SD: standard deviation; SLNB: sentinel lymph node biopsy; AL: axillary lymphadenectomy; RJ: Rio de Janeiro. *Patients who were under treatment due to progression of the disease at the time of the one-year teleconsultation; **exclusive axillary lymphadenectomy, without breast approach, for occult breast cancer; ***the difference in the total sample is due to the total number of women undergoing mastectomy; ^athe total value may change due to missing values.

Table 2. Treatment in progress at the time of the one-year teleconsultation after breast cancer surgery (n=362).

Variables	n (%) ^a
Treatment in progress	
No	52 (14.4)
Adjuvant	300 (82.9)
Palliative	10 (2.8)
Hormone therapy	
No	64 (17.7)
Yes	298 (82.3)
Chemotherapy	
No	351 (97.0)
Yes	11 (3.0)
Radiotherapy	
No	351 (97.0)
Yes	11 (3.0)
Targeted therapy	
No	347 (95.9)
Yes	15 (4.1)

^aThe same patient could be undergoing more than one treatment simultaneously, except in cases of chemotherapy and hormone therapy or chemotherapy and radiotherapy.

Lucena *et al.*¹⁸ carried out a cross-sectional cohort in the same institution of the present study and evaluated 182 women after one year of surgical treatment for breast cancer; they observed that 58.2% of the interviewees reported paresthesia in the ICBN. However, in the aforementioned study, the majority of patients only underwent SLNB (58.2%), while in our study the vast majority were submitted to AL (88.1%).

In the study by Siqueira *et al.*¹⁹, 47.2% of the interviewees reported sensory alteration related to ICBN injury after the mean time of 5.06 years (± 1.8) between surgery and evaluation. The difference in the results can be justified by the fact that the evaluation of patients in the research of Siqueira *et al.*¹⁹ was carried out in a longer interval, in addition to a higher percentage of SLNB (39.8%), compared to the present study.

Persistent pain or painful syndrome after treatment of breast cancer related to intercostobrachial neuralgia (pain in the ICBN)

or in the phantom breast is another common symptom, affecting 25–60% of patients, and has been associated with a decrease in quality of life.

In addition to surgery, adjuvant therapies, such as radiotherapy and systemic treatments, are also risk factors for painful syndrome, as they can cause damage to nerve fibers^{20–22}.

In the study sample, 23.8% of the patients complained of pain at the time of the physical therapy teleconsultation, most related to the upper limb homolateral to surgery (14.4%). Pain was the second most frequent reason (41.8%) for referral of patients to face-to-face outpatient consultation.

Authors of a meta-analysis comprised of 18 observational studies with 6,364 patients with persistent pain after breast cancer surgery found that the prevalence of this symptom was 31.0% (95%CI: 23–41%) in 1–2 years²³.

The complication responsible for the highest number of referrals to face-to-face consultation after one year of surgery

Table 3. Complications reported in the one-year teleconsultation after breast cancer surgery (n=362).

Variables	n (%)**
Paresthesia in the ICBN	
No	47 (12.9)
Yes	312 (87.1)
ICB	
No	347 (96.9)
Yes	11 (3.1)
AWS	
No	357 (98.6)
Yes	5 (1.4)
ROM	
No	359 (99.2)
Yes	3 (0.8)
Pain	
No	276 (76.2)
Yes	86 (23.8)
Pain site	
HUL	52 (14.4)
Chest wall/breast/thorax	17 (4.7)
Other	17 (4.7)
Does not apply	276 (76.2)
Phantom breast	
No	198 (56.9)
Yes	77 (22.1)
Does not apply**	73 (21.0)
Subjective sensation of lymphedema	
No	347 (95.9)
Yes	15 (4.1)
Lymphedema	
No	337 (93.1)
Yes	25 (6.9)

ICBN: intercostobrachial nerve; ICB: intercostobrachial neuralgia; AWS: axillary web syndrome; ROM: reduced range of motion; HUL: homolateral upper limb. *Each patient may have presented more than one complication; **women undergoing conservative surgery and immediate breast reconstruction; *the total value may change due to missing values.

Table 4. Life habits and home routine at the time of the one-year teleconsultation after breast cancer surgery (n=362).

Variables	n (%) ^a
Independence in personal care	
Yes	361 (99.7)
No	1 (0.3)
Household chores	
Carries it out completely	287 (79.3)
Carries it out partially	73 (20.2)
Does not carry it out	2 (0.6)
Practice of home exercises with UL	
Yes	301 (83.1)
No	61 (16.9)
Frequency of home exercises with UL	
Regular*	181 (50.0)
Irregular	120 (33.1)
Never	61 (16.9)
Practice of physical exercises	
Yes	100 (28.7)
No	248 (71.3)

UL: upper limbs. *At least once a day; *the total value may change due to missing values.

Table 5. Complications confirmed in the face-to-face consultation (43 women).

Variables	n (%)*
Lymphedema	25 (58.1)
Pain	18 (41.8)
Paresthesia in the ICBN	2 (4.6)
AWS	1 (2.3)
ROM	1 (2.3)

ICBN: intercostobrachial nerve; AWS: axillary web syndrome; ROM: reduced range of motion. *Each patient may have presented more than one complication.

was lymphedema, reported by 25 women (6.9% of the sample), through the perception of alteration in the volume of HUL. This prevalence was confirmed in the face-to-face evaluation.

According to recent research, the incidence of lymphedema after surgical treatment of breast cancer varies according to the characteristics of the studied population, being associated, in general, with high body mass index/obesity, a higher number of lymph nodes removed, radiotherapy on the lymph node chain, and taxane-based chemotherapy²⁴⁻²⁶.

In the study by Furlan *et al.*²⁷, whose objective was to evaluate the circumference and the sensation of swelling in the upper limb homolateral to surgery right after the procedure and within 24 months, the authors identified that, in the first year, of the 152 patients followed up, 23.7% had a feeling of limb edema; 21.1% had a difference greater than 2 cm at a single point; and 5.9%, circumference greater than 2 cm at two points, comparing the affected limb and contralateral limb.

In the present study, in the one-year teleconsultation, only 4.1% of patients reported a subjective sensation of lymphedema and 6.9% (n=25) reported lymphedema because they noticed changes in the volume of the affected limb, which was confirmed in all of these 25 patients in the face-to-face evaluation.

Konish *et al.*²⁸ found cumulative incidence of lymphedema in high-risk patients of approximately 3.0% in one year. Conversely, in the study by Paramanandam *et al.*²⁹, of the 149 patients guided as for the usual care with arm, skin, drain, and daily shoulder exercises since the first postoperative day, the cumulative incidence in one year was 25.0%. When evaluating 580 patients submitted to breast surgery and postoperative radiotherapy with or without systemic treatment, Kim *et al.*³⁰ found a cumulative incidence of 10.5% after one year of radiotherapy, but in this sample the majority of patients (84.5%) underwent conservative breast surgery and sentinel lymph node biopsy (78.4%), and less than half of the patients (37.4%) did not receive chemotherapy.

In the cohort study conducted by Fabro *et al.*³¹, on 174 women with a mean age of 58 years, 29.5% of the patients reported the subjective sensation of edema in the upper limb homolateral to surgery after approximately eight months postoperatively, corroborating our findings, although the evaluation time was slightly shorter in the aforementioned study³¹.

ICBN paresthesia and lymphedema were also the most common complications found in the study conducted by Abass *et al.*³² In a sample of 96 patients, the authors identified paresthesia as the most frequent complication (21.9%), followed by lymphedema (9.4%), in the average time of 18 months of follow-up.

Although most patients in the present study reported total independence for personal care (99.7%) and carrying out household chores, without assistance (79.3%), demonstrating being active in the day-to-day, when questioned about the practice of exercises with the UL, 83.1% stated practicing them as recommended since the physical therapy consultation of the first postoperative day,

but half of these (50.0%) followed a regular routine (at least once a day). Only 28.7% of the interviewees reported practicing some kind of sport activity at the time of teleconsultation.

Marchito *et al.*³³ also observed that patients submitted to surgical treatment of breast cancer adhere to the preventive physical therapy guidelines (skin care and exercise with UL) after surgery, especially in the early months, and that adherence to these guidelines reduced in the following months, mainly due to household chores.

Lee *et al.*³⁴, in a study carried out in Malaysia on the practice of global physical activity by breast cancer survivors, found that physical activity levels in this population were inadequate since diagnosis and that they significantly reduced ($p=0.04$) over three years after cancer discovery. Among the interviewees, 48.1% were active at the time of diagnosis, 39.8% in one year, and 35.3% in the third year.

Groef *et al.*³⁵ assessed the levels of global physical activity within two years after breast cancer surgery and found that in none of the domains (occupational, sports, and domestic) there was a return to preoperative activity levels.

The studies whose authors address teleconsultation as a modality of care show that this strategy is effective, accessible, and viable for monitoring patients in breast cancer treatment. Singleton *et al.*³⁶ suggested that interventions through teleconsultation had wide range, high acceptance by survivors of breast cancer, and were effective in improving quality of life, self-efficacy, fatigue, and psychological suffering.

Nápoles *et al.*³⁷ found positive evidence related to viability, acceptability, and efficacy, with significant improvement in fatigue, psychological suffering, and emotional well-being as well as benefits related to greater knowledge of recommended care. The authors also observed an improvement in symptoms and in the level of physical activity in breast cancer survivors. Macedo *et al.*¹³ evaluated patients' opinion on teleconsultations for follow-up of breast cancer, and showed good acceptance with patients feeling safe, satisfied, and comfortable.

A recent guideline on telerehabilitation in patients with breast cancer suggests that this modality is present from the initial moments of the preoperative period and through individualized programs, with prescription of exercises in the postoperative and at the long term³⁸. Moreover, authors of systematic reviews have shown that this form of care is cost-effective in public health, especially for people living in rural areas^{39,40}.

Although this is a cross-sectional study, with no information since the preoperative period, its strength is presenting the prevalence of the main complications one year after breast cancer surgery evaluated through teleconsultation. The predominant symptoms or complications described in this period may guide physical therapists and other healthcare professionals in their medium- and long-term conduct. In addition, we showed that the number of face-to-face consultations required after teleconsultation was low, which makes the online modality a viable resource after surgical treatment of breast cancer.

CONCLUSIONS

The most frequent complications reported by patients in the one-year teleconsultation after breast surgery were paresthesia, pain, and phantom breast sensation. The greatest reason for referrals to face-to-face consultation was lymphedema, with diagnostic confirmation of all cases. With such findings, this modality of care shows a possibility of effective follow-up in the postoperative period of breast cancer.

AUTHORS' CONTRIBUTION

JFC: Data curation, Formal analysis, Investigation, Methodology, Writing – original draft. FOM: Conceptualization, Data curation,

Formal analysis, Investigation, Methodology, Project administration, Supervision, Writing – original draft, Writing – review & editing. EANF: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Writing – original draft, Writing – review & editing. FOF: Data curation, Investigation, Writing – original draft, Writing – review & editing. DMT: Data curation, Investigation, Writing – original draft, Writing – review & editing. JFOTO: Data curation, Investigation, Writing – review & editing. SAS: Data curation, Investigation, Writing – review & editing. RMC: Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project administration, Supervision, Writing – original draft, Writing – review & editing.

REFERENCES

1. Santos M de O, Lima FC da S de, Martins LFL, Oliveira JFP, Almeida LM de, Cancela M de C. Estimativa de Incidência de Câncer no Brasil, 2023-2025. *Rev Bras Cancerol.* 2023;69(1):e-213700. <https://doi.org/10.32635/2176-9745.RBC.2023v69n1.3700>
2. Bergmann A, Ribeiro MJP, Pedrosa E, Nogueira EA, Oliveira ACG. Fisioterapia em mastologia oncológica: rotinas do Hospital do Câncer III / INCA. *Rev Bras Cancerol.* 2006;52(1):97-109. <https://doi.org/10.32635/2176-9745.RBC.2006v52n1.1906>
3. Bruce J, Mazuquin B, Mistry P, Rees S, Canaway A, Hossain A, et al. Exercise to prevent shoulder problems after breast cancer surgery: the PROSPER RCT. *Health Technol Assess* 2022;26(15):1-124. <https://doi.org/10.3310/JKNZ2003>
4. Ribeiro IL, Moreira RFC, Ferrari AV, Alburquerque-Sendín F, Camargo PR, Salvini TF. Effectiveness of early rehabilitation on range of motion, muscle strength and arm function after breast cancer surgery: a systematic review of randomized controlled trials. *Clin Rehabil.* 2019;33(12):1876-86. <https://doi.org/10.1177/0269215519873026>
5. Rizzi SKLA, Haddad CAS, Giron PS, Figueira PVG, Estevão A, Elias S, et al. Early Free Range-of-Motion Upper Limb Exercises After Mastectomy and Immediate Implant-Based Reconstruction Are Safe and Beneficial: A Randomized Trial. *Ann Surg Oncol.* 2020;27(12):4750-9. <https://doi.org/10.1245/s10434-020-08882-z>
6. De Groef A, Van Kampen M, Dieltjens E, Christiaens MR, Neven P, Geraerts I, et al. Effectiveness of postoperative physical therapy for upper-limb impairments after breast cancer treatment: a systematic review. *Arch Phys Med Rehabil.* 2015;96(6):1140-53. <https://doi.org/10.1016/j.apmr.2015.01.006>
7. Rett MT, Moura DP, Oliveira FB, Domingos HYB, Oliveira MMF, Gallo RBS, et al. Fisioterapia após cirurgia de câncer de mama melhora a amplitude de movimento e a dor ao longo do tempo. *Fisioter Pesqui.* 2022;29(1):46-52. <https://doi.org/10.1590/1809-2950/21001929012022PT>
8. Rizzi SKLA, Cerqueira MTAS, Gomes NO, Baiocchi JMT, Aguiar SS, Bergmann A. Nota Técnica da Associação Brasileira de Fisioterapia em Oncologia sobre os Atendimentos de Fisioterapia em Oncologia frente à Pandemia de Covid-19. *Rev Bras Cancerol.* 2020;66(TemaAtual):e-1973. <https://doi.org/10.32635/2176-9745.RBC.2020v66nTemaAtual.973>
9. Conselho Federal de Fisioterapia e Terapia Ocupacional (BR). RESOLUÇÃO nº 516, de 20 de março de 2020. Dispõe sobre a suspensão temporária do Artigo 15, inciso II e Artigo 39 da Resolução COFFITO nº424/2013 e Artigo 15, inciso II e Artigo 39 da Resolução COFFITO nº 425/2013 e estabelece outras providências durante o enfrentamento da crise provocada pela Pandemia do COVID-19. *Diário Oficial da União.* 23 mar. 2020.
10. BRASIL. Lei nº 14.510, de 27 de dezembro de 2022. Altera a Lei nº 8.080, de 19 de setembro de 1990, para autorizar e disciplinar a prática da tele saúde em todo o território nacional, e a Lei nº 13.146, de 6 de julho de 2015; e revoga a Lei nº 13.989, de 15 de abril de 2020. *Diário Oficial da União.* 28 dez. 2022.
11. Bergmann A, Ribeiro MJP, Pedrosa E, Nogueira EA, Oliveira ACG. Fisioterapia em Mastologia oncológica: rotinas do Hospital do Câncer III / INCA. *Rev Bras Cancerol.* 2006;52(1):97-109. <https://doi.org/10.32635/2176-9745.RBC.2006v52n1.1906>
12. Agência Nacional de Vigilância Sanitária. Nota Técnica GVIMS/ GGES/ANVISA nº 04/2020: orientações para serviços de saúde: medidas de prevenção e controle que devem ser adotadas durante a assistência aos casos suspeitos ou confirmados de infecção pelo novo coronavírus (SARS-CoV-2) [Internet]. Brasília, DF: ANVISA; 2020 [acessado em 2025 abr 10]. Disponível em: <https://www.gov.br/anvisa/pt-br/centraisdeconteudo/publicacoes/servicosdesaude/notas-tecnicas/notas-tecnicas-vigentes/NOTATCNICAGVIMS0420covid1925.06.2024.pdf>
13. Macedo FO, Ferreira FO, Torres DM, Saraiva SA, Oliveira JFOT de, Fabro EAN, et al. Acompanhamento Fisioterapêutico após Tratamento Cirúrgico do Câncer de Mama por Teleconsulta: Percepção e Adesão das Pacientes. *Rev Bras Cancerol.* 2023;69(4):e-074091. <https://doi.org/10.32635/2176-9745.RBC.2023v69n4.4091>
14. Kuschnir R, Chorny A, Lira AML, Sonoda G, Fonseca TMP. Regionalização no estado do Rio de Janeiro: o desafio de aumentar acesso e diminuir desigualdades. In: Ugá MAD et al. (orgs.). *A gestão do SUS no âmbito estadual: o caso do Rio de Janeiro* [online]. Rio de Janeiro: Editora FIOCRUZ; 2010. p. 215-240.
15. Olivencia-Delgado MN, Jusino-Álamo JF, De Miranda-Sánchez E, Quiñones-Rodríguez JI. From cadaveric dissection to the operating room: a unilateral double intercostobrachial nerve and the implications in axillary lymph node dissection. *Cureus.* 2023;15(3):e36647. <https://doi.org/10.7759/cureus.36647>

16. Chang PJ, Asher A, Smith SR. A Targeted Approach to Post-Mastectomy Pain and Persistent Pain following Breast Cancer Treatment. *Cancers (Basel)*. 2021 Oct 16;13(20):5191. <https://doi.org/10.3390/cancers13205191>.
17. Andersen KG, Duriaud HM, Kehlet H, Aasvang EK. The Relationship between sensory loss and persistent pain 1 year after breast cancer surgery. *J Pain*. 2017;18(9):1129-38. <https://doi.org/10.1016/j.jpain.2017.05.002>
18. Lucena RN, Bona AF, Macedo FO, Bergmann A, Aguiar SS, Lou MBA. Perfil das Pacientes Avaliadas no Ambulatório de Fisioterapia em um Hospital de Referência em Oncologia após um Ano de Cirurgia para o Câncer de Mama. In: VI Congresso Brasileiro de Fisioterapia em Oncologia 2019. *Rev Bras Cancerol*. 2019;65.1(Suplemento 2):123.
19. Siqueira TC, Frágoas SP, Pelegrini A, de Oliveira AR, da Luz CM. Factors associated with upper limb dysfunction in breast cancer survivors. *Support Care Cancer*. 2021;29(4):1933-40. <https://doi.org/10.1007/s00520-020-05668-7>
20. Wang K, Yee C, Tam S, Drost L, Chan S, Zaki P, et al. Prevalence of pain in patients with breast cancer post-treatment: A systematic review. *Breast*. 2018;42:113-27. <https://doi.org/10.1016/j.breast.2018.08.105>
21. Murugappan A, Khanna A. Interventional Treatment Options for Post-mastectomy Pain. *Curr Oncol Rep*. 2023;25(10):1175-79. <https://doi.org/10.1007/s11912-023-01435-z>
22. Manfuku M, Nishigami T, Mibu A, Yamashita H, Imai R, Kanamori H, et al. Predictors of persistent post-surgical pain intensity and interference at 1 year after breast cancer surgery: assessing central sensitization, central sensitivity symptoms, and psychological factors. *Breast Cancer*. 2023;30(2):271-81. <https://doi.org/10.1007/s12282-022-01420-7>
23. Wang L, Cohen JC, Devasenapathy N, Hong BY, Kheyson S, Lu D, et al. Prevalence and intensity of persistent post-surgical pain following breast cancer surgery: a systematic review and meta-analysis of observational studies. *Br J Anaesth*. 2020;125(3):346-57. <https://doi.org/10.1016/j.bja.2020.04.088>
24. Klein I, Friger M, David MB, Shahar D. Risk factors for long-term arm morbidities following breast cancer treatments: A systematic review. *Oncotarget*. 2023;14:921-42. <https://doi.org/10.18632/oncotarget.28539>
25. Wang L, Chen H, Li Y, Wang H, Liu N, Yu M, et al. Body mass index increases the risk of breast cancer-related lymphedema at 6-18 months after surgery: a retrospective study. *Support Care Cancer*. 2023;31(5):278. <https://doi.org/10.1007/s00520-023-07721-7>
26. Hara Y, Otsubo R, Shinohara S, Morita M, Kuba S, Matsumoto M, et al. Lymphedema After Axillary Lymph Node Dissection in Breast Cancer: Prevalence and Risk Factors-A Single-Center Retrospective Study. *Lymphat Res Biol*. 2022;20(6):600-6. <https://doi.org/10.1089/lrb.2021.0033>
27. Furlan C, Matheus CN, Jales RM, Derchain SFM, Bennini JR Jr, Sarian LO. Longitudinal, Long-Term Comparison of Single- versus Multipoint Upper Limb Circumference Periodical Measurements as a Tool to Predict Persistent Lymphedema in Women Treated Surgically for Breast Cancer: An Optimized Strategy to Early Diagnose Lymphedema and Avoid Permanent Sequelae in Breast Cancer Survivors. *Ann Surg Oncol*. 2021;28(13):8665-76. <https://doi.org/10.1245/s10434-021-10290-w>
28. Konishi T, Tanabe M, Michihata N, Matsui H, Nishioka K, Fushimi K, et al. Risk factors for arm lymphedema following breast cancer surgery: a Japanese nationwide database study of 84,022 patients. *Breast Cancer*. 2023;30(1):36-45. <https://doi.org/10.1007/s12282-022-01395-5>
29. Paramanandam VS, Dylke E, Clark GM, Daptardar AA, Kulkarni AM, Nair NS, et al. Prophylactic Use of Compression Sleeves Reduces the Incidence of Arm Swelling in Women at High Risk of Breast Cancer-Related Lymphedema: A Randomized Controlled Trial. *J Clin Oncol*. 2022;40(18):2004-12. <https://doi.org/10.1200/JCO.21.02567>
30. Kim JS, Kim JH, Chang JH, Kim DW, Shin KH. Prediction of breast cancer-related lymphedema risk after postoperative radiotherapy via multivariable logistic regression analysis. *Front Oncol*. 2022;12:1026043. <https://doi.org/10.3389/fonc.2022.1026043>
31. Fabro EAN, Bergmann A, do Amaral E Silva B, Padula Ribeiro AC, de Souza Abrahão K, da Costa Leite Ferreira MG, et al. Post-mastectomy pain syndrome: incidence and risks. *Breast*. 2012;21(3):321-5. <https://doi.org/10.1016/j.breast.2012.01.019>
32. Abass MO, Gismalla MDA, Alsheikh AA, Elhassan MMA. Axillary Lymph Node Dissection for Breast Cancer: Efficacy and Complication in Developing Countries. *J Glob Oncol*. 2018;4:1-8. <https://doi.org/10.1200/JGO.18.00080>
33. Marchito L de O, Fabro EAN, Macedo FO, Costa RM, Lou MB de A. Prevenção e Cuidado do Linfedema após Câncer de Mama: Entendimento e Adesão às Orientações Fisioterapêuticas. *Rev Bras Cancerol*. 2019;65(1):e-03273. <https://doi.org/10.32635/2176-9745.RBC.2019v65n1.273>
34. Lee YL, Islam T, Danaee M, Taib NA; MyBCC study group. A longitudinal study of physical activity among Malaysian breast cancer survivors. *PLoS One*. 2022;17(11):e0277982. <https://doi.org/10.1371/journal.pone.0277982>
35. De Groef A, Geraerts I, Demeyer H, Van der Gucht E, Dams L, de Kinkelder C, et al. Physical activity levels after treatment for breast cancer: Two-year follow-up. *Breast*. 2018;40:23-8. <https://doi.org/10.1016/j.breast.2018.04.009>
36. Singleton AC, Raeside R, Hyun KK, Partridge SR, Di Tanna GL, Hafiz N, et al. Electronic Health Interventions for Patients With Breast Cancer: Systematic Review and Meta-Analyses. *J Clin Oncol*. 2022;40(20):2257-70. <https://doi.org/10.1200/JCO.21.01171>
37. Nápoles AM, Santoyo-Olsson J, Chacón L, Stewart AL, Dixit N, Ortiz C. Feasibility of a mobile phone app and telephone coaching survivorship care planning program among Spanish-speaking breast cancer survivors. *JMIR Cancer*. 2019;5(2):e13543. <https://doi.org/10.2196/13543>
38. Vladymyrov OA, Semykopna TV, Vakulenko DV, Syvak OV, Budnyk MM. Telerehabilitation Guidelines for Patients with Breast Cancer. *Int J Telerehab*. 2024;27. <https://doi.org/10.5195/ijt.2024.6642>
39. Delgoshaei B, Mobinizadeh M, Mojdekar R, Afzal E, Arabloo J, Mohamadi E. Telemedicine: A systematic review of economic evaluations. *Med J Islam Repub Iran*. 2017;31:113. <https://doi.org/10.14196/mjiri.31.113>
40. Rinaldi G, Hijazi A, Haghparsat-Bidgoli H. Cost and cost-effectiveness of mHealth interventions for the prevention and control of type 2 diabetes mellitus: A systematic review. *Diabetes Res Clin Pract*. 2020;162:108084. <https://doi.org/10.1016/j.diabres.2020.108084>



A prospective evaluation of breast satisfaction and expectation in preoperative immediate breast reconstruction patients**

Flávia Kuroda^{1,*} , Cícero de Andrade Urban¹ , Maíra Teixeira Dória¹ , Íris Rabinovich¹ ,
Mateus Ricardo Lourenço² , Bernardo Sobreiro³ , Marcelo Loureiro³ 

ABSTRACT

Introduction: Breast reconstruction has an important positive contribution to the quality of life of breast cancer patients. However, a large proportion of breast cancer survivors have unmet expectations surrounding reconstruction. This study aimed to delineate factors affecting preoperative native breast satisfaction and expectations with surgery in immediate breast reconstruction (IBR) patients. **Methods:** This is a prospective cross-sectional trial with breast cancer patients undergoing oncology surgery following breast reconstruction enrolled from 2019 to 2021 at the Hospital Nossa Senhora das Graças in Curitiba, Brazil. Two groups were studied: patients who underwent mastectomy following IBR with implant; and those who underwent breast conservative therapy (BCT) following oncoplastic surgery (OP). All patients completed a patient-reported outcome, the BREAST-Q Breast Reconstruction Expectations Module, prior to surgery. **Results:** Seventy-nine patients with breast cancer were included: 49 OP and 30 mastectomy following IBR. The mastectomy with IBR implants group had a better satisfaction with their native breast than the OP group ($p=0.001$). Women in the OP group had higher expectations for breast appearance when clothed than the mastectomy with IBR implant group ($p=0.030$). Patients aged 50 years and older with a university education or higher level expected that their breast appearance would match almost the same after ten years ($p=0.001$). **Conclusions:** Our results highlight the importance of establishing realistic expectations prior to surgery. Understanding which factors affect patients' satisfaction with native breasts and their expectation toward surgery in the preoperative set could improve preoperative counseling and management of patients' expectations regarding breast reconstruction.

KEYWORDS: expectation; breast reconstruction; satisfaction; quality of life; breast cancer.

INTRODUCTION

Breast cancer care involves highly complex procedures such as surgery in conjunction with oncoplastic techniques and breast reconstruction^{1,2}. Over the past 20 years, there have been many innovations and advancements that elevate the quality of breast reconstruction following a mastectomy or breast conservative surgery. Several methods and surgical techniques were developed such as tissue expanders; shaped, integrated valve; textured saline or silicone gel implants that have undergone significant improvements; a novel and innovative oncoplastic approach described based upon an oncoplastic algorithm; fluorescent laser angiography; acellular dermal matrices; and current techniques

for fat grafting that have revolutionized breast reconstruction. These advancements focus on improving surgical and aesthetic outcomes as well as reducing adverse events³.

There is general agreement that breast reconstruction makes a significant positive contribution to the quality of life of many women who have undergone mastectomy for breast cancer⁴⁻⁶. Patients' satisfaction is one of the most important endpoints whose overriding goal is to meet their expectations and improve their quality of life. However, a large proportion of breast cancer survivors have unmet expectations surrounding reconstruction after mastectomy, particularly in relation to appearance. Approximately 42% of women who underwent breast

¹Hospital Nossa Senhora das Graças – Curitiba (PR), Brazil.

²Faculdade Evangélica Mackenzie – Curitiba (PR), Brazil.

³Universidade Positivo, Post-Graduation Program in Biotechnology – Curitiba (PR), Brazil.

*Corresponding author: flaviakuroda@hotmail.com

Conflict of interests: nothing to declare. Funding: none.

Received on: 01/16/2025 – Accepted on: 03/21/2025.

**The content of this manuscript was presented in part at the XXIII Brazil Breast Congress in April 2021 and published at Mastology, 2021;31(Suppl 1):92.

reconstruction after mastectomy reported their reconstruction to be worse than expected⁷.

There is little consensus about what impact specific factors have on women's satisfaction with the breast reconstruction process and outcomes. Concerning expectation and satisfaction issues, several instrumentals were validated but most of them are general questionnaires that do not specify body and psychological changes experienced by breast cancer patients^{8,9} and compare different surgical approaches. Among the available patient-reported outcome measurement tools, the BREAST-Q has established itself as the gold standard, being most frequently used in the literature¹⁰⁻¹². The BREAST-Q questionnaire was developed especially for breast cancer patients undergoing breast surgery. Independent modules are available for different surgical interventions (e.g., mastectomy, breast reconstruction, or conservative surgery). Each module consists of a preoperative and a postoperative questionnaire¹³. In 2012, a specific preoperative expectation for breast reconstruction module was added to the BREAST-Q set of questionnaires that cover a range of breast surgical procedures. The expectation module covers a thorough range of questions about how the patient expects to feel in the first week, first year, and ten years after breast reconstruction surgery⁴.

A preoperative assessment of quality of life, satisfaction, and expectation can aid the surgeon in an accurate clinical evaluation and may allow for early identification of patients with a higher risk of regret^{14,15}. Furthermore, these assessments improve patient education, shared medical decision-making, patient perception of outcomes¹⁶, and provide a point of reference for assessing change after a procedure¹⁵. Besides, it is an important predictor of health outcomes and health-related quality of life^{16,17}. Unrecognized or unfulfilled expectations have been shown to correlate with patients' low satisfaction and poor overall outcomes in any type of surgery¹⁷⁻²⁰. Despite its importance, few studies to date have focused on measuring expectations and satisfaction prior to oncological breast surgery using systematically the validated BREAST-Q²¹. A systematic review of literature did not find consistent evidence to support a link between patients' expectations and degrees of satisfaction with breast reconstruction outcomes⁴. A recent study that evaluated patients' expectations using the preoperative BREAST-Q expectation score was a retrospective chart review that included mainly delayed reconstruction¹⁷.

The present study aimed to delineate factors affecting preoperative native breast satisfaction and expectations toward surgery using the BREAST-Q in patients before oncological breast surgery following IBR.

METHODS

This is a prospective cross-sectional trial with breast cancer patients undergoing oncology surgery (mastectomy or breast conservative therapy) following breast reconstruction or oncoplastic

surgery enrolled from November 2019 to October 2021 at the Hospital Nossa Senhora das Graças, Breast Unit, in Curitiba, Brazil. All patients had *in situ* or invasive carcinoma diagnosed by core biopsy or vacuum-assisted biopsy⁹. We excluded patients who refused participation in the study, who would undergo prophylactic mastectomy or preoperative radiotherapy, and those who had local recurrence or metastasis at the time of analysis⁹.

Two independent groups of patients undergoing oncology surgery were studied. The first included patients who underwent mastectomy following IBR with definitive anatomical form-stable implant. Here, contralateral symmetrization was performed using different techniques according to the necessity in each individualized case and the possibility of obtaining better symmetry with the reconstructed breast: reduction mammoplasty, mastopexy, augmentation mammoplasty, or mastopexy associated with implant⁹. The second group underwent breast conservative therapy (BCT) following level 2 oncoplastic techniques (bilateral surgeries with mammoplasty techniques).

This study was approved by the Internal Review Board of Positivo University, Curitiba, Brazil, on September 19, 2019.

All patients were invited to complete the patient-reported outcome BREAST-Q Expectations Module and Preoperative Breast Reconstruction or Preoperative Reduction/Mastopexy Module already translated into Portuguese. They signed informed consent and answered the questionnaire in paper format prior to the surgical procedure.

The BREAST-Q Preoperative Breast Reconstruction Module comprises two domains: satisfaction (i.e., satisfaction with breasts) and quality of life (psychosocial, physical, and sexual well-being), consisting of five scales. The score from each scale is transferred into a 100-point scale. Thus, BREAST-Q question values were transformed and scored using the QScore, a statistical program developed specifically for the BREAST-Q that provides a total scale score, ranging from 0 to 100, in which a higher score suggests a better quality of life or satisfaction⁸⁻¹⁰.

The BREAST-Q Preoperative Expectation short-form module is composed of five scales and assesses:

1. Pain;
2. Appearance when clothed after one year;
3. Appearance of breast symmetry after one year;
4. Sensation of breast after one year; and
5. Appearance of breast symmetry after ten years.

Response options for all scales are on a 3-point Likert-type scale, where 1 represents unlikely, 2 likely, and 3 very likely¹⁷.

Item responses for each section of the modules are summed and transformed to give a score for each scale (0–100), using a standardized conversion template¹⁷. Statistical analysis was performed using the Statistical Package for Social Sciences (SPSS) program. For quantitative variables, expressed with mean \pm standard deviation, the Mann-Whitney U test was applied. For

qualitative variables, expressed with numbers and percentages. Fisher's exact test was used. Sociodemographic and clinical characteristics were compared between groups. A $p < 0.050$ was considered statistically significant.

RESULTS

Overall, 79 breast cancer patients completed the preoperative BREAST-Q questionnaire. Patients were divided into two groups: BCT following oncoplastic surgery ($n=49$) and mastectomy following IBR with implant ($n=30$). Table 1 summarizes the sociodemographic characteristics of the cohort. The mean age was 52.6

standard deviation ± 12.3 years. Most patients were considered overweight or obese (64.5%) and 16 women had previously been submitted to breast aesthetic surgery (20.3%).

Table 2 shows BREAST-Q expectation and satisfaction rates for both groups. The mastectomy with IBR implants group had a better satisfaction with their native breasts than BCT oncoplastic group ($p=0.001$). There was no statistically significant difference between groups regarding the other parameters.

When we compared BREAST-Q reconstruction expectations rate, women in BCT following oncoplastic group had higher expectations for breast appearance when clothed than the mastectomy with IBR implant group (93.4 ± 16.3 vs. 82.9 ± 26.5 ; $p=0.030$).

Table 1. Sociodemographic characteristics of study cohort.

Characteristic	BCT+Oncoplastic n (%)	Mastectomy with IBR implants n (%)	p-value
Age, mean±SD (year)	52.3±12.8	53±11.6	0.82
BMI, mean±SD	27.4±4.7	25.5±4.1	0.07
Weight characteristics			
Normal	14 (28.6)	14 (46.7)	0.16
Overweight	26 (53.1)	14 (46.7)	
Obese	9 (18.4)	2 (6.7)	
Menopausal status			
Postmenopausal	25 (51.0)	17 (56.7)	0.65
Premenopausal	24 (49.0)	13 (43.3)	
HRT			
Yes	16 (32.7)	9 (30.0)	1.00
No	33 (67.3)	21 (70.0)	
Education level			
Unfinished primary school	5 (10.2)	1 (3.3)	0.81
Full primary school	1 (2.0)	2 (6.7)	
High school	9 (18.4)	7 (23.3)	
College degree	18 (36.7)	6 (20.0)	
Specialization, postgraduate degree	16 (32.7)	14 (46.7)	
Family history			
Yes	24 (49.0)	13 (43.3)	0.65
No	25 (51.0)	17 (56.7)	
Previous aesthetic breast surgery			
Yes	7 (14.3)	9 (30.0)	0.15
No	42 (85.7)	21 (70.0)	
Neoadjuvant chemotherapy			
Yes	17 (34.7)	9 (30.0)	0.81
No	32 (65.3)	21 (70.0)	
Smoking			
Yes	3 (6.1)	5 (16.7)	0.25
No	46 (93.9)	25 (83.3)	

BCT: breast conservative therapy; IBR: immediate breast reconstruction; SD: standard deviation; BMI: body mass index; HRT: hormonal reposiotion therapy.

Table 2. BREAST-Q satisfaction and expectation rates between the two groups.

BREAST-Q satisfaction			
	BCT+Oncoplastics (n=49)	Mastectomy with IBR implants (n=30)	p-value
	Mean±SD	Mean±SD	
Psychosocial well-being	55.5±16.7	71.9±23.0	0.920
Sexual well-being	61.3±23.0	61±21.6	0.950
Physical well-being	69.1±16.4	68.2±22.4	0.850
Satisfaction with breast	55.5±16.7	71.9±23	0.001
BREAST-Q expectations			
	BCT + Oncoplastics (n=49)	Mastectomy with IBR implants (n=30)	p-value
	Mean±SD	Mean±SD	
Expectations for pain	63.2±18.9	56.7±23.9	0.190
Expectations for breast appearance when clothed	93.4±16.3	82.9±6.5	0.030
	n (%)	n (%)	p-value
Expectation for breast appearance when unclothed after one year			
Will look very different	1 (2.0)	0 (0.0)	0.623
Will look similar	35 (71.4)	24 (80.0)	
Will look exactly the same	6 (12.2)	4 (13.3)	
Don't know	7 (14.3)	2 (6.7)	
Expectations for breast sensation after one year			
Almost no sensation	3 (6.1)	6 (20.0)	0.001
Will have some sensation	10 (20.4)	13 (43.3)	
Will have normal sensation	25 (51.0)	3 (10.0)	
Don't know	21 (42.9)	8 (26.7)	
Expectation for breast appearance after ten years			
Will not match	9 (18.4)	6 (20.0)	0.721
Will match almost	21 (42.9)	11 (36.7)	
Will match exactly	3 (6.1)	4 (13.3)	
Don't know	16 (32.7)	9 (30.0)	

BCT: breast conservative therapy; IBR: immediate breast reconstruction (implant based); SD: standard deviation.
 Bold indicates statistically significant p-values.

Most patients in both groups expected that breast appearance (symmetry) when unclothed would look similar after one year (71.4% for BCT and 80.0% for mastectomy group) and after ten years would match almost the same as it did right after the reconstruction (42.9% for BCT and 36.7% for mastectomy group). In the BCT with oncoplastic group, 51.0% of patients expected that the breast would have normal sensation after one year, whereas 43.3% of women in the mastectomy with IBR group expected to have some sensation ($p=0.001$).

Table 3 shows logistic regression analysis and results. Previous aesthetic breast surgery and neoadjuvant chemotherapy were significant predictors of preoperative physical well-being. Patients 50 years or older and with a university degree or higher level of education expected that their breast appearance would match almost the same after ten years ($p=0.001$) (Table 4).

DISCUSSION

Patients' satisfaction with their breasts is an important metric for the evaluation of outcomes in breast surgery¹⁵. Many factors affect aesthetics and satisfaction with each native breast; it is difficult to capture in existing assessments. In our study, the mastectomy with IBR implant group had better satisfaction with their native breast than the BCT oncoplastic group ($p=0.001$). Despite all these variables and nonspecific factors, it is essential to have baseline scores representative of patients' self-perception (15) before treatment in order to assess whether quality of life will change postoperatively.

Patients in the oncoplastic group had worse preoperative psychosocial well-being (55.5±16.3) than the breast reconstruction group (71.9±23.0), and in both groups, we found low physical well-being scores. It is important to consider that preoperative

Table 3. BREAST-Q satisfaction and reconstruction expectations according to different factors.

	Psychosocial well-being	Sexual well-being	Physical well-being	Satisfaction with breast	Expectations for pain	Expectations for breast appearance when clothed
Age (years)						
<39	73.8±18.3	67.6±17.2	70.1±23.2	65.0±18.9	70.0±14.7	87.5±21.6
40–49	66.7±19.9	55.2±22.4	64.1±17.3	55.7±19.2	64.4±19.7	91.6±15.5
50–59	66.9±19.4	64.4±19.7	69.0±17.7	66.0±20.6	63.4±18.0	84.4±29.6
>60	76.0±20.2	61.3±27.6	72.6±19.1	61.6±23.3	49.5±24.7	93.3±15.6
p-value	0.31	0.40	0.51	0.37	0.26	0.53
Educational level						
High school or less	68.7±23.4	57.4±27.8	69.3±20.1	60.6±21.8	54.1±20.6	92.5±16.0
University or more	71.1±18.0	62.9±19.7	68.6±18.3	62.1±20.5	63.4±21.0	88.2±23.0
p-value	0.62	0.33	0.89	0.78	0.90	0.42
Previous aesthetic breast surgery						
Yes	73.7±18.6	70.9±18.1	77.2±15.1	68.4±22.8	53.4±27.7	82.9±29.1
No	69.5±20.1	58.8±22.9	66.6±19.1	59.9±20.1	62.8±18.7	91.1±18.6
p-value	0.45	0.06	0.044*	0.14	0.11	0.17
Weight characteristic						
Normal	75.5±20.5	65.9±22	74.3±18.0	65.7±20.4	62.9±19.7	90.0±18.4
Overweight	65.9±18.6	57.9±24.6	64.9±17.7	59.7±21.9	59.7±21.9	90.7±21.8
Obese	73.6±19.4	60.1±12.1	69.6±21.8	60.6±12.8	58.5±23.5	84.2±25.8
p-value	0.12	0.37	0.12	0.43	0.78	0.66
Neoadjuvant chemotherapy						
Yes	65.3±19.9	55.6±24.9	62.7±14.7	60.5±21.2	65.0±18.4	92.4±15.1
No	72.9±19.3	64.1±20.6	71.7±19.8	62.3±20.7	58.5±22.2	88.0±23.6
p-value	0.11	0.12	0.04*	0.67	0.21	0.40

*Statistically significant ($p < 0.050$).

patients are not “normal”, as they have undergone the physical and psychological trauma associated with being diagnosed with breast cancer²² — a unique entity and a life-changing moment for each patient. The low physical well-being score may be explained by pain secondary to the tumor itself or pain after biopsy before cancer resection²². A study by Roth et al.²³ showed that women who reported higher preoperative levels of distress and anxiety were significantly less satisfied with the outcomes of breast reconstruction^{23,24}. Clearly, many clinical and non-clinical factors influence a woman's satisfaction with psychosocial and physical breast reconstruction outcomes, making a single measurement of satisfaction challenging⁴. Differently, Builes Ramírez et al. identified no anthropometric and clinical variables related to satisfaction and quality of life in breast cancer women before their surgical procedure²⁵. In our study, we found that variations in expectations such as previous aesthetic breast surgery and neoadjuvant chemotherapy were significant predictors of preoperative physical well-being.

The assessment and management of patients' expectations may improve their perception of outcomes¹⁶. When we compare

the two different types of surgery, in the BCT with oncoplastic group, 51.0% of patients expected that the breast would have normal sensation after one year, whereas 43.3% of women in mastectomy with IBR group expected to have some sensation ($p = 0.001$). A review by Sisco et al.²⁶ reported that sensory outcomes in nipple-sparing mastectomy varied, with normal sensation self-reported in the range 10.0–43.0%^{26,27}. However, it has now become clear that nipple sensation is largely or completely lost in most cases. A Swedish prospective study that quantitatively examined tactile, thermal, and nociceptive cutaneous sensitivity before and after nipple-sparing mastectomy found total loss of touch sensation in the nipple in 62.0% of patients, while touch sensation was impaired in the remaining 38.0%^{27,28}. These findings highlight the importance of managing patients' expectations about breast and nipple sensations after mastectomy to reduce the risk of dissatisfaction with the surgery.

Interestingly, we identified that most women in both groups expected that breast appearance (symmetry) when unclothed would look similar after one year (71.4% for BCT and 80.0% for

Table 4. Analysis of BREAST-Q reconstruction expectation for breast appearance after ten years according to different factors.

Expectation for breast appearance after ten years					
	Will not match	Will match almost	Will match exactly	Don't know	p-value
Age (years)					
<39	4	6	0	1	0.03*
40–49	5	7	0	12	
50–59	5	9	2	6	
>60	1	10	5	6	
Educational level					
High school or less	2	9	1	13	0.04*
University or more	13	23	6	12	
Previous aesthetic breast surgery					
Yes	6	6	2	2	0.09
No	9	26	5	23	
Weight characteristic					
Normal	8	13	2	5	0.18
Overweight	6	17	3	14	
Obese	1	2	2	6	
Neoadjuvant chemotherapy					
Yes	6	11	0	9	0.27
No	9	21	7	16	

*Statistically significant ($p < 0.050$).

mastectomy group) and after ten years would match almost the same as it did right after reconstruction (42.9% for BCT and 36.7% for mastectomy group). Overall, the aesthetic outcomes decline over time, especially if chemotherapy and radiotherapy are required. Furthermore, breast cancer patients using adjuvant endocrine therapy can vary their weight resulting in asymmetry, impacting patient-reported outcomes. In breast-conserving therapy, a prospective study by Hennigs et al.²⁹ showed that the change in the aesthetic outcome is still measurable over four years after the surgical procedure with a subjective evaluation^{29,30}. In breast reconstruction with implant, several authors have described a trend of deterioration over time, with a decline in aesthetic outcomes, an increase in capsular contracture, and an overall decrease in patient satisfaction^{10,31,32}. Seth and Cordeiro contradict these results demonstrating that prosthetic breast reconstruction outcomes do not deteriorate over time. This stability is apparent in both long-term surgeon and patient report outcomes data measured in the same patients¹⁰. Despite the differences in the literature, we delineated factors such as patients aged 50 years and older with university education or higher who expect their breast appearance to match almost the same after ten years ($p = 0.001$). These findings emphasize the importance of managing patient expectations about breast and nipple sensation after mastectomy and aesthetic outcomes over time to reduce the risk of dissatisfaction with the surgery.

It is important to consider that most data were collected during the coronavirus disease (COVID-19) that was first reported in Wuhan (China), in December 2019. The COVID-19 pandemic became one of the main international concerns regarding its impact on mental health³³. A study that included 3,000 Brazilian population from 25 states showed that almost half of participants expressed symptoms of depression (46.4%), anxiety (30.7%), and stress (42.2%) in this period³³. Mental illness during the pandemic associated with the diagnosis of breast cancer may have adversely affected the satisfaction and quality of life scores found in our study.

A strength of this work is that it is the first prospective study that provides a useful perspective on the patients' feelings prior to breast cancer surgery, using the objective, validated, and reliable BREAST-Q questionnaire. The recruitment of this population included all breast cancer patients who underwent oncological surgery with IBR or oncoplastic surgery. We excluded those who underwent a prophylactic mastectomy and delayed breast reconstruction to get a homogenous cohort. This comparison enables surgeons to adopt an individualized approach according to the technique to be employed.

In contrast, this study also has several limitations. Our population was restricted to a single center, limiting the generalizability of data. As a cross-sectional study, there is an important element of selection bias to consider. We only included patients

who agreed to participate in the study; they probably had a better quality of life and satisfaction score than those who refused to take part.

Further studies are needed to evaluate the effect of preoperative patient expectations on patient-reported outcomes following breast reconstruction to determine whether preoperative expectations can be modified to produce long-term satisfaction after surgery. It is well documented that patients' satisfaction with their breasts correlates more strongly with their satisfaction with the information they received prior to surgery and with their plastic surgeon^{24,34,35}. Failure to recognize and understand what patients expect from their surgical procedure often leads to dissatisfaction and poor overall outcome for them²⁴.

CONCLUSIONS

This study's results highlight the need to improve education and informed decision-making about breast reconstruction. Patients demonstrated high expectations for breast appearance after reconstruction and expected it not to change over time.

Multiple factors influence preoperative breast satisfaction and expectation prior to surgery. Understanding which factors affect patients' satisfaction with native breasts and their expectations with the surgery in the preoperative set could improve preoperative counseling and the management of patients' expectations of subsequent breast reconstruction and reduce the risk of dissatisfaction with the surgery.

AUTHORS' CONTRIBUTIONS

FK: Conceptualization, Investigation, Methodology, Project administration, Resources, Validation, Visualization, Writing – original draft, Writing – review and editing, Manuscript review. CAU: Conceptualization, Methodology, Project administration, Resources, Supervision, Manuscript review. MTD: Investigation, Validation, Visualization, Writing – original draft, Writing – review and editing. IR: Investigation, Validation, Visualization, Writing – original draft, Writing – review and editing. MRL: Data curation, Formal analysis, Software. BS: Data curation, Formal analysis, Software. ML: Conceptualization, Methodology, Project administration, Resources, Supervision, Manuscript review.

REFERENCES

1. Fisher B, Anderson S, Bryant J, Margolese RG, Deutsch M, Fisher ER, et al. Twenty-year follow-up of a randomized trial comparing total mastectomy, lumpectomy, and lumpectomy plus irradiation for the treatment of invasive breast cancer. *N Engl J Med*. 2002;347(16):1233–41. <https://doi.org/10.1056/NEJMoa022152>
2. Freitas-Júnior R, Gagliato DM, Moura Filho JWC, Gouveia PA, Rahal RMS, Paulinelli RR, et al. Trends in breast cancer surgery at Brazil's public health system. *J Surg Oncol*. 2017;115(5):544–9. <https://doi.org/10.1002/jso.24572>
3. Nahabedian M. Innovations and advancements in breast reconstruction. *Gland Surg*. 2015;4(2):91–2. <https://doi.org/10.3978/j.issn.2227-684X.2015.04.02>
4. Flitcroft K, Brennan M, Spillane A. Women's expectations of breast reconstruction following mastectomy for breast cancer: a systematic review. *Support Care Cancer*. 2017;25(8):2631–61. <https://doi.org/10.1007/s00520-017-3712-x>
5. Cordeiro PG. Breast reconstruction after surgery for breast cancer. *N Engl J Med*. 2008;359(15):1590–601. <https://doi.org/10.1056/NEJMct0802899>
6. Rowland J, Desmond K, Meyerowitz B, Belin TR, Wyatt GE, Ganz PA. Role of breast reconstructive surgery in physical and emotional outcomes among breast cancer survivors. *J Natl Cancer Inst*. 2000;92(17):1422–9. <https://doi.org/10.1093/jnci/92.17.1422>
7. Steffen LE, Johnson A, Levine BJ, Mayer DK, Avis NE. Met and unmet expectations for breast reconstruction in early posttreatment breast cancer survivors. *Plast Surg Nurs*. 2017;37(4):146–53. <https://doi.org/10.1097/PSN.0000000000000205>
8. Cardoso MJ, Santos AC, Cardoso J, Barros H, Oliveira MC. Choosing observers for evaluation of aesthetic results in breast cancer conservative treatment. *Int J Radiat Oncol Biol Phys*. 2005;61(3):879–81. <https://doi.org/10.1016/j.ijrobp.2004.06.257>
9. Kuroda F, Urban C, Zucca-Matthes G, Oliveira VM, Arana GH, Iera M, et al. Evaluation of aesthetic and quality-of-life results after immediate breast reconstruction with definitive form-stable anatomical implants. *Plast Reconstr Surg*. 2016;137(2):278e–86e. <https://doi.org/10.1097/01.prs.0000475746.17968.f4>
10. Seth AK, Cordeiro PG. Stability of long-term outcomes in implant-based breast reconstruction: an evaluation of 12-year surgeon- and patient-reported outcomes in 3489 nonirradiated and irradiated implants. *Plast Reconstr Surg*. 2020;146(3):474–84. <https://doi.org/10.1097/PRS.00000000000007117>
11. Pusic AL, Klassen AF, Scott AM, Klok JA, Cordeiro PG, Cano SJ. Development of a new patient-reported outcome measure for breast surgery: the BREAST-Q. *Plast Reconstr Surg*. 2009;124(2):345–53. <https://doi.org/10.1097/PRS.0b013e3181aee807>
12. Pusic AL, Matros E, Fine N, Buchel E, Gordillo GM, Hamill JB, et al. Patient-reported outcomes 1 year after immediate breast reconstruction: results of the Mastectomy Reconstruction Outcomes Consortium study. *J Clin Oncol*. 2017;35(22):2499–506. <https://doi.org/10.1200/JCO.2016.69.9561>
13. Stolpner I, Heil J, Riedel F, Wallwiener M, Schäfer B, Feißt M, et al. Long-term patient satisfaction and quality of life after breast-conserving therapy: a prospective study using the BREAST-Q. *Ann Surg Oncol*. 2021;28(13):8742–51. <https://doi.org/10.1245/s10434-021-10377-4>

14. Zhong T, Hu J, Bagher S, O'Neill AC, Beber B, Hofer SOP, et al. Decision regret following breast reconstruction: the role of self-efficacy and satisfaction with information in the preoperative period. *Plast Reconstr Surg*. 2013;132(5):724e-34e. <https://doi.org/10.1097/PRS.0b013e3182a3bf5d>
15. Klifto KM, Aravind P, Major M, Payne RM, Shen W, Rosson GD, et al. Differences between Breast Cancer Reconstruction and Institutionally Established Normative Data Using the BREAST-Q Reconstruction Module: a comparative study. *Plast Reconstr Surg*. 2020;145(6):1371-9. <https://doi.org/10.1097/PRS.00000000000006810>
16. Pusic AL, Klassen AF, Snell L, Cano SJ, McCarthy C, Scott A, et al. Measuring and managing patient expectations for breast reconstruction: Impact on quality of life and patient satisfaction. *Expert Rev Pharmacoecon Outcomes Res*. 2012;12(2):149-58. <https://doi.org/10.1586/erp.11.10>
17. Morzycki A, Corkum J, Joukhadar N, Samargandi O, Williams JG, Frank SG. The impact of delaying breast reconstruction on patient expectations and health-related quality of life: an analysis using the BREAST-Q. *Plast Surg (Oakv)*. 2020;28(1):46-56. <https://doi.org/10.1177/2292550319880924>
18. Noble PC, Conditt MA, Cook KF, Mathis KB. The John Insall Award: patient expectations affect satisfaction with total knee arthroplasty. *Clin Orthop Relat Res*. 2006;452:35-43. <https://doi.org/10.1097/01.blo.0000238825.63648.1e>
19. Frank Henn 3rd R, Kang L, Tashjian RZ, Green A. Patients' preoperative expectations predict the outcome of rotator cuff repair. *J Bone Joint Surg Am*. 2007;89(9):1913-9. <https://doi.org/10.2106/JBJS.F.00358>
20. Burton KE, Wright V, Richards J. Patients' expectations in relation to outcome of total hip replacement surgery. *Ann Rheum Dis*. 1979;38(5):471-4. <https://doi.org/10.1136/ard.38.5.471>
21. Snell L, McCarthy C, Klassen A, Cano S, Rubin L, Hurley K, et al. Clarifying the expectations of patients undergoing implant breast reconstruction: a qualitative study. *Plast Reconstr Surg*. 2010;126(6):1825-30. <https://doi.org/10.1097/PRS.0b013e3181f44580>
22. Mundy LR, Homa K, Klassen AF, Pusic AL, Kerrigan CL. Breast cancer and reconstruction: normative data for interpreting the BREAST-Q. *Plast Reconstr Surg*. 2017;139(5):1046e-55e. <https://doi.org/10.1097/PRS.00000000000003241>
23. Roth RS, Lowery JC, Davis J, Wilkins EG. Psychological factors predict patient satisfaction with postmastectomy breast reconstruction. *Plast Reconstr Surg*. 2007;119(7):2008-15. <https://doi.org/10.1097/01.prs.0000260584.09371.99>
24. Yip JM, Watson DI, Tiggemann M, Hsia S, Smallman AE, Dean NR. Determinants of breast reconstruction outcome: how important is volume symmetry? *J Plast Reconstr Aesthet Surg*. 2015;68(5):679-85. <https://doi.org/10.1016/j.bjps.2014.12.038>
25. Builes Ramírez S, Acea Nebril B, García Novoa A, Cereijo C, Bouzón A, Mosquera Oses J. Evaluation of the preoperative perception of quality of life and satisfaction of women with breast cancer using the BREAST-Q™ questionnaire. *Cir Esp (Engl Ed)*. 2020;98(4):212-8. <https://doi.org/10.1016/j.ciresp.2019.10.004>
26. Sisco M, Yao KA. Nipple-sparing mastectomy: a contemporary perspective. *J Surg Oncol*. 2016;113(8):883-90. <https://doi.org/10.1002/jso.24209>
27. Galimberti V, Vicini E, Corso G, Morigi C, Fontana S, Sacchini V, et al. Nipple-sparing and skin-sparing mastectomy: review of aims, oncological safety and contraindications. *Breast*. 2017;34(Suppl 1):S82-4. <https://doi.org/10.1016/j.breast.2017.06.034>
28. Gahm J, Hansson P, Brandberg Y, Wickman M. Breast sensibility after bilateral risk-reducing mastectomy and immediate breast reconstruction: a prospective study. *J Plast Reconstr Aesthet Surg*. 2013;66(11):1521-7. <https://doi.org/10.1016/j.bjps.2013.06.054>
29. Hennigs A, Hartmann B, Rauch G, Golatta M, Tabatabai P, Domschke C, et al. Long-term objective esthetic outcome after breast-conserving therapy. *Breast Cancer Res Treat*. 2015;153(2):345-51. <https://doi.org/10.1007/s10549-015-3540-y>
30. Heil J, Riedel F, Golatta M, Hennigs A. Measurement and optimizing cosmetic outcomes for breast excisions/factors influencing aesthetic outcomes of breast conservation surgery. In: Klimberg V, Kovacs T, Rubio I, eds. *oncoplastic breast surgery techniques for the general surgeon*. Cham: Springer International Publishing; 2020. p. 93-106. https://doi.org/10.1007/978-3-030-40196-2_4
31. Nelson JA, Allen Jr RJ, Polanco T, Shamsunder M, Patel AR, McCarthy CM, et al. Long-term patient-reported outcomes following postmastectomy breast reconstruction: an 8-year examination of 3268 patients. *Ann Surg*. 2019;270(3):473-83. <https://doi.org/10.1097/SLA.00000000000003467>
32. Yueh JH, Slavin SA, Adesiyun T, Nyame TT, Gautam S, Morris DJ, et al. Patient satisfaction in postmastectomy breast reconstruction: a comparative evaluation of DIEP, TRAM, latissimus flap, and implant techniques. *Plast Reconstr Surg*. 2010;125(6):1585-95. <https://doi.org/10.1097/PRS.0b013e3181cb6351>
33. Serafim AP, Durães RSS, Rocca CCA, Gonçalves PD, Saffi F, Cappellozza A, et al. Exploratory study on the psychological impact of COVID-19 on the general Brazilian population. *PLoS One*. 2021;16(2):e0245868. <https://doi.org/10.1371/journal.pone.0245868>
34. Ho AL, Klassen AF, Cano S, Scott AM, Pusic AL. Optimizing patient-centered care in breast reconstruction: the importance of preoperative information and patient-physician communication. *Plast Reconstr Surg*. 2013;132(2):212e-20e. <https://doi.org/10.1097/PRS.0b013e31829586fa>
35. Zhong T, McCarthy C, Min S, Zhang J, Beber B, Pusic AL, et al. Patient satisfaction and health-related quality of life after autologous tissue breast reconstruction: a prospective analysis of early postoperative outcomes. *Cancer*. 2012;118(6):1701-9. <https://doi.org/10.1002/cncr.26417>



Trends in mastectomy performance for early breast cancer in a public institution with limited access: a retrospective cohort

Francisco Pimentel Cavalcante^{1,2} , Tallita Moniele Gomes Pinheiro³ , Felipe Pereira Zerwes^{2,4} ,
Eduardo Camargo Millen^{2,5} , Andre Mattar^{2,6} , Marcelo Antonini^{2,7} , Fabrício Palermo Brenelli^{2,8} ,
Guilherme Garcia Novita⁹ , Antônio Luiz Frasson^{2,10} , Ruffo Freitas-Junior¹¹ 

ABSTRACT

Objective: To assess trends in breast surgery, Breast-conserving surgery (BCS) and mastectomy, in an institution with limited access to health resources. **Methods:** A retrospective cohort study was carried out in patients who underwent surgery for non-metastatic breast cancer between 2012 and 2019 at the Hospital Geral de Fortaleza (HGF), an institution that exclusively treats patients from the Brazilian public health system (SUS). The main objective of the study was to evaluate the rates of mastectomy in the period, with or without immediate reconstruction, as well as BCS rates. The χ^2 test, with Bonferroni adjustment, was applied to the relative frequency of the procedures performed to test for statistical significance in the evolution of the frequencies of surgeries over the years. **Results:** A total of 805 patients underwent surgical treatment for non-metastatic breast cancer, with an average of 100 surgeries per year (range 85–118) during the study period. Mastectomy was performed in 552 cases (68.57%), while 253 patients underwent BCS (31.42%). Among the patients who underwent mastectomy, 181 (32.78%) had immediate reconstruction, with the highest proportion using implants (92.26%). No statistical difference was observed between mastectomies with or without reconstruction throughout the period ($p=0.6635$), with a statistically significant difference between BCS ($p=0.04281$) and mastectomies. **Conclusion:** There was no increase in the rates of mastectomies, with and without immediate reconstruction, over the years, but a trend towards an increase in BCS. Further studies are needed to better understand this trend in settings with limited access to health care.

KEYWORDS: breast neoplasms; mastectomy; mastectomy, subcutaneous; mastectomy, segmental; mammoplasty.

INTRODUCTION

Breast cancer is the most common malignant neoplasm in women, with an estimated 2 million cases annually worldwide^{1,2}. In Brazil, the National Cancer Institute (INCA) estimates that there will be more than 73,000 cases of breast cancer in 2023³. The prognosis of the disease, on the other hand, has improved significantly in recent decades, with a significant impact on mortality. The advent of organized screening, significant improvements

in cancer treatment, and a better understanding of biology are responsible for this impact^{4,5}.

Breast cancer surgery has evolved substantially over the years: there has been a gradual replacement of more radical techniques, such as the mastectomy proposed by Halsted, by less invasive approaches⁶. Breast-conserving surgery (BCS), associated with modern multimodal treatment, has similar local recurrence and overall survival rates compared to mastectomy^{7,8}. However,

¹Hospital Geral de Fortaleza – Fortaleza (CE), Brazil.

²Brazilian Breast Association Team – São Paulo (SP), Brazil.

³Santa Casa de Misericórdia de Fortaleza – Fortaleza (CE), Brazil.

⁴Pontifícia Universidade Católica do Rio Grande do Sul – Porto Alegre (RS), Brazil.

⁵Americas Oncologia – Rio de Janeiro (RJ), Brazil.

⁶Hospital da Mulher – São Paulo (SP), Brazil.

⁷Hospital do Servidor Público Estadual – São Paulo (SP), Brazil.

⁸Universidade Estadual de Campinas – Campinas (SP), Brazil.

⁹Oncoclínicas – São Paulo (SP), Brazil.

¹⁰Hospital Israelita Albert Einstein – São Paulo (SP), Brazil.

¹¹Universidade Federal de Goiás, Hospital das Clínicas – Goiânia (GO), Brazil.

*Corresponding author: fpimentelcavalcante@gmail.com

Conflict of interests: nothing to declare. Funding: none.

Received on: 01/23/2025. Accepted on: 03/21/2025.

despite the oncological safety of BCS, a trend towards increasing mastectomy rates, especially bilateral mastectomy, has been observed in developed countries in recent years. The advent of genetic knowledge, the possibility of immediate reconstruction and the false impression of greater safety have been related to the increase in mastectomy rates⁹⁻¹⁴. This trend, however, has not yet been adequately evaluated among patients with limited access to health care systems, where these technologies are not universally available.

The objective of this study was to characterize the trends in surgical procedures in non-metastatic breast cancer, especially mastectomies, with or without immediate breast reconstruction, compared to BCS, between 2012 and 2019, in a single Brazilian institution that provides exclusive treatment to patients of the Unified Health System (SUS).

METHODS

The main objective of this study was to analyze trends in mastectomy and BCS rates, between 2012 and 2019, in patients treated for non-metastatic breast cancer at the Hospital Geral de Fortaleza (HGF), a Brazilian public institution that provides care exclusively to patients covered by SUS. In Brazil, 80% of the population's medical care is provided by SUS. The study was initiated after approval by the institution's Research Ethics Committee (CAAE 29325720.7.0000.5040).

The records of surgical treatment of patients with breast cancer were evaluated, which included: BCS (with or without oncoplastic surgery), total mastectomy without immediate reconstruction, and mastectomy associated with immediate reconstruction, total or with preservation of the nipple-areola complex. The type of reconstruction performed, implants (permanent or temporary expanders) or myocutaneous flaps, and the specific date (year) of each procedure were also evaluated. The decision on each procedure was individually decided by the institution's breast surgeons, as well as the surgical technique used, including oncoplasty and the type of reconstruction. The institution's adjuvant and neoadjuvant treatments followed international guidelines. There was no genetic counseling or testing available during the study period. Patients with benign or indeterminate lesions, as well as other malignant breast lesions, such as melanoma or sarcoma, were excluded from the analysis, as were cases without adequate information on surgical treatment, clinical stage IV at the time of diagnosis or who did not undergo surgery.

Data were tabulated in a spreadsheet compatible with the SPSS-IBM version 20.0 application, used for data analysis. Data were expressed as absolute frequencies and percentages. The χ^2 test, with Bonferroni adjustment, was applied to the relative frequency of the procedures performed to determine the statistical significance in the evolution of frequencies by year. 95%CI with $p < 0.05$ was used to determine the relationship of the variables

with the trends in the rates measured during the study period under a log-linear model (Poisson regression).

RESULTS

After applying the study inclusion criteria, 805 patients undergoing surgical treatment for non-metastatic breast cancer were included for analysis. The average number of surgeries during this period was 100 procedures per year, ranging from 85 cases in 2012 to 118 in 2015 (Figure 1).

The most frequently performed surgical procedure was total mastectomy, in 552 cases (68.57%), while 253 patients underwent BCS (31.42%). In 2012, among the 85 surgeries performed, 44 (51.76%) cases were mastectomies without reconstruction, 19 (22.35%) mastectomies with reconstruction, and 22 (25.88%) BCS. On the other hand, in the last year of the analysis, in 2019, among 106 surgical procedures performed, 41 (38.68%) cases were BCS, 43 (40.57%) mastectomies without reconstruction, and 22 (20.75%) mastectomies with reconstruction. The year with the highest proportion of BCS was 2018 ($n=41$; 40.20%), while 2014 had the highest proportion of mastectomies ($n=70$; 81.40%). Among patients undergoing mastectomy, 181 (32.78%) had immediate reconstruction, where the highest proportion were with isolated implants (92.26%) and 14 cases with flaps (7.73%) (Figure 2).

When analyzing the evolution of mastectomies, no statistically significant increase in this procedure was observed over the period, nor was there a statistically significant difference between mastectomies with or without reconstruction ($p=0.663$). On the other hand, there was a statistically significant difference in conservative surgery over the years ($p=0.042$). Individually evaluating the proportions of each type of surgery during the study period, a statistically significant change was found only for conservative surgery ($p=0.001$; Bonferroni adjustment: $p=0.003$), compared to mastectomy without reconstruction ($p=0.623$; Bonferroni adjustment: $p=0.299$) and mastectomy with reconstruction ($p=0.591$; Bonferroni adjustment: $p=0.663$), as shown in Table 1. Finally, using




Figure 1. Evolution of total surgeries annually between 2012 and 2019.

Poisson regression, a variation in the profile of surgeries over the years was identified (<0.001), but without a significant difference between the types of surgery or immediate reconstruction (Table 2).

DISCUSSION

In this study, no overall increase in mastectomies was observed in an institution that treated patients in the Brazilian public system between 2012 and 2019. Conversely, an upward trend in BCS rates was observed throughout the study period. BCS is the preferred treatment for early breast cancer, replacing radical mastectomy in most cases, following the results of several randomized studies. A study from Denmark, for example, assessed the prevalence of BCS between 1982 and 2002, with

an increase from less than 1% to 25% of conservative surgery, with a significant increase triggering the increase after the publication of the Danish Breast Cancer Cooperative Group study in 1988¹⁵. This trend, however, has recently changed in developed countries. A retrospective American study evaluated the temporal trend of mastectomies in more than 1 million women treated in centers accredited by the American Cancer Society and the American College of Surgeons Commission on Cancer between 1998 and 2011, using the National Cancer Database (NCDB), and it observed a 34% increase in mastectomies, with an odds ratio of 1.34 (95%CI 1.31–1.38)¹⁴. Other studies also identified an increase in bilateral mastectomies in the United States, with stability in conservative surgeries^{12,13,16}.

There are several hypotheses that may explain these differences, including the availability of immediate reconstruction. In Brazil, immediate breast reconstruction after mastectomy was guaranteed by a recent law for the public health system (Law No. 12,802, of 2013)¹⁷. However, in practice, it is possible that the impact has not yet been significant. In our analysis, only 32% of cases underwent immediate reconstruction, which may have affected the results, despite it being available in the institution.




Figure 2. Proportion (%) of surgeries (breast-conserving therapy, in yellow; total mastectomy, in blue; and mastectomy and immediate reconstruction, in red) annually during the period of 2012 to 2019.

Table 2. Poisson regression for evaluation of surgeries annually.

	p-value
Year of surgery	<0.001
Type of surgery	
Mastectomy (No-BR + IBR)	0.907
BCS	1.000
Reconstruction	
No-BR	1.000
Implant	0.950
Myocutaneous flap	0.924

No-BR: Total mastectomy without reconstruction; IBR: Mastectomy and immediate reconstruction; BCS: Breast-conserving surgery; p: Value of significance after Poisson regression.

Table 1. Evolution of surgeries (BCS, No-BR and IBR) annually between 2012 and 2019 (number of surgeries and proportion).

Year	No-BR(n)	%	IBR (n)	%	BCS (n)	%	Total
2012	44	51.76	19	22.35	22	25.88	85
2013	49	50.00	20	20.41	29	29.59	98
2014	48	55.81	22	25.58	16	18.60	86
2015	56	47.46	30	25.42	32	27.12	118
2016	52	53.61	18	18.56	27	27.84	97
2017	40	35.40	28	24.78	45	39.82	113
2018	39	38.24	22	21.57	41	40.20	102
2019	43	40.57	22	20.75	41	38.68	106
Total	371	-	181	-	253	-	805
p-Value		p=0.299		p=0.663		p=0.003	

In line with our findings, a recent Brazilian study reported an increase in breast reconstructions in Brazil between 2008 and 2014, reaching 29%, consistent with the results of this study¹⁸. In fact, the availability of immediate reconstruction has been related to mastectomies in some studies, especially in cases of unilateral cancer and contralateral risk-reducing mastectomy. In this sense, a retrospective study using the NCDB demonstrated that the mastectomy rate, specifically contralateral mastectomy, increased by 7% for each percentage point increase in reconstruction¹¹. The rates of immediate reconstruction, on the other hand, were similar to those observed in some countries. For example, a European study evaluated 2,315 patients with early breast cancer between 2002 and 2016, in which the authors showed that the rate of immediate reconstruction was 34.3%¹⁹. Other factors, including barriers to access to breast reconstruction, were highlighted through a Canadian systematic review, which identified that, in rural areas, the high costs of the procedure, insufficient reimbursement by insurance companies, non-acceptance of the procedure by the patient, tumor characteristics and lack of patient awareness about reconstruction were related to non-performance of reconstruction²⁰.

The false impression that mastectomy could be a safer treatment may also explain the increase in mastectomies²¹. A study showed that only 38.1% of patients with unilateral breast cancer knew that contralateral prophylactic surgery had no effect on survival²². In fact, many patients opt for mastectomy, whether unilateral or bilateral, even after the proven observation of a reduction in ipsilateral recurrences or new contralateral tumors in patients undergoing conservative surgeries, observed over the years, possibly because of the advent of systemic therapy²³. Another reason patients may opt for bilateral mastectomy, especially in cases of unilateral breast cancer with indication for mastectomy and reconstruction with implants, is the possibility of better symmetry: a study conducted at Memorial Sloan Kettering Cancer Center using Breast-Q in 3,489 breasts of women who opted for mastectomy with bilateral reconstruction with implants showed that the latter procedure had the best aesthetic result over a 12-year follow-up²².

The advent of genetic counseling and multigene testing, which evaluate genes that are related to a greater hereditary predisposition for patients with unilateral breast cancer to develop recurrence or contralateral breast cancer, may also influence mastectomy rates. Recent studies conducted with thousands of patients have identified genes with a high predisposition for the incidence of breast cancer, whether first or second primary, after treatment for breast cancer, especially BRCA1 and BRCA2²⁴⁻²⁷. Some studies have also demonstrated a significant reduction in new tumors after bilateral mastectomy, as well as an impact on mortality in women with BRCA mutations and unilateral breast cancer^{28,29}. However, in many places around the world, access to financial resources for health care is a significant impediment, as is the case for patients in the Brazilian public health system, who were the patients treated in this analysis. Furthermore, several regions

of Brazil do not have the availability of cancer genetic specialists even for women with access to the private supplementary health network³⁰. This issue, however, does not seem to be a problem only in countries with limited resources, with significant disparities occurring in developed countries in the management of breast cancer patients. An observational study presented at the 2023 American Society of Clinical Oncology (ASCO) meeting involving more than 1 million women who were diagnosed with any type of cancer between 2013 and 2019 assessed the prevalence of germline testing in American patients, with only 6.8% of them undergoing genetic testing, with tests performed below expectations even in cases where the test is recommended in guidelines³¹. Furthermore, patients of Asian, Hispanic or Black ethnicity had a lower proportion of tests performed³¹.

Another fact that stands out in our study is the high volume of mastectomies during this recent period. This is possibly because diagnosis still occurs in stages II and III, with great frequency, as a consequence of inadequate screening in patients in the public system. A Brazilian study conducted with 4,912 patients diagnosed with breast cancer in Brazil in 28 institutions in 2001 and 2006 demonstrated that approximately 75% of cases were stages II and III, higher than that observed in high-income countries, and approximately 80% of these cases were treated in the public system³². Despite this, there was paradoxically a trend towards an increase in BCS in our analysis during the study period, which may hypothetically reflect the advent of neoadjuvant therapies and a greater possibility of BCS in initially ineligible patients. In Brazil, during the study period, patients in the public system had access to adequate systemic treatment, including anti-HER2 therapy since 2013, although pertuzumab is not yet available for non-metastatic disease. Several studies have, in fact, demonstrated an increase in the rate of breast conservation and oncological safety in performing BCS in patients who were ineligible at the time of diagnosis^{33,34}. The use of oncoplastic techniques frequently used in our institution may also have contributed to this trend, possibly favoring an increase in the rate of breast conservation even in extreme cases (tumors larger than 5 cm, for example), being an alternative to mastectomy³⁵.

Our study had some limitations, including the fact that it was a retrospective study at a single institution that treats patients in the public system, with limited resources. Therefore, our results should not be extrapolated in a generalized manner, especially in relation to institutions that treat patients with access to the private and supplementary health network in Brazil. For these patients, there is a tendency for bilateral mastectomies to increase³⁶, similar to what happens in some high-income countries. The analysis of a single institution may also not reflect other institutions, even public ones, or other cities or regions. Finally, factors specific to the surgical team, such as the incorporation of oncoplastic techniques and clinical-pathological characteristics of the disease, such as staging and younger age, for example, may have influenced decision-making and were not evaluated in this study³⁷.

To the best of our knowledge, this is the first study to assess the contemporary trend of breast surgeries (mastectomy and BCS) for the treatment of early breast cancer in an institution with limited resources — in this case, in Brazil. These data reveal the importance of discussion, as well as public policies for the incorporation of new technologies, including genetic testing. As in other places in the world, access and disparity are relevant problems in Brazil, whether for surgical treatment, including immediate reconstruction, or even for the incorporation of new drugs with high costs for systemic treatment.

CONCLUSIONS

We did not see an increase in mastectomy rates during the study period (2012–2019) in a public institution with limited resources. However, a trend of increasing BCS rates was demonstrated over the years at HGF. Although these results may not be generalizable, suggest that rising mastectomy trends may not be consistent across countries, especially with differences in access to health care systems, and warrant further studies and discussion on this scenario.

AUTHORS' CONTRIBUTION

FPC: Conceptualization, Data curation, Formal Analysis, Funding acquisition, Investigation, Methodology, Project administration, Resources, Software, Supervision, Validation, Visualization, Writing – original draft, Writing – review & editing. TMGP: Conceptualization, Data curation, Formal Analysis, Funding acquisition, Investigation, Methodology, Project administration, Resources, Software, Supervision, Validation, Visualization, Writing – original draft, Writing – review & editing. FPZ: Conceptualization, Validation, Visualization, Writing – original draft, Writing – review & editing. ECM: Conceptualization, Validation, Visualization, Writing – original draft, Writing – review & editing. AM: Conceptualization, Validation, Visualization, Writing – original draft, Writing – review & editing. MA: Conceptualization, Validation, Visualization, Writing – original draft, Writing – review & editing. FPB: Conceptualization, Validation, Visualization, Writing – original draft, Writing – review & editing. GGN: Conceptualization, Validation, Visualization, Writing – original draft, Writing – review & editing. ALF: Conceptualization, Validation, Visualization, Writing – original draft, Writing – review & editing. RFJ: Conceptualization, Validation, Visualization, Writing – original draft, Writing – review & editing.








REFERENCES

1. Siegel RL, Miller KD, Wagle NS, Jemal A. Cancer statistics, 2023. *CA Cancer J Clin.* 2023;73(1):17-48. <https://doi.org/10.3322/caac.21763>
2. Bray F, Laversanne M, Sung H, Ferlay J, Siegel RL, Soerjomataram I, et al. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2024;74(3):229-63. <https://doi.org/10.3322/caac.21834>
3. Santos M de O, Lima FC da S de, Martins LFL, Oliveira JFP, Almeida LM de, Cancela M de C. Estimativa de incidência de câncer no Brasil, 2023-2025. *Rev Bras Cancerol.* 2023;69(1):e-213700. <https://doi.org/10.32635/2176-9745.RBC.2023v69n1.3700>
4. Balic M, Thomssen C, Gnant M, Harbeck N. St. Gallen/Vienna 2023: Optimization of treatment for patients with primary breast cancer - a brief summary of the consensus discussion. *Breast Care.* 2023;18(3):213-22. <https://doi.org/10.1159/000530584>
5. Gradishar WJ, Moran MS, Abraham J, Abramson V, Aft R, Agnese D, et al. NCCN Guidelines® Insights: Breast Cancer, Version 4.2023. *J Natl Compr Canc Netw.* 2023;21(6):594-608. <https://doi.org/10.6004/jnccn.2023.0031>
6. Halsted WS. I. The results of radical operations for the cure of carcinoma of the breast. *Ann Surg.* 1907;46(1):1-19. <https://doi.org/10.1097/00000658-190707000-00001>
7. Fisher B, Anderson S. Conservative surgery for the management of invasive and noninvasive carcinoma of the breast: NSABP trials. National Surgical Adjuvant Breast and Bowel Project. *World J Surg.* 1994;18(1):63-9. <https://doi.org/10.1007/BF00348193>
8. Veronesi U, Zucali R, Luini A. Local control and survival in early breast cancer: the Milan trial. *Int J Radiat Oncol Biol Phys.* 1986;12(5):717-20. [https://doi.org/10.1016/0360-3016\(86\)90027-1](https://doi.org/10.1016/0360-3016(86)90027-1)
9. Neuburger J, Macneill F, Jeevan R, van der Meulen JH, Cromwell DA. Trends in the use of bilateral mastectomy in England from 2002 to 2011: retrospective analysis of hospital episode statistics. *BMJ Open.* 2013;3(8):e003179. <https://doi.org/10.1136/bmjopen-2013-003179>
10. Metcalfe K, Gershman S, Ghadirian P, Lynch HT, Snyder C, Tung N, et al. Contralateral mastectomy and survival after breast cancer in carriers of BRCA1 and BRCA2 mutations: retrospective analysis. *BMJ.* 2014;348:g226. <https://doi.org/10.1136/bmj.g226>
11. Albornoz CR, Matros E, Lee CN, Hudis CA, Pusic AL, Elkin E, et al. Bilateral mastectomy versus breast-conserving surgery for early-stage breast cancer: the role of breast reconstruction. *Plast Reconstr Surg.* 2015;135(6):1518-26. <https://doi.org/10.1097/PRS.0000000000001276>
12. Tuttle TM, Habermann EB, Grund EH, Morris TJ, Virnig BA. Increasing use of contralateral prophylactic mastectomy for breast cancer patients: a trend toward more aggressive surgical treatment. *J Clin Oncol.* 2007;25(33):5203-9. <https://doi.org/10.1200/JCO.2007.12.3141>
13. Habermann EB, Abbott A, Parsons HM, Virnig BA, Al-Refaie WB, Tuttle TM. Are mastectomy rates really increasing in the United States? *J Clin Oncol.* 2010;28(21):3437-41. <https://doi.org/10.1200/JCO.2009.27.6774>
14. Kummerow KL, Du L, Penson DF, Shyr Y, Hooks MA. Nationwide trends in mastectomy for early-stage breast cancer. *JAMA Surg.* 2015;150(1):9-16. <https://doi.org/10.1001/jamasurg.2014.2895>

15. Ahern TP, Larsson H, Garne JP, Cronin-Fenton DP, Sørensen HT, Lash TL. Trends in breast-conserving surgery in Denmark, 1982-2002. *Eur J Epidemiol.* 2008;23(2):109-14. <https://doi.org/10.1007/s10654-007-9207-1>
16. Gomez SL, Lichtensztajn D, Kurian AW, Telli ML, Chang ET, Keegan TH, et al. Increasing mastectomy rates for early-stage breast cancer? Population-based trends from California. *J Clin Oncol.* 2010;28(10):e155-7; author reply e158. <https://doi.org/10.1200/JCO.2009.26.1032>
17. Brasil. Legislação Informatizada - Lei nº 12.802, de 24 de abril de 2013 - Publicação Original [Internet]. Brasil; 2013 [accessed July 29, 2023]. Available at: <https://www2.camara.leg.br/legin/fed/lei/2013/lei-12802-24-abril-2013-775848-publicacaooriginal-139619-pl.html>
18. Freitas-Júnior R, Gagliato DM, Moura Filho JWC, Gouveia PA, Rahal RMS, Paulinelli RR, et al. Trends in breast cancer surgery at Brazil's public health system. *J Surg Oncol.* 2017;115(5):544-9. <https://doi.org/10.1002/jso.24572>
19. Fancellu A, Sanna V, Cottu P, Feo CF, Scanu AM, Farina G, et al. Mastectomy patterns, but not rates, are changing in the treatment of early breast cancer. Experience of a single European institution on 2315 consecutive patients. *Breast.* 2018;39:1-7. <https://doi.org/10.1016/j.breast.2018.02.003>
20. Retrouvey H, Solaja O, Gagliardi AR, Webster F, Zhong T. Barriers of access to breast reconstruction: a systematic review. *Plast Reconstr Surg.* 2019;143(3):465e-476e. <https://doi.org/10.1097/PRS.0000000000005313>
21. Huang J, Chagpar A. Factors associated with decision to undergo contralateral prophylactic mastectomy versus unilateral mastectomy. *Am J Surg.* 2019;218(1):170-4. <https://doi.org/10.1016/j.amjsurg.2018.11.040>
22. Jagi R, Hawley ST, Griffith KA, Janz NK, Kurian AW, Ward KC, et al. Contralateral prophylactic mastectomy decisions in a population-based sample of patients with early-stage breast cancer. *JAMA Surg.* 2017;152(3):274-82. <https://doi.org/10.1001/jamasurg.2016.4749>
23. Nichols HB, Berrington de González A, Lacey JV Jr, Rosenberg PS, Anderson WF. Declining incidence of contralateral breast cancer in the United States from 1975 to 2006. *J Clin Oncol.* 2011;29(12):1564-9. <https://doi.org/10.1200/JCO.2010.32.7395>
24. Breast Cancer Association Consortium. Breast Cancer Risk Genes - Association Analysis in More than 113,000 Women. *N Engl J Med.* 2021;384(5):428-39. <https://doi.org/10.1056/NEJMoa1913948>
25. Hu C, Hart SN, Gnanaolivu R, Huang H, Lee KY, Na J, et al. A population-based study of genes previously implicated in breast cancer. *N Engl J Med.* 2021;384(5):440-51. <https://doi.org/10.1056/NEJMoa2005936>
26. Yadav S, Boddicker NJ, Na J, Polley EC, Hu C, Hart SN, et al. Contralateral breast cancer risk among carriers of germline pathogenic variants in *ATM*, *BRCA1*, *BRCA2*, *CHEK2*, and *PALB2*. *J Clin Oncol.* 2023;41(9):1703-13. <https://doi.org/10.1200/JCO.22.01239>
27. Morra A, Mavaddat N, Muranen TA, Ahearn TU, Allen J, Andrulis IL, et al. The impact of coding germline variants on contralateral breast cancer risk and survival. *Am J Hum Genet.* 2023;110(3):475-86. <https://doi.org/10.1016/j.ajhg.2023.02.003>
28. Li X, You R, Wang X, Liu C, Xu Z, Zhou J, et al. Effectiveness of prophylactic surgeries in BRCA1 or BRCA2 mutation carriers: a meta-analysis and systematic review. *Clin Cancer Res.* 2016;22(15):3971-81. <https://doi.org/10.1158/1078-0432.CCR-15-1465>
29. Valachis A, Nearchou AD, Lind P. Surgical management of breast cancer in BRCA-mutation carriers: a systematic review and meta-analysis. *Breast Cancer Res Treat.* 2014;144(3):443-55. <https://doi.org/10.1007/s10549-014-2890-1>
30. Buzaid AC, Achatz MI, Amorim GLS, Barrios CH, Carvalho FM, Cavalcante FP, et al. Challenges in the journey of breast cancer patients in Brazil. Desafios na jornada de pacientes com câncer de mama no Brasil. *Braz J Oncol.* 2020;16:e20200021. <https://doi.org/10.5935/2526-8732.20200021>
31. Kurian AW, Abrahamse P, Furgal A, Ward KC, Hamilton AS, Hodan R, et al. Germline genetic testing after cancer diagnosis. *JAMA.* 2023;330(1):43-51. <https://doi.org/10.1001/jama.2023.9526>
32. Simon SD, Bines J, Werutsky G, Nunes JS, Pacheco FC, Segalla JG, et al. Characteristics and prognosis of stage I-III breast cancer subtypes in Brazil: The AMAZONA retrospective cohort study. *Breast.* 2019;44:113-9. <https://doi.org/10.1016/j.breast.2019.01.008>
33. Early Breast Cancer Trialists' Collaborative Group (EBCTCG). Long-term outcomes for neoadjuvant versus adjuvant chemotherapy in early breast cancer: meta-analysis of individual patient data from ten randomised trials. *Lancet Oncol.* 2018;19(1):27-39. [https://doi.org/10.1016/s1470-2045\(17\)30777-5](https://doi.org/10.1016/s1470-2045(17)30777-5)
34. Mamtani A, Sevilimedu V, Le T, Morrow M, Barrio AV. Is local recurrence higher among patients who downstage to breast conservation after neoadjuvant chemotherapy? *Cancer.* 2022;128(3):471-8. <https://doi.org/10.1002/cncr.33929>
35. Paulinelli RR, Ribeiro LFJ, Santos TD, Caires EMS, Pontes MGM, Faria BM, et al. Oncoplastic mammoplasty with disguised geometric compensation. *Surg Oncol.* 2021;39:101660. <https://doi.org/10.1016/j.suronc.2021.101660>
36. Cavalcante FP, Silva PG de B, Lima MVA. Trends in bilateral mastectomy for cases of unilateral breast cancer in a Brazilian institute over a 10-year period. *Mastology.* 2021;31:1-6. <https://doi.org/10.29289/2594539420210030>
37. Kurian AW, Ward KC, Abrahamse P, Hamilton AS, Deapen D, Morrow M, et al. Association of germline genetic testing results with locoregional and systemic therapy in patients with breast cancer. *JAMA Oncol.* 2020;6(4):e196400. <https://doi.org/10.1001/jamaoncol.2019.6400>



Vacuum-assisted biopsy and excision of breast lesions: review and current indications

Letícia Martins de Araújo Campos Linhares^{1*} , José Tadeu Campos de Avelar¹ ,
Gabriela Ramos Alves¹ , Thaís Paiva Moraes¹ , Elizabeth Asuncion Sanchez de Ayub¹ ,
Jairo Luis Coelho Junior¹ , Renata Capanema Saliba Franco¹ 

ABSTRACT

Vacuum-assisted biopsy is an advance in breast diagnostics because it is a less invasive and more practical approach than conventional surgery, capable of diagnosing and treating certain lesions. Despite the still limited availability of the method, related to its cost and the number of professionals qualified to perform it, the potential of vacuum-assisted biopsy and excision in the practice of mastologists is unquestionable. The attending physician is expected to understand the methods, as well as the indications pertinent to them. Recognizing the impossibility of exhausting the subject, the objective of this study is to conduct a narrative review, summarizing the indications for vacuum-assisted breast biopsy and excision currently, according to the available scientific evidence.

KEYWORDS: biopsy; large-core needle; image-guided biopsy.

INTRODUCTION

The evolution of breast biopsy techniques began in 1960, with the advent of fine-needle aspiration biopsy (FNA), widely used until the 1990s, enabling cytopathological analysis of the lesion and the possible distinction between malignancy and benignity. Incorporated into the routine diagnosis of breast lesions, together with clinical evaluation and imaging examination, FNAB had a non-diagnosis rate of 40%¹. Given the diagnostic limitations of the method, a new puncture technology was developed: core needle biopsy, also known as core biopsy. This new technique consists of an automatic device, with a large-caliber needle, capable of biopsying an entire fragment of the lesion and thus providing histological study, providing greater detail on benignity or malignancy, as well as the diagnosis of *in situ* or invasive lesions. In the mid-1990s, core biopsy became the standard intervention for diagnosing breast lesions with superiority based on a successful biopsy rate of 99% of cases, compared to 60-75% for FNA², and a sensitivity of 80-93%, compared to 65-82% for FNA³.

Approved by the Food and Drug Administration (FDA) in April 1995, vacuum-assisted biopsy, also known by the abbreviation

VAB, has been used in medical practice in Brazil ever since, and has been progressively incorporated into the list of mandatory minimum coverage for health insurance plans in the country by the National Supplementary Health Agency (ANS). With the latest update in February 2021, ANS began granting access to VAB via ultrasound, stereotactics and magnetic resonance imaging in certain clinical contexts^{4,5}.

VAB is a safe, fast and effective technique for excising breast fragments, whose equipment basically consists of a biopsy needle coupled to a rotational cutter, a suction chamber and a device capable of creating a vacuum. The procedure is performed through a single incision in the skin, with the biopsy needle being inserted only once, since the material collected in each cycle is collected by suction into the reservoir chamber⁶. The effectiveness of VAB is based on its greater technical precision and broader sampling compared to core biopsy, since VAB needles range from 8G to 14G, with the capacity to collect between 40 mg and 300 mg of tissue per fragment, while core biopsy has an average collection of 17 mg of tissue per fragment⁷.

¹Mater Dei Rede de Saúde – Belo Horizonte (MG), Brazil.

*Corresponding author: leticialinhares_@hotmail.com

Conflict of interests: nothing to declare. Funding: none.

Received on: 01/15/2024. Accepted on: 02/06/2025.

Explored in the context of breast biopsy for years, VAB has become a consolidated and improved technique, which culminated in the idea of also making it an option for therapeutic proposal for some types of breast lesions. With the approval of the FDA in 2002, vacuum-assisted excision, recognized by the abbreviation VAE, proposes the complete excision of breast lesions using needles larger in diameter than those used in VAB, but maintaining its outpatient nature, under local anesthesia to avoid surgical excision, whose complexity and costs are significantly greater^{6,7}.

METHODS

This study aimed to address the topic of vacuum-assisted biopsy and excision of breast lesions, with an emphasis on current indications for the procedure. The search was conducted in the PubMed, Medline, Embase, and Cochrane databases, using the following search terms: “Biopsy”, “Large-Core Needle”, and “Image-guided biopsy”. Articles published between 2006 and 2023, in English and Portuguese that directly addressed breast biopsy and vacuum-assisted excision of breast lesions were included. The period from 2006 to 2023 was chosen to cover the development and consolidation of the techniques, considering from the approval of vacuum-assisted excision by the Food and Drug Administration (FDA) in 2002 to the most recent studies available. The inclusion criteria considered original articles, reviews, clinical practice guidelines, and relevant studies related to the topic. Duplicate publications, articles that dealt with techniques unrelated to the topic, and those unavailable in full text were excluded. The articles identified in the search were screened in two stages: initially, through the analysis of titles and abstracts; then, by reading them in full to determine if the inclusion criteria were met. After applying the inclusion and exclusion criteria, 20 articles were selected for analysis, considering their relevance and alignment with the objective of this study. In addition to the articles, the normative resolutions of the National Supplementary Health Agency (ANS) were analyzed, especially with regard to the coverage of procedures by supplementary health in Brazil, to complement the analysis on clinical applicability. Because this is a narrative review, the absence of systematic criteria and the subjective choice of articles may limit the reproducibility of the study and introduce biases. However, efforts were made to ensure the inclusion of relevant and updated publications, with the aim of providing a comprehensive and informative view on the topic.

RESULTS AND DISCUSSION

Core needle biopsy vs. vacuum-assisted biopsy

Core needle biopsy is the routine diagnostic method because it provides a quality diagnosis, is easy to perform, is minimally

invasive and has a lower cost than surgical excision. In this scenario, both core biopsy and vacuum-assisted aspiration are feasible options, although the clinical decision to choose one or the other is sometimes not simple and carries doubts about the possibility of false negatives and diagnostic underestimation^{8,9}. Compared to core biopsy, vacuum-assisted biopsy offers larger samples, lower false negative rates, lower rebiopsy rates and lower diagnostic underestimation^{7,10}. In this scenario, the indications for vacuum-assisted aspiration have grown and have been reaffirmed by studies and positive statistical numbers.

Regarding concerns about false-negative results and diagnostic underestimation, the diagnostic upgrade rate is the parameter for comparison between methods. The upgrade rate is defined by the presence of malignancy after surgical excision of the lesion previously biopsied or during its clinical-imaging follow-up¹¹.

Regarding the use of VAB in the diagnostic investigation of breast lesions with malignant potential, comparative studies show that, in cases of atypical ductal hyperplasia, the diagnostic upgrade rate for ductal carcinoma *in situ* was reduced from 50% to 20%; but in cases of ductal carcinoma *in situ*, the diagnostic upgrade rate for invasive ductal carcinoma was 30% to 10%, when comparing the results of core biopsy and VAB, respectively¹².

Other studies analyzing the upgrade rate in VAB show variable trends towards malignancy in specimen analysis or in clinical follow-up. A retrospective study showed an upgrade rate of 0.4% and a confidence interval between 0.1% and 2.1% when analyzing lesions of uncertain behavior without atypia, while the upgrade rate for lesions of uncertain behavior with atypia, such as flat epithelial atypia, atypical ductal or lobular hyperplasia, and lobular carcinoma *in situ*, was 4.7% with a confidence interval between 2.9% and 7.5%. The same study showed that the upgrade rate is significantly increased when two or more lesions coexist in the same biopsied area¹¹.

VAB has a higher cost per biopsy needle when compared to core biopsy needle; therefore, for lesions in which VAB is not the first method of choice, cost-effectiveness should be taken into consideration and core biopsy should be considered. On the other hand, in lesions for which VAB is used, the higher cost of the needle is outweighed by the benefits of diagnostic accuracy. Grady et al.¹³ refers to a study conducted in 2015 reviewing data from the American Society of Breast Surgeons between 2001 and 2014, containing data from 31,451 patients, in which information on biopsy, rebiopsy, instrument used to perform the biopsy and the cost per breast cancer diagnosis in each situation were evaluated, using a linear mathematical model. The study resulted in an average cost per case diagnosed by core biopsy of \$4,346 (4,327–4,366) and of \$3,742 (3,732–3,752) to \$4,779 (4,750–4,809) per VAB, with variation related to the brand of the device used. The study concluded that VAB would be more cost-effective, provided that the best-performing devices were chosen.

Vacuum-assisted biopsy vs. Vacuum-assisted excision

In addition to its investigative role, VAB also has an established therapeutic role. Unlike vacuum biopsy, which has a purely diagnostic purpose and does not require complete removal of the lesion, VAE aims to replace diagnostic surgical biopsy by removing the lesion in its entirety. The VAE technique stipulates that the aspiration be orthogonal, rather than oblique, to the lesion, and it also stipulates complete removal of the lesion and its periphery to ensure greater diagnostic accuracy¹⁴. To perform VAE, the needles used are larger in size, ranging from 7G to 10G, with the capacity to obtain large and varied amounts of tissue. There is no consensus on the size of the lesion to be approached, and according to Park et al.¹⁵, there is no limit to the size to be excised, which should be guided by the characteristics of the lesion, its location and the patient's particularities. At the end of the procedure, radiological verification showing the absence of the target lesion is mandatory, as well as subsequent assessment of imaging and histopathological compatibility.

Vacuum-assisted excision vs. conventional surgical approach

VAE is a subject of broad debate due to the high variability in the rates of upgrade to malignancy of lesions treated with the technique. The defense of vacuum excision with subsequent clinical follow-up is based on the personalization of treatment and advocates the de-escalation of invasive procedures associated with strategies to not underestimate the patient's subsequent specific risk of developing breast cancer, while opposition to vacuum aspiration points to the lack of standardized protocols and robust evidence as factors that prevent the widespread use of the method¹¹. Based on the BI-RADS (Breast Imaging Reporting and Data System) classification system, an upper safety limit can be inferred to be an underestimation rate of up to 2% after vacuum-assisted aspiration procedures, which would be equivalent to a lesion classified as BI-RADS 3, in terms of probability of malignancy. Therefore, breast lesions with upgrade rates $\leq 2\%$ after the procedure – a group generally composed of lesions without atypia – would be eligible for clinical follow-up in to avoid surgical approaches subsequent to biopsy.

In cases of lesions with atypia, carcinomas *in situ*, invasive carcinomas and lesions with clinical-radiological-histopathological discordance, conventional surgery is mandatory in the therapeutic management, since vacuum aspiration violates the basic principle of oncological surgery, which is the ability to perform a single-block excision of the lesion, in addition to the possibility of evaluating the margins of the specimen, which must be free of neoplasia⁸. However, there are studies aimed at proving the role of excision by vacuum aspiration even in these scenarios.

Valadares et al.¹⁶ published in 2023 a retrospective study evaluating data from 116 patients who underwent vacuum aspiration

of BI-RADS 4, BI-RADS 5 breast lesions or lesions with a diagnosis of uncertain malignant potential (B3) in a previous core biopsy, with anatomopathological results of *in situ* or invasive breast carcinoma of the aspirated specimens. These patients subsequently underwent conventional surgery to evaluate the existence of residual disease, and the result of the study was promising, mainly for low or moderate grade pT1a and pT1b tumors, which provides data to support the selection of criteria for future prospective trials on the subject. Regarding data on cost and effectiveness, Whitworth et al.¹⁷ conducted a retrospective cohort study, analyzing the amounts spent on the treatment of benign lesions and high-risk lesions treated by vacuum aspiration compared to similar lesions treated surgically. The study showed that VAE, in both the scenario of benign lesions and in the scenario of high-risk lesions, has a cost approximately 60% lower than the surgical approach, without compromising the quality of treatment.

Indications for vacuum-assisted biopsy

VAB has proven to be an accurate technique for treating suspicious breast calcifications, as shown in Figure 1, significantly reducing the need for rebiopsy (8), since this is a small breast alteration that can be seen primarily only on mammography, making core biopsy even more technically difficult and with less favorable success rates. The same logic applies to tiny, non-palpable lesions and intracystic lesions, for which core biopsy needle shots are less likely to be accurate in relation to the lesion and, therefore, have lower diagnostic sensitivity, making VAB the best option in this scenario^{1,5,6}. With advances in imaging in the context of the breast, magnetic resonance imaging has become an important imaging method for detecting lesions suspected of malignancy. With an emphasis on lesions seen only by this method, areas of nodular or non-nodular enhancement, with no corresponding second-look ultrasound directed at such areas, have become lesions with indication for magnetic resonance-guided VAB¹⁸. Finally, some individual details of the cases make VAB




Figure 1. (A) Suspicious calcifications seen on mammography (red rim); (B) Image showing biopsy needle positioned below lesion; (C) Mammography image of biopsied tissue containing calcifications.

the method of choice, since, as it does not require triggering or advancement, it is safer when investigating nodules close to the chest, close to the skin, the nipple or implants⁶.

Indications for vacuum-assisted excision

Regarding nodules that show growth, such as phyllodes tumor, studies demonstrate that VAE has a recurrence rate of 5% to 17% when evaluating lesions up to 3.3 cm, with lower recurrence rates in lesions less than 1.5 cm, which makes the procedure an alternative to surgery for benign tumors¹⁹. The method can be used to treat fibroadenoma, the most common benign breast tumor, with good results^{10,12}, and is also a potential treatment option for minor gynecomastia, for which there is no need for surgical reduction of the skin or the nipple-areola complex²⁰. The indication for excision by vacuum aspiration extends to other breast lesions, such as radial scar and flat epithelial atypia, as long as they do not present atypia¹⁴. VAE is still the treatment used for intraductal papillary lesions, as shown in Figure 2, with studies showing resolution in 97% of cases, in addition to preserving the viability of the function and sensitivity of the nipple-areola complex, which is often compromised after a conventional surgical approach⁸. However, it is important to emphasize that surgery is still the rule in cases of lesions with atypia, carcinomas in situ and lesions with clinical-radiological-histopathological discordance.

Complications

The possible complications of core biopsy and VAB are similar and are represented by hematoma, pain, infection, pneumothorax and skin injury. Hematoma is the main complication after vacuum-assisted aspirations, and small-volume hematomas do not require treatment. Surgery is necessary

for hemostasis or debridement if there is suspicion of active bleeding or large-volume hematoma causing severe pain or secondary infection^{5,10}. A 2019 retrospective study²¹, including 4,776 patients undergoing VAB, identified complication in 6.7% biopsy cases. Of these, 96.2% were mild complications, which included hematoma that did not require treatment, mild pain, nausea, dizziness and itching or skin irritation, while only 3.8% were moderate complications, which included hematoma or bleeding requiring compressive treatment, vasovagal reaction or observation in the emergency room. Finally, in this study, no patient displayed serious complications, which would include post-biopsy infection, hematoma or bleeding requiring surgical treatment and death, arguing in favor of the safety of the method.

CONCLUSIONS

Vacuum-assisted aspiration has proven to be a highly valuable and potential procedure. Approved by the ANS in 2021, it is mandatory for health insurance plans to provide histopathological studies of non-palpable lesions, breast microcalcifications, intraductal or intracystic lesions suspected of being papillomas, nodular or non-nodular enhancements seen on magnetic resonance imaging, categorized as 4 or 5 in the BI-RADS classification, or in cases of nodules smaller than 2 cm, also categorized as 4 or 5 in the BI-RADS classification, in scenarios where doubts remain after core biopsy⁴. It is a comprehensive diagnostic and therapeutic method that because of its minimally invasive nature and association with high efficacy tends to contribute greatly to mastology^{5,10,11}. Future studies will be essential to guide, on the basis of robust evidence, protocols that optimize the application of the technique for each type of breast lesion, ensuring greater precision and clinical benefit.

AUTHORS' CONTRIBUTION

LMACL: Conceptualization, Data curation, Methodology, Project administration, Writing – original draft, Writing – review & editing. GRA: Data curation, Formal Analysis, Supervision, Validation, Writing – review & editing. TPM: Data curation, Formal Analysis, Supervision, Validation, Writing – review & editing. JLCJ: Formal Analysis, Supervision, Validation, Writing – review & editing. EASA: Formal Analysis, Supervision, Validation, Writing – review & editing. RCSF: Formal Analysis, Supervision, Validation, Writing – review & editing. ADS: Formal Analysis, Supervision, Validation, Writing – review & editing. WJAJ: Formal Analysis, Supervision, Validation, Writing – review & editing. FMOC: Formal Analysis, Supervision, Validation, Writing – review & editing. JTCA: Conceptualization, Formal Analysis, Methodology, Project administration, Supervision, Validation, Writing – review & editing.




Figure 2. (A) Mixed intraductal lesion on USG; (B) Metal clip inserted after the procedure at the biopsy site (red rim); (C) Macroscopic image of the biopsied tissue; (D) Mammography image after the procedure with confirmation of clip at the biopsy site.

REFERENCES

1. Bennett IC, Saboo A. The Evolving Role of Vacuum Assisted Biopsy of the Breast: A Progression from Fine-Needle Aspiration Biopsy. *World J Surg*. 2019;43(4):1054-61. <https://doi.org/10.1007/s00268-018-04892-x>
2. Willems SM, van Deurzen CHM, van Diest PJ. Diagnosis of breast lesions: fine-needle aspiration cytology or core needle biopsy? A review. *J Clin Pathol*. 2012;65(4):287-92.
3. Lieske B, Ravichandran D, Wright D. Role of fine-needle aspiration cytology and core biopsy in the preoperative diagnosis of screen-detected breast carcinoma. *Br J Cancer*. 2006;95(1):62-6.
4. Brasil. Ministério da Saúde. Agência Nacional de Saúde Suplementar. Resolução Normativa - RN nº 465, de 24 de fevereiro de 2021. Atualiza o Rol de Procedimentos e Eventos em Saúde que estabelece a cobertura assistencial obrigatória a ser garantida nos planos privados de assistência à saúde contratados a partir de 1º de janeiro de 1999 e naqueles adaptados conforme previsto no artigo 35 da Lei nº 9.656, de 3 de junho de 1998; fixa as diretrizes de atenção à saúde; e revoga a Resolução Normativa - RN nº 428, de 7 de novembro de 2017, a Resolução Normativa - RN nº 453, de 12 de março de 2020, a Resolução Normativa - RN nº 457, de 28 de maio de 2020 e a RN nº 460, de 13 de agosto de 2020.
5. Li SJ, Hao XP, Hua B, Wang JD, Fan ZM; Chinese Society of Breast Surgery. Clinical practice guidelines for ultrasound-guided vacuum-assisted breast biopsy: Chinese Society of Breast Surgery (CSBrS) practice guidelines 2021. *Chin Med J (Engl)*. 2021;134(12):1390-92. <https://doi.org/10.1097/CM9.0000000000001508>
6. Bennett IC. The changing role of vacuum-assisted biopsy of the breast: a new prototype of minimally invasive breast surgery. *Clin Breast Cancer*. 2017;17(5):323-25. <https://doi.org/10.1016/j.clbc.2017.03.001>
7. Park HL, Hong J. Vacuum-assisted breast biopsy for breast cancer. *Gland Surg*. 2014;3(2):120-27. <https://doi.org/10.3978/j.issn.2227-684X.2014.02.03>
8. Wang O, Zhang W, Chen S, Cao F, Chen L, Chen H. A Multicenter, Randomized, Controlled Study of the Breast Biopsy and Circumferential Excision System for Breast Lesions. *Clin Breast Cancer*. 2023;23(6):640-8. <https://doi.org/10.1016/j.clbc.2023.05.007>
9. Zhang Y, Li J, Mo M, Shen J, Ren H, Li S, et al. The comparison of efficacy and safety evaluation of vacuum-assisted Elite 10-G system and the traditional BARD 14-G core needle in breast diagnosis: an open-label, parallel, randomized controlled trial. *Int J Surg*. 2023;109(5):1180-7. <https://doi.org/10.1097/JS9.0000000000000257>
10. Dinç Elibol F, Dere Y, Belli AK, Elibol C, Dere Ö, Nazlı O. Both a biopsy method and a therapeutic procedure in BI-RADS 4A and 4B lesions: ultrasound-guided vacuum-assisted breast biopsy. *Turk J Surg*. 2020;36(1):65-71. <https://doi.org/10.5578/turksurg.4472>
11. Girardi V, Guaragni M, Ruzzenenti N, Palmieri F, Fogazzi G, Cozzi A, et al. B3 lesions at vacuum-assisted breast biopsy under ultrasound or mammography guidance: a single-center experience on 3634 consecutive biopsies. *Cancers (Basel)*. 2021;13(21):5443. <https://doi.org/10.3390/cancers13215443>
12. O'Flynn EAM, Wilson ARM, Michell MJ. Image-guided breast biopsy: state-of-the-art. *Clin Radiol*. 2010;65(4):259-70. <https://doi.org/10.1016/j.crad.2010.01.008>
13. Grady I, Vasquez T, Tawfik S, Grady S. Ultrasound-guided core-needle versus vacuum-assisted breast biopsy: a cost analysis based on the American Society of Breast Surgeons' Mastery of Breast Surgery Registry. *Ann Surg Oncol*. 2017;24(3):676-82. <https://doi.org/10.1245/s10434-016-5607-3>
14. Miranda BMM, Bitencourt AGV. Vacuum-assisted excision of breast lesions in surgical de-escalation: where are we? *Radiol Bras*. 2023;56(3):150-6. <https://doi.org/10.1590/0100-3984.2022.0078-en>
15. Park HL, Kwak JY, Jung H, Lee SH, Shim JY, Kim JY, et al. Is Mammotome excision feasible for benign breast mass bigger than 3 cm in greatest dimension? *J Korean Surg Soc*. 2006;70:25-9.
16. Valadares CN, Couto HL, Soares AN, Toppa PH, Ricardo BP, McIntosh SA, et al. Potential role of vacuum-assisted procedures in resecting breast cancers and highlighting selection criteria to support future trials. *Front Oncol*. 2023;13:1239574. <https://doi.org/10.3389/fonc.2023.1239574>
17. Whitworth P, Hogan A, Ferko N, Son D, Wang F, Xiong Y, et al. Reduced hospital costs for ultrasound-guided vacuum-assisted excision compared with open surgery in patients with benign breast masses and high-risk lesions. *J Breast Imaging*. 2020;2(5):452-61. <https://doi.org/10.1093/jbi/wbaa055>
18. Takahama N, Tozaki M, Ohgiya Y. Current status of MRI-guided vacuum-assisted breast biopsy in Japan. *Breast Cancer*. 2021;28(6):1188-94. <https://doi.org/10.1007/s12282-020-01107-x>
19. Shang QJ, Li N, Zhang MK, He Y, Liu G, Wang ZL. Ultrasound-guided vacuum-assisted excisional biopsy to treat benign phyllodes tumors. *Breast*. 2020;49:242-5. <https://doi.org/10.1016/j.breast.2019.12.008>
20. Couto HL, Valadares CN, Pellegrini Junior O, Oliveira TCF, El Bacha PMG, Ferreira SG. Minimally invasive treatment of gynecomastia by ultrasound-guided vacuum-assisted excision: report of a case series. *Mastology*. 2021;31:1-7.
21. Lin LLY, Gao Y, Lewin AA, Toth HK, Heller SL, Moy L. Overstated harms of breast cancer screening? A large outcomes analysis of complications associated with 9-gauge stereotactic vacuum-assisted breast biopsy. *AJR Am J Roentgenol*. 2019;212(4):925-32. <https://doi.org/10.2214/ajr.18.20421>

